

## References

- Abramson,S., Miller,R.G., and Phillips,R.A. (1977). Identification in Adult Bone-Marrow of Pluripotent and Restricted Stem-Cells of Myeloid and Lymphoid Systems. *Journal of Experimental Medicine* 145, 1567-1579.
- Adam,P.J., Regan,C.P., Hautmann,M.B., and Owens,G.K. (2000). Positive- and negative-acting Kruppel-like transcription factors bind a transforming growth factor beta control element required for expression of the smooth muscle cell differentiation marker SM22 alpha in vivo. *Journal of Biological Chemistry* 275, 37798-37806.
- Ai,W.D., Liu,Y., Langlois,M., and Wang,T.C. (2004). Kruppel-like factor 4 (KLF4) represses histidine decarboxylase gene expression through an upstream Sp1 site and downstream gastrin responsive elements. *Journal of Biological Chemistry* 279, 8684-8693.
- Akashi,K., Traver,D., Kondo,M., and Weissman,I.L. (1999). Lymphoid development from hematopoietic stem cells. *International Journal of Hematology* 69, 217-226.
- Akashi,K., Traver,D., Miyamoto,T., and Weissman,I.L. (2000). A clonogenic common myeloid progenitor that gives rise to all myeloid lineages. *Nature* 404, 193-197.
- Akashi,K., He,X., Chen,J., Iwasaki,H., Niu,C., Steenhard,B., Zhang,J.W., Haug,J., and Li,L.H. (2003). Transcriptional accessibility for genes of multiple tissues and hematopoietic lineages is hierarchically controlled during early hematopoiesis. *Blood* 101, 383-390.
- Alheim,K. and Bartfai,T. (1998). The interleukin-1 system: Receptors, ligands, and ICE in the brain and their involvement in the fever response. *Neuroimmunomodulation* 840, 51-58.
- Anderson,K.L., Smith,K.A., Pio,F., Torbett,B.E., and Maki,R.A. (1998). Neutrophils deficient in PU.1 do not terminally differentiate or become functionally competent. *Blood* 92, 1576-1585.
- Anderson,K.L., Smith,K.A., Conners,K., McKercher,S.R., Maki,R.A., and Torbett,B.E. (1998). Myeloid development is selectively disrupted in PU.1 null mice. *Blood* 91, 3702-3710.
- Angerer,N.D., Du,Y., Nalbant,D., and Williams,S.C. (1999). A short conserved motif is required for repressor domain function in the myeloid-specific transcription factor CCAAT enhancer-binding protein epsilon. *Journal of Biological Chemistry* 274, 4147-4154.
- Arredouani,M., Yang,Z.P., Ning,Y.Y., Qin,G.Z., Soininen,R., Tryggvason,K., and Kobzik,L. (2004). The scavenger receptor MARCO is required for lung defense against pneumococcal pneumonia and inhaled particles. *Journal of Experimental Medicine* 200, 267-272.
- Asada,M., Yamada,T., Ichijo,H., Delia,D., Miyazono,K., Fukumuro,K., and Mizutani,S. (1999). Apoptosis inhibitory activity of cytoplasmic p21(Cip1/WAF1) in monocytic differentiation. *Embo Journal* 18, 1223-1234.
- Asada,M., Ohmi,K., Delia,D., Enosawa,S., Suzuki,S., Yuo,A., Suzuki,H., and Mizutani,S. (2004). Brap2 functions as a cytoplasmic retention protein for p21 during monocyte differentiation. *Molecular and Cellular Biology* 24, 8236-8243.
- Asada,M., Mizutani,S., Ichijo,H., and Suzuki,H. (2004). Cytoplasmic p21 inhibits stress-activated MAP kinase pathway. *Journal of Pharmacological Sciences* 94, 136P.
- Aurigemma,R.E., Blair,D.G., and Ruscetti,S.K. (1992). Transactivation of Erythroid Transcription Factor Gata-1 by A Myb-Ets-Containing Retrovirus. *Journal of Virology* 66, 3056-3061.
- Barreda,D.R. and Belosevic,M. (2001). Transcriptional regulation of hemopoiesis. *Developmental and Comparative Immunology* 25, 763-789.
- Barreda,D.R., Hannington,P.C., and Belosevic,M. (2004). Regulation of myeloid development and function by colony stimulating factors. *Developmental and Comparative Immunology* 28, 509-554.
- Bavisotto,L., Kaushansky,K., Lin,N., and Hromas,R. (1991). Antisense Oligonucleotides from the Stage-Specific Myeloid Zinc Finger Gene Mzf-1 Inhibit Granulopoiesis Invitro. *Journal of Experimental Medicine* 174, 1097-1101.
- Beato,M., Chavez,S., and Truss,M. (1996). Transcriptional regulation by steroid hormones. *Steroids* 61, 240-251.
- Behm,C.A. and Ovington,K.S. (2000). The role of eosinophils in parasitic helminth infections: Insights from genetically modified mice. *Parasitology Today* 16, 202-209.

## References

- Besson,A., Assoian,R.K., and Roberts,J.M. (2004). Regulation of the cytoskeleton: an oncogenic function for CDK inhibitors? *Nat. Rev. Cancer* 4, 948-955.
- Bierie,B. and Moses,H.L. (2006). TGF beta: the molecular Jekyll and Hyde of cancer. *Nature Reviews Cancer* 6, 506-520.
- Bleasle,K., Lewis,A., and Raymon,H.K. (2003). Emerging treatments for asthma. *Expert Opin. Emerg. Drugs* 8, 71-81.
- Borzillo,G.V., Ashmun,R.A., and Sherr,C.J. (1990). Macrophage Lineage Switching of Murine Early Pre-B Lymphoid-Cells Expressing Transduced Fms Genes. *Molecular and Cellular Biology* 10, 2703-2714.
- Boyd,A.W. and Schrader,J.W. (1982). Derivation of Macrophage-Like Lines from the Pre-Lymphoma-B Abls 8.1 Using 5-Azacytidine. *Nature* 297, 691-693.
- Bradford,G.B., Williams,B., Rossi,R., and Bertoncello,I. (1997). Quiescence, cycling, and turnover in the primitive hematopoietic stem cell compartment. *Experimental Hematology* 25, 445-453.
- Braun,S.E., Mantel,C., Rosenthal,M., Cooper,S., Liu,L., Robertson,K.A., Hromas,R., and Broxmeyer,H.E. (1998). A positive effect of p21(cip1/waf1) in the colony formation from murine myeloid progenitor cells as assessed by retroviral-mediated gene transfer. *Blood Cells Molecules and Diseases* 24, 138-148.
- Brembeck,F.H. and Rustgi,A.K. (2000). The tissue-dependent keratin 19 gene transcription is regulated by GKLF/KLF4 and Sp1. *Journal of Biological Chemistry* 275, 28230-28239.
- Brett,J., Schmidt,A.M., Yan,S.D., Zou,Y.S., Weidman,E., Pinsky,D., Nowyngrod,R., Neeper,M., Przysiecki,C., Shaw,A., Migheli,A., and Stern,D. (1993). Survey of the Distribution of A Newly Characterized Receptor for Advanced Glycation End-Products in Tissues. *American Journal of Pathology* 143, 1699-1712.
- Cantor,A.B. and Orkin,S.H. (2001). Hematopoietic development: a balancing act. *Current Opinion in Genetics & Development* 11, 513-519.
- Chabot,B., Stephenson,D.A., Chapman,V.M., Besmer,P., and Bernstein,A. (1988). The Proto-Oncogene C-Kit Encoding A Transmembrane Tyrosine Kinase Receptor Maps to the Mouse W-Locus. *Nature* 335, 88-89.
- Chen,X.M., Johns,D.C., Geiman,D.E., Marban,E., Dang,D.T., Hamlin,G., Sun,R.G., and Yang,V.W. (2001). Kruppel-like factor 4 (Gut-enriched Kruppel-like factor) inhibits cell proliferation by blocking G(1)/S progression of the cell cycle. *Journal of Biological Chemistry* 276, 30423-30428.
- Chen,X.M., Whitney,E.M., Gao,S.Y., and Yang,V.W. (2003). Transcriptional profiling of Kruppel-like factor 4 reveals a function in cell cycle regulation and epithelial differentiation. *Journal of Molecular Biology* 326, 665-677.
- Chen,Z.Y., Shie,J.L., and Tseng,C.C. (2002). STAT1 is required for IFN-gamma-mediated gut-enriched Kruppel-like factor expression. *Experimental Cell Research* 281, 19-27.
- Chen,Z.Y., Shie,J.L., and Tseng,C.C. (2002). Gut-enriched Kruppel-like factor represses ornithine decarboxylase gene expression and functions as checkpoint regulator in colonic cancer cells. *Journal of Biological Chemistry* 277, 46831-46839.
- Cheng,T., Rodrigues,N., Shen,H.M., Yang,Y.G., Dombkowski,D., Sykes,M., and Scadden,D.T. (2000). Hematopoietic stem cell quiescence maintained by p21(cip1/waf1). *Science* 287, 1804-1808.
- Cheshier,S.P., Morrison,S.J., Liao,X.S., and Weissman,I.L. (1999). In vivo proliferation and cell cycle kinetics of long-term self-renewing hematopoietic stem cells. *Proceedings of the National Academy of Sciences of the United States of America* 96, 3120-3125.
- Chono,H., Yoshioka,H., Ueno,A., and Kato,I. (2001). Removal of inhibitory substances with recombinant fibronectin-CH-296 plates enhances the retroviral transduction efficiency of CD34(+)CD38(-) bone marrow cells. *Journal of Biochemistry* 130, 331-334.
- Coccia,E.M., Del Russo,N., Stellacci,E., Testa,U., Marziali,G., and Battistini,A. (1999). STAT1 activation during monocyte to macrophage maturation: role of adhesion molecules. *International Immunology* 11, 1075-1083.
- Coffman,R.L., Seymour,B.W.P., Hudak,S., Jackson,J., and Rennick,D. (1989). Antibody to Interleukin-5 Inhibits Helminth-Induced Eosinophilia in Mice. *Science* 245, 308-310.
- Colvin,G.A., Lambert,J.F., Abedi,M., Dooner,M.S., Demers,D., Moore,B.E., Greer,D., Aliotta,J.M., Pimentel,J., Cerny,J., Lum,L.G., and Quesenberry,P.J. (2004). Differentiation hotspots: the deterioration of hierarchy and stochasticity. *Blood Cells Molecules and Diseases* 32, 34-41.

## References

- Cross,M.A., Heyworth,C.M., Murrell,A.M., Bockamp,E.O., Dexter,T.M., and Green,A.R. (1994). Expression of Lineage Restricted Transcription Factors Precedes Lineage-Specific Differentiation in A Multipotent Hematopoietic Progenitor-Cell Line. *Oncogene* 9, 3013-3016.
- Dang,D.T., Bachman,K.E., Mahatan,C.S., Dang,L.H., Giardiello,F.M., and Yang,V.W. (2000). Decreased expression of the gut-enriched Kruppel-like factor gene in intestinal adenomas of multiple intestinal neoplasia mice and in colonic adenomas of familial adenomatous polyposis patients. *Febs Letters* 476, 203-207.
- Dang,D.T., Mahatan,C.S., Dang,L.H., Agboola,I.A., and Yang,V.W. (2001). Expression of the gut-enriched Kruppel-like factor (Kruppel-like factor 4) gene in the human colon cancer cell line RKO is dependent on CDX2. *Oncogene* 20, 4884-4890.
- Dang,D.T., Chen,X.M., Feng,J., Torbenson,M., Dang,L.H., and Yang,V.W. (2003). Overexpression of Kruppel-like factor 4 in the human colon cancer cell line RKO leads to reduced tumorigenicity. *Oncogene* 22, 3424-3430.
- Danielian,P.S., White,R., Lees,J.A., and Parker,M.G. (1992). Identification of A Conserved Region Required for Hormone Dependent Transcriptional Activation by Steroid-Hormone Receptors. *Embo Journal* 11, 1025-1033.
- Danielian,P.S., White,R., Hoare,S.A., Fawell,S.E., and Parker,M.G. (1993). Identification of Residues in the Estrogen-Receptor That Confer Differential Sensitivity to Estrogen and Hydroxytamoxifen. *Molecular Endocrinology* 7, 232-240.
- DeKoter,R.P., Walsh,J.C., and Singh,H. (1998). PU.1 regulates both cytokine-dependent proliferation and differentiation of granulocyte/macrophage progenitors. *Embo Journal* 17, 4456-4468.
- DeKoter,R.P. and Singh,H. (2000). Regulation of B lymphocyte and macrophage development by graded expression of PU.1. *Science* 288, 1439-1441.
- Demoulin,J.B. and Renauld,J.C. (1998). Interleukin 9 and its receptor: an overview of structure and function. *Int. Rev. Immunol.* 16, 345-364.
- Deng,C.X., Zhang,P.M., Harper,J.W., Elledge,S.J., and Leder,P. (1995). Mice Lacking P21(C/P1/Waf1) Undergo Normal Development, But Are Defective in G1 Checkpoint Control. *Cell* 82, 675-684.
- Dent,L.A., Strath,M., Mellor,A.L., and Sanderson,C.J. (1990). Eosinophilia in Transgenic Mice Expressing Interleukin-5. *Journal of Experimental Medicine* 172, 1425-1431.
- deSauvage,F.J., CarverMoore,K., Luoh,S.M., Ryan,A., Dowd,M., Eaton,D.L., and Moore,M.W. (1996). Physiological regulation of early and late stages of megakaryocytopoiesis by thrombopoietin. *Journal of Experimental Medicine* 183, 651-656.
- Di Cunto,F., Topley,G., Calautti,E., Hsiao,J., Ong,L., Seth,P.K., and Dotto,G.P. (1998). Dual pattern of p21 regulation and function in primary keratinocyte differentiation. *Journal of Investigative Dermatology* 110, 538.
- Driggers,P.H., Ennist,D.L., Gleason,S.L., Mak,W.H., Marks,M.S., Levi,B.Z., Flanagan,J.R., Appella,E., and Ozato,K. (1990). An Interferon Gamma-Regulated Protein That Binds the Interferon-Inducible Enhancer Element of Major Histocompatibility Complex Class-I Genes. *Proceedings of the National Academy of Sciences of the United States of America* 87, 3743-3747.
- Du,J., Stankiewicz,M.J., Liu,Y., Xi,Q., Schmitz,J.E., Lekstrom-Himes,J.A., and Ackerman,S.J. (2002). Novel combinatorial interactions of GATA-1, PU.1, and C/EBP epsilon isoforms regulate transcription of the gene encoding eosinophil granule major basic protein. *Journal of Biological Chemistry* 277, 43481-43494.
- Du,X.X. and Williams,D.A. (1997). Interleukin-11: Review of molecular, cell biology, and clinical use. *Blood* 89, 3897-3908.
- Duarte,R.F. and Franf,D.A. (2002). The synergy between stem cell factor (SCF) and granulocyte colony-stimulating factor (G-CSF): molecular basis and clinical relevance. *Leuk. Lymphoma* 43, 1179-1187.
- Dublet,B., Ruello,A., Pederzoli,M., Hajjar,E., Courbebaisse,M., Canteloup,S., Reuter,N., and Witko-Sarsat,V. (2005). Cleavage of p21/WAF1/CIP1 by proteinase 3 modulates differentiation of a monocytic cell line. *Journal of Biological Chemistry* 280, 30242-30253.
- Eldeiry,W.S., Tokino,T., Velculescu,V.E., Levy,D.B., Parsons,R., Trent,J.M., Lin,D., Mercer,W.E., Kinzler,K.W., and Vogelstein,B. (1993). Waf1, A Potential Mediator of P53 Tumor Suppression. *Cell* 75, 817-825.
- Enver,T. and Greaves,M. (1998). Loops, lineage, and leukemia. *Cell* 94, 9-12.

## References

- Fairbairn,L.J., Cowling,G.J., Reipert,B.M., and Dexter,T.M. (1993). Suppression of Apoptosis Allows Differentiation and Development of A Multipotent Hematopoietic-Cell Line in the Absence of Added Growth-Factors. *Cell* 74, 823-832.
- Farrar,M.A. and Schreiber,R.D. (1993). The Molecular Cell Biology of Interferon-Gamma and Its Receptor. *Annual Review of Immunology* 11, 571-611.
- Fehr,T., Schoedon,G., Odermatt,B., Holtschke,T., Schneemann,M., Bachmann,M.F., Mak,T.W., Horak,I., and Zinkernagel,R.M. (1997). Crucial role of interferon consensus sequence binding protein, but neither of interferon regulatory factor 1 nor of nitric oxide synthesis for protection against murine listeriosis. *Journal of Experimental Medicine* 185, 921-931.
- Feil,R., Brocard,J., Mascrez,B., LeMeur,M., Metzger,D., and Chambon,P. (1996). Ligand-activated site-specific recombination in mice. *Proceedings of the National Academy of Sciences of the United States of America* 93, 10887-10890.
- Feil,R., Wagner,J., Metzger,D., and Chambon,P. (1997). Regulation of Cre recombinase activity by mutated estrogen receptor ligand-binding domains. *Biochemical and Biophysical Research Communications* 237, 752-757.
- Feinberg,M.W., Cao,Z.X., Wara,A.K., Lebedeva,M.A., SenBanerjee,S., and Jain,M.K. (2005). Kruppel-like factor 4 is a mediator of proinflammatory signaling in macrophages. *Journal of Biological Chemistry* 280, 38247-38258.
- Feldmann,M. and Maini,R.N. (2001). Anti-TNF alpha therapy of rheumatoid arthritis: what have we learned? *Annual Review of Immunology* 19, 163-196.
- Filippi,M.D., Harris,C.E., Meller,J., Gu,Y., Zheng,Y., and Williams,D.A. (2004). Localization of Rac2 via the C terminus and aspartic acid 150 specifies superoxide generation, actin polarity and chemotaxis in neutrophils. *Nature Immunology* 5, 744-751.
- Finkelman,F.D., SheaDonohue,T., Goldhill,J., Sullivan,C.A., Morris,S.C., Madden,K.B., Gause,W.C., and Urban,J.F. (1997). Cytokine regulation of host defense against parasitic gastrointestinal nematodes: Lessons from studies with rodent models. *Annual Review of Immunology* 15, 505-533.
- Foster,K.W., Ren,S.R., Louro,I.D., Lobo-Ruppert,S.M., McKie-Bell,P., Grizzle,W., Hayes,M.R., Broker,T.R., Chow,L.T., and Ruppert,J.M. (1999). Oncogene expression cloning by retroviral transduction of adenovirus E1A-immortalized rat kidney RK3E cells: Transformation of a host with epithelial features by c-MYC and the zinc finger protein GKLF. *Cell Growth & Differentiation* 10, 423-434.
- Foster,K.W., Frost,A.R., McKie-Bell,P., Lin,C.Y., Engler,J.A., Grizzle,W.E., and Ruppert,J.M. (2000). Increase of GKLF messenger RNA and protein expression during progression of breast cancer. *Cancer Research* 60, 6488-6495.
- Foster,K.W., Liu,Z.L., Nail,C.D., Li,X.N., Fitzgerald,T.J., Bailey,S.K., Frost,A.R., Louro,I.D., Townes,T.M., Paterson,A.J., Kudlow,J.E., Lobo-Ruppert,S.M., and Ruppert,J.M. (2005). Induction of KLF4 in basal keratinocytes blocks the proliferation - differentiation switch and initiates squamous epithelial dysplasia. *Oncogene* 24, 1491-1500.
- Fujiwara,Y., Browne,C.P., Cunniff,K., Goff,S.C., and Orkin,S.H. (1996). Arrested development of embryonic red cell precursors in mouse embryos lacking transcription factor GATA-1. *Proceedings of the National Academy of Sciences of the United States of America* 93, 12355-12358.
- Fuller,J.F., McAdara,J., Yaron,Y., Sakaguchi,M., Fraser,J.K., and Gasson,J.C. (1999). Characterization of HOX gene expression during myelopoiesis: Role of HOX A5 in lineage commitment and maturation. *Blood* 93, 3391-3400.
- Gallily,R. and Feldman,M. (1967). Role of Macrophages in Induction of Antibody in X-Irradiated Animals. *Immunology* 12, 197-&.
- Gartel,A.L. and Tyner,A.L. (1999). Transcriptional regulation of the p21((WAF1/CIP1)) gene. *Experimental Cell Research* 246, 280-289.
- Geiman,D.E., Han,T.T., Johnson,J.M., and Yang,V.W. (2000). Transactivation and growth suppression by the gut-enriched Kruppel-like factor (Kruppel-like factor 4) are dependent on acidic amino acid residues and protein-protein interaction. *Nucleic Acids Research* 28, 1106-1113.
- Ghanem,L. and Steinman,R. (2005). A proapoptotic function of p21 in differentiating granulocytes. *Leukemia Research* 29, 1315-1323.

## References

- Giese,N.A., Gabriele,L., Doherty,T.M., Klinman,D.M., TadesseHeath,L., Contursi,C., Epstein,S.L., and Morse,H.C. (1997). Interferon (IFN) consensus sequence-binding protein, a transcription factor of the IFN regulatory factor family, regulates immune responses in vivo through control of interleukin 12 expression. *Journal of Experimental Medicine* 186, 1535-1546.
- Graf,T., McNagny,K., Brady,G., and Frampton,J. (1992). Chicken Erythroid-Cells Transformed by the Gag-Myb-Ets-Encoding E26 Leukemia-Virus Are Multipotent. *Cell* 70, 201-213.
- Greene,G.L. and Press,M.F. (1986). Structure and Dynamics of the Estrogen-Receptor. *Journal of Steroid Biochemistry and Molecular Biology* 24, 1-7.
- Hein,J., Kempf,V.A.J., Diebold,J., Bucheler,N., Preger,S., Horak,I., Sing,A., Kramer,U., and Autenrieth,I.B. (2000). Interferon consensus sequence binding protein confers resistance against Yersinia enterocolitica. *Infection and Immunity* 68, 1408-1417.
- Heyworth,C., Pearson,S., May,G., and Enver,T. (2002). Transcription factor-mediated lineage switching reveals plasticity in primary committed progenitor cells. *Embo Journal* 21, 3770-3781.
- Higaki,Y., Schullery,D., Kawata,Y., Shnyreva,M., Abrass,C., and Bomsztyk,K. (2002). Synergistic activation of the rat laminin gamma 1 chain promoter by the gut-enriched Kruppel-like factor (GKLF/KLF4) and Sp1. *Nucleic Acids Research* 30, 2270-2279.
- Hinnebusch,B.F., Siddique,A., Henderson,J.W., Malo,M.S., Zhang,W., Athaide,C.P., Abedrapo,M.A., Chen,X., Yang,V.W., and Hodin,R.A. (2004). Enterocyte differentiation marker intestinal alkaline phosphatase is a target gene of the gut-enriched Kruppel-like factor. *Am. J. Physiol Gastrointest. Liver Physiol* 286, G23-G30.
- Hirasawa,R., Shimizu,R., Takahashi,S., Osawa,M., Takayanagi,S., Kato,Y., Onodera,M., Minegishi,N., Yamamoto,M., Fukao,K., Taniguchi,H., Nakauchi,H., and Iwama,A. (2002). Essential and instructive roles of GATA factors in eosinophil development. *Journal of Experimental Medicine* 195, 1379-1386.
- Hodgson,G.S. and Bradley,T.R. (1979). Properties of Hematopoietic Stem-Cells Surviving 5-Fluorouracil Treatment - Evidence for A Pre-Cfu-S Cell. *Nature* 281, 381-382.
- Holmes,M., Turner,J., Fox,A., Chisholm,O., Crossley,M., and Chong,B. (1999). hFOG-2, a novel zinc finger protein, binds the co-repressor mCtBP2 and modulates GATA-mediated activation. *Journal of Biological Chemistry* 274, 23491-23498.
- Holtschke,T., Lohler,J., Kanno,Y., Fehr,T., Giese,N., Rosenbauer,F., Lou,J., Knobeloch,K.P., Gabriele,L., Waring,J.F., Bachmann,M.F., Zinkernagel,R.M., Morse,H.C., Ozato,K., and Horak,I. (1996). Immunodeficiency and chronic myelogenous leukemia-like syndrome in mice with a targeted mutation of the ICSBP gene. *Cell* 87, 307-317.
- Htun,H., Holth,L.T., Walker,D., Davie,J.R., and Hager,G.L. (1999). Direct visualization of the human estrogen receptor alpha reveals a role for ligand in the nuclear distribution of the receptor. *Molecular Biology of the Cell* 10, 471-486.
- Hu,H.M., Baer,M., Williams,S.C., Johnson,P.F., and Schwartz,R.C. (1998). Redundancy of C/EBP alpha, -beta, and -delta in supporting the lipopolysaccharide-induced transcription of IL-6 and monocyte chemoattractant protein-1. *Journal of Immunology* 160, 2334-2342.
- Hu,M., Krause,D., Greaves,M., Sharkis,S., Dexter,M., Heyworth,C., and Enver,T. (1997). Multilineage gene expression precedes commitment in the hemopoietic system. *Genes & Development* 11, 774-785.
- Iwasaki-Arai,J., Iwasaki,H., Miyamoto,T., Watanabe,S., and Akashi,K. (2003). Enforced granulocyte/macrophage colony-stimulating factor signals do not support lymphopoiesis, but instruct lymphoid to myelomonocytic lineage conversion. *Journal of Experimental Medicine* 197, 1311-1322.
- Iwasaki,H., Mizuno,S., Mayfield,R., Shigematsu,H., Arinobu,Y., Seed,B., Gurish,M.F., Takatsu,K., and Akashi,K. (2005). Identification of eosinophil lineage - committed progenitors in the murine bone marrow. *Journal of Experimental Medicine* 201, 1891-1897.
- Jaubert,J., Cheng,J., and Segre,J.A. (2003). Ectopic expression of Kruppel like factor 4 (Klf4) accelerates formation of the epidermal permeability barrier. *Development* 130, 2767-2777.
- Jenkins,T.D., Opitz,O.G., Okano,J., and Rustgi,A.K. (1998). Transactivation of the human keratin 4 and Epstein-Barr virus ED-L2 promoters by gut-enriched Kruppel-like factor. *Journal of Biological Chemistry* 273, 10747-10754.
- Katz,J.P., Yang,V.W., and Kaestner,K.H. (2000). The zinc-finger transcription factor Klf4 in the regulation of gastrointestinal epithelial proliferation and differentiation in vivo. *Gastroenterology* 118, A289.

## References

- Katz,J.P., Perreault,N., Goldstein,B.G., Lee,C.S., Labosky,P.A., Yang,V.W., and Kaestner,K.H. (2002). The zinc-finger transcription factor Klf4 is required for terminal differentiation of goblet cells in the colon. *Development* 129, 2619-2628.
- Katz,J.P., Perreault,N., Goldstein,B.G., Lee,C.S., Yang,V.W., and Kaestner,K.H. (2002). The zinc-finger transcription factor Klf4 is required for proper goblet cell differentiation in the colon. *Gastroenterology* 122, A95.
- Katz,J.P., Perreault,N., Goldstein,B.G., and Kaestner,K.H. (2003). The zinc-finger transcription factor Klf4 is required for normal gastric epithelial proliferation and differentiation in vivo. *Gastroenterology* 124, A92.
- Katz,J.P., Perreault,N., Goldstein,B.G., Actman,L., McNally,S.R., Silberg,D.G., Furth,E.E., and Kaestner,K.H. (2005). Loss of Klf4 in mice causes altered proliferation and differentiation and precancerous changes in the adult stomach. *Gastroenterology* 128, 935-945.
- Katze,M.G., He,Y., and Gale,M., Jr. (2002). Viruses and interferon: a fight for supremacy. *Nat. Rev. Immunol.* 2, 675-687.
- Kaushansky,K. (2006). Lineage-specific hematopoietic growth factors - Reply. *New England Journal of Medicine* 355, 527.
- Keller,G., Paige,C., Gilboa,E., and Wagner,E.F. (1985). Expression of A Foreign Gene in Myeloid and Lymphoid-Cells Derived from Multipotent Hematopoietic Precursors. *Nature* 318, 149-154.
- Kirstein,M., Brett,J., Radoff,S., Ogawa,S., Stern,D., and Vlassara,H. (1990). Advanced Protein Glycosylation Induces Transendothelial Human Monocyte Chemotaxis and Secretion of Platelet-Derived Growth-Factor - Role in Vascular-Disease of Diabetes and Aging. *Proceedings of the National Academy of Sciences of the United States of America* 87, 9010-9014.
- Knoops,L. and Renauld,J.C. (2004). IL-9 and its receptor: From signal transduction to tumorigenesis. *Growth Factors* 22, 207-215.
- Ko,J., Gendron-Fitzpatrick,A., and Splitter,G.A. (2002). Susceptibility of IFN regulatory factor-1 and IFN consensus sequence binding protein-deficient mice to brucellosis. *Journal of Immunology* 168, 2433-2440.
- Kondo,M., Weissman,I.L., and Akashi,K. (1997). Identification of clonogenic common lymphoid progenitors in mouse bone marrow. *Cell* 91, 661-672.
- Kondo,M., Scherer,D.C., Miyamoto,T., King,A.G., Akashi,K., Sugamura,K., and Weissman,I.L. (2000). Cell-fate conversion of lymphoid-committed progenitors by instructive actions of cytokines. *Nature* 407, 383-386.
- Kondo,M., Wagers,A.J., Manz,M.G., Prohaska,S.S., Scherer,D.C., Beilhack,G.E., Shizuru,J.A., and Weissman,I.L. (2003). Biology of hematopoietic stem cells and progenitors: Implications for clinical application. *Annual Review of Immunology* 21, 759-806.
- Konig,S., Schmidt,M., Alfken,J., Hoppe,G., Nagel,S., Horak,I., Levi,B.Z., Neubauer,A., and Wittig,B. (1997). A possible role of interferon consensus sequence binding protein (ICSBP) in regulating MHC class II expression through class II transactivator (CIITA). *Blood* 90, 3445.
- Kopf,M., Legros,G., Bachmann,M., Lamers,M.C., Bluethmann,H., and Kohler,G. (1993). Disruption of the Murine Il-4 Gene Blocks Th2 Cytokine Responses. *Nature* 362, 245-248.
- Kopf,M., Baumann,H., Freer,G., Freudenberg,M., Lamers,M., Kishimoto,T., Zinkernagel,R., Bluethmann,H., and Kohler,G. (1994). Impaired Immune and Acute-Phase Responses in Interleukin-6-Deficient Mice. *Nature* 368, 339-342.
- Kopf,M., Brombacher,F., Hodgkin,P.D., Ramsay,A.J., Milbourne,E.A., Dai,W.J., Ovington,K.S., Behm,C.A., Kohler,G., Young,I.G., and Matthaei,K.I. (1996). IL-5-Deficient mice have a developmental defect in CD5(+) B-1 cells and lack eosinophilia but have normal antibody and cytotoxic T cell responses. *Immunity* 4, 15-24.
- Korzenik,J.R. and Podolsky,D.K. (2006). Evolving knowledge and therapy of inflammatory bowel disease. *Nature Reviews Drug Discovery* 5, 197-209.
- Krause,D.S., Theise,N.D., Collector,M.I., Henegariu,O., Hwang,S., Gardner,R., Neutzel,S., and Sharkis,S.J. (2001). Multi-organ, multi-lineage engraftment by a single bone marrow-derived stem cell. *Cell* 105, 369-377.
- Krawczenko,A., Kieda,C., and Dus,D. (2005). The biological role and potential therapeutic application of interleukin 7. *Archivum Immunologiae et Therapiae Experimentalis* 53, 518-525.

## References

- Krishnaraju,K., Hoffman,B., and Liebermann,D.A. (1997). Lineage-specific regulation of hematopoiesis by HOX-B8 (HOX-2.4): Inhibition of granulocytic differentiation and potentiation of monocytic differentiation. *Blood* 90, 1840-1849.
- Kuhn,R., Lohler,J., Rennick,D., Rajewsky,K., and Muller,W. (1993). Interleukin-10-Deficient Mice Develop Chronic Enterocolitis. *Cell* 75, 263-274.
- Kuhn,R., Schwenk,F., Aguet,M., and Rajewsky,K. (1995). Inducible Gene Targeting in Mice. *Science* 269, 1427-1429.
- Kulessa,H., Frampton,J., and Graf,T. (1995). Gata-1 Reprograms Avian Myelomonocytic Cell-Lines Into Eosinophils, Thromboblasts, and Erythroblasts. *Genes & Development* 9, 1250-1262.
- Kunjathoor,V.V., Febbraio,M., Podrez,E.A., Moore,K.J., Andersson,L., Koehn,S., Rhee,J.S., Silverstein,R., Hoff,H.F., and Freeman,M.W. (2002). Scavenger receptors class A-I/II and CD36 are the principal receptors responsible for the uptake of modified low density lipoprotein leading to lipid loading in macrophages. *J. Biol. Chem.* 277, 49982-49988.
- Kuo,C.T., Veselits,M.L., and Leiden,J.M. (1997). LKLF: A transcriptional regulator of single-positive T cell quiescence and survival. *Science* 277, 1986-1990.
- Kurzrock,R. (2005). Thrombopoietic factors in chronic bone marrow failure states: The platelet problem revisited. *Clinical Cancer Research* 11, 1361-1367.
- Lacaud,G., Carlsson,L., and Keller,G. (1998). Identification of a fetal hematopoietic precursor with B cell, T cell, and macrophage potential. *Immunity* 9, 827-838.
- Lampinen,M., Carlson,M., Hakansson,L.D., and Venge,P. (2004). Cytokine-regulated accumulation of eosinophils in inflammatory disease. *Allergy* 59, 793-805.
- Larson,R.A. and Le Beau,M.M. (2005). Therapy-related myeloid leukaemia: A model for leukemogenesis in humans. *Chemico-Biological Interactions* 153, 187-195.
- Lawrence,H.J., Helgason,C.D., Sauvageau,G., Fong,S., Izon,D.J., Humphries,R.K., and Largman,C. (1997). Mice bearing a targeted interruption of the homeobox gene HOXA9 have defects in myeloid, erythroid, and lymphoid hematopoiesis. *Blood* 89, 1922-1930.
- Lee,S.L., Wang,Y., and Milbrandt,J. (1996). Unimpaired macrophage differentiation and activation in mice lacking the zinc finger transcription factor NGFI-A (EGR1). *Molecular and Cellular Biology* 16, 4566-4572.
- Leng,C., Lusatis,S., Horak,I.D., Pezzutto,A., and Mapara,M.Y. (2004). ICSBP gene deficient mice are highly susceptible to the development of graft-versus-host disease (GVHD) following administration of delayed donor lymphocyte infusions (DLI). *Blood* 104, 329B.
- Levy,D.A. and Frondoza,C. (1983). Immunity to Intestinal Parasites - Role of Mast-Cells and Goblet Cells. *Federation Proceedings* 42, 1750-1755.
- Li,M.O., Wan,Y.Y., Sanjabi,S., Robertson,A.K., and Flavell,R.A. (2006). Transforming growth factor-beta regulation of immune responses. *Annu. Rev. Immunol.* 24, 99-146.
- Li,Y., McClintick,J., Zhong,L., Edenberg,H.J., Yoder,M.C., and Chan,R.J. (2005). Murine embryonic stem cell differentiation is promoted by SOCS-3 and inhibited by the zinc finger transcription factor Klf4. *Blood* 105, 635-637.
- Lieschke,G.J., Grail,D., Hodgson,G., Metcalf,D., Stanley,E., Cheers,C., Fowler,K.J., Basu,S., Zhan,Y.F., and Dunn,A.R. (1994). Mice Lacking Granulocyte-Colony-Stimulating Factor Have Chronic Neutropenia, Granulocyte and Macrophage Progenitor-Cell Deficiency, and Impaired Neutrophil Mobilization. *Blood* 84, 1737-1746.
- Lin,C.S., Lim,S.K., DAgati,V., and Costantini,F. (1996). Differential effects of an erythropoietin receptor gene disruption on primitive and definitive erythropoiesis. *Genes & Development* 10, 154-164.
- Littlewood,T.D., Hancock,D.C., Danielian,P.S., Parker,M.G., and Even,G.I. (1995). A Modified Estrogen-Receptor Ligand-Binding Domain As An Improved Switch for the Regulation of Heterologous Proteins. *Nucleic Acids Research* 23, 1686-1690.
- Lofts,F.J. and Pettengell,R. (1998). Myeloid growth factors in oncology. *Expert Opin. Investig. Drugs* 7, 1955-1976.

## References

- Louahed,J., Zhou,Y.H., Maloy,W.L., Rani,P.U., Weiss,C., Tomer,Y., Vink,A., Renauld,J.C., Van Snick,J., Nicolaides,N.C., Levitt,R.C., and Haczku,A. (2001). Interleukin 9 promotes influx and local maturation of eosinophils. *Blood* 97, 1035-1042.
- Lowe,S.W. and Ruley,H.E. (1993). Stabilization of the P53 Tumor Suppressor Is Induced by Adenovirus-5 E1A and Accompanies Apoptosis. *Genes & Development* 7, 535-545.
- Mackarehtschian,K., Hardin,J.D., Moore,K.A., Boast,S., Goff,S.P., and Lemischka,I.R. (1995). Targeted Disruption of the Flk2/Flt3 Gene Leads to Deficiencies in Primitive Hematopoietic Progenitors. *Immunity* 3, 147-161.
- Magram,J., Connaughton,S.E., Warrier,R.R., Carvajal,D.M., Wu,C.Y., Ferrante,J., Stewart,C., Sarmiento,U., Faherty,D.A., and Gately,M.K. (1996). IL-12-deficient mice are defective in IFN gamma production and type 1 cytokine responses. *Immunity* 4, 471-481.
- Manz,M.G., Traver,D., Miyamoto,T., Weissman,I.L., and Akashi,K. (2001). Dendritic cell potentials of early lymphoid and myeloid progenitors. *Blood* 97, 3333-3341.
- Masumi,A., Tamaoki,S., Wang,I.M., Ozato,K., and Komuro,K. (2002). IRF-8/ICSBP and IRF-1 cooperatively stimulate mouse IL-12 promoter activity in macrophages. *Febs Letters* 531, 348-353.
- McArthur,G.A., Rohrschneider,L.R., and Johnson,G.R. (1994). Induced Expression of C-Fms in Normal Hematopoietic-Cells Shows Evidence for Both Conservation and Lineage Restriction of Signal-Transduction in Response to Macrophage-Colony-Stimulating Factor. *Blood* 83, 972-981.
- McArthur,G.A., Longmore,G.D., Klingler,K., and Johnson,G.R. (1995). Lineage-Restricted Recruitment of Immature Hematopoietic Progenitor Cells in Response to Epo After Normal Hematopoietic-Cell Transfection with Epor. *Experimental Hematology* 23, 645-654.
- McInnes,I., Martin,R., Zimmerman-Gorska,I., Nayiager,S., Sun,G., Patel,A., and Appleton,B. (2004). Safety and efficacy of a human monoclonal antibody to IL-15 (AMG 714) in patients with rheumatoid arthritis (RA): Results from a multicenter, randomized, double-blind, placebo-controlled trial. *Arthritis and Rheumatism* 50, S241.
- McKenzie,G.J., Emson,C.L., Bell,S.E., Anderson,S., Fallon,P., Zurawski,G., Murray,R., Grencis,R., and McKenzie,A.N.J. (1998). Impaired development of Th2 cells in IL-13-deficient mice. *Immunity* 9, 423-432.
- McKercher,S.R., Torbett,B.E., Anderson,K.L., Henkel,G.W., Vestal,D.J., Baribault,H., Klemsz,M., Feeney,A.J., Wu,G.E., Paige,C.J., and Maki,R.A. (1996). Targeted disruption of the PU.1 gene results in multiple hematopoietic abnormalities. *Embo Journal* 15, 5647-5658.
- McNagny,K. and Graf,T. (2002). Making eosinophils through subtle shifts in transcription factor expression. *Journal of Experimental Medicine* 195, f43-f47.
- McNagny,K.M., Sieweke,M.H., Doderlein,G., Graf,T., and Nerlov,C. (1998). Regulation of eosinophil-specific gene expression by a C/EBP-Ets complex and GATA-1. *Embo Journal* 17, 3669-3680.
- Metcalf,D. (1993). Hematopoietic Regulators - Redundancy Or Subtlety. *Blood* 82, 3515-3523.
- Metcalf,D. (1999). Stem cells, pre-progenitor cells and lineage-committed cells: Are our dogmas correct? *Hematopoietic Stem Cells* 872, 289-304.
- Metcalf,D. (1999). Cellular hematopoiesis in the twentieth century. *Seminars in Hematology* 36, 5-12.
- Michieli,P., Chedid,M., Lin,D., Pierce,J.H., Mercer,W.E., and Givol,D. (1994). Induction of Waf1/Cip1 by A P53-Independent Pathway. *Cancer Research* 54, 3391-3395.
- Migliaccio,A.R., Rana,R.A., Sanchez,M., Lorenzini,R., Centurione,L., Bianchi,L., Vannucchi,A.M., Migliaccio,G., and Orkin,S.H. (2003). GATA-1 as a regulator of mast cell differentiation revealed by the phenotype of the GATA-1(low) mouse mutant. *Journal of Experimental Medicine* 197, 281-296.
- Miller,K.A., Eklund,E.A., Peddinghaus,M.L., Cao,Z.J., Fernandes,N., Turk,P.W., Thimmapaya,B., and Weitzman,S.A. (2001). Kruppel-like factor 4 regulates laminin alpha 3A expression in mammary epithelial cells. *Journal of Biological Chemistry* 276, 42863-42868.
- Mitre,E. and Nutman,T.B. (2006). Basophils, basophilia and helminth infections. *Chem. Immunol. Allergy* 90, 141-156.
- Mizutani,K., Takeuchi,S., Ohashi,Y., Yakushiji,M., Nishimura,H., Takahashi,T., Maruhashi,T., Ueda,K., Noda,K., Watanabe,Y., Kawana,T., Terashima,Y., Ochiai,K., Goto,S., Takaku,F., and Motoyoshi,K. (2003).

## References

- Clinical usefulness of macrophage colony-stimulating factor for ovarian cancers: Long-term prognosis after five years. *Oncology Reports* 10, 127-131.
- Montecino-Rodriguez,E., Leathers,H., and Dorshkind,K. (2001). Bipotential B-macrophage progenitors are present in adult bone marrow. *Nat. Immunol.* 2, 83-88.
- Moore,K.W., de Waal,M.R., Coffman,R.L., and O'Garra,A. (2001). Interleukin-10 and the interleukin-10 receptor. *Annu. Rev. Immunol.* 19, 683-765.
- Moreaugachelin,F., Tavitian,A., and Tambourin,P. (1988). Spi-1 Is A Putative Oncogene in Virally Induced Murine Erythroleukemias. *Nature* 331, 277-280.
- Moritz,T., Patel,V.P., and Williams,D.A. (1994). Bone-Marrow Extracellular-Matrix Molecules Improve Gene-Transfer Into Human Hematopoietic-Cells Via Retroviral Vectors. *Journal of Clinical Investigation* 93, 1451-1457.
- Moritz,T., Hanenberg,H., Kato,I., and Williams,D.A. (1996). Fibronectin improves retroviral gene transfer by co-localizing vector and target cells. *Experimental Hematology* 24, 204.
- Morrison,S.J. and Weissman,I.L. (1994). The Long-Term Repopulating Subset of Hematopoietic Stem-Cells Is Deterministic and Isolatable by Phenotype. *Immunity* 1, 661-673.
- Morrison,S.J., Shah,N.M., and Anderson,D.J. (1997). Regulatory mechanisms in stem cell biology. *Cell* 88, 287-298.
- Motoyoshi,K. (1998). Biological activities and clinical application of M-CSF. *International Journal of Hematology* 67, 109-122.
- Murphy,J.E., Tedbury,P.R., Homer-Vanniasinkam,S., Walker,J.H., and Ponnambalam,S. (2005). Biochemistry and cell biology of mammalian scavenger receptors. *Atherosclerosis* 182, 1-15.
- Nakahata,T. and Ogawa,M. (1982). Identification in Culture of A Class of Hematopoietic Colony-Forming-Units with Extensive Capability to Self-Renew and Generate Multipotential Hematopoietic Colonies. *Proceedings of the National Academy of Sciences of the United States of America-Biological Sciences* 79, 3843-3847.
- Nakahata,T. and Ogawa,M. (1982). Clonal Origin of Murine Hematopoietic Colonies with Apparent Restriction to Granulocyte-Macrophage-Megakaryocyte (Gmm) Differentiation. *Journal of Cellular Physiology* 111 , 239-246.
- Nathan,C. (2006). Neutrophils and immunity: challenges and opportunities. *Nature Reviews Immunology* 6, 173-182.
- Nelson,N., Marks,M.S., Driggers,P.H., and Ozato,K. (1993). Interferon Consensus Sequence-Binding Protein, A Member of the Interferon Regulatory Factor Family, Suppresses Interferon-Induced Gene-Transcription. *Molecular and Cellular Biology* 13, 588-599.
- Nelson,N., Kanno,Y., Hong,C., Contursi,C., Fujita,T., Fowlkes,B.J., OConnell,E., Li,J.H., Paul,W.E., Jankovic,D., Sher,A.F., Coligan,J.E., Thornton,A., Appella,E., Yang,Y.L., and Ozato,K. (1996). Expression of IFN regulatory factor family proteins in lymphocytes - Induction of Stat-1 and IFN consensus sequence binding protein expression by T cell activation. *Journal of Immunology* 156, 3711-3720.
- Nemunaitis,J. (1998). Use of macrophage colony-stimulating factor in the treatment of fungal infections. *Clinical Infectious Diseases* 26, 1279-1281.
- Nerlov,C. and Graf,T. (1998). PU.1 induces myeloid lineage commitment in multipotent hematopoietic progenitors. *Genes & Development* 12, 2403-2412.
- Nguyen,H.Q., Hoffmanliebermann,B., and Liebermann,D.A. (1993). The Zinc Finger Transcription Factor Egr-1 Is Essential for and Restricts Differentiation Along the Macrophage Lineage. *Cell* 72, 197-209.
- Nishinakamura,R., Miyajima,A., Mee,P.J., Tybulewicz,V.L.J., and Murray,R. (1996). Hematopoiesis in mice lacking the entire granulocyte-macrophage colony-stimulating factor/interleukin-3/interleukin-5 functions. *Blood* 88, 2458-2464.
- Noti,J.D., Johnson,A.K., and Dillon,J.D. (2005). The leukocyte integrin gene CD11d is repressed by gut-enriched Kruppel-like factor 4 in myeloid cells. *Journal of Biological Chemistry* 280, 3449-3457.
- Nuez,B., Michalovich,D., Bygrave,A., Ploemacher,R., and Grosveld,F. (1995). Defective Hematopoiesis in Fetal Liver Resulting from Inactivation of the Eklf Gene. *Nature* 375, 316-318.

## References

- Ogawa,M. and Nakahata,T. (1982). Identification in Culture of A New Class of Hematopoietic Colony-Forming-Units with Extensive Capability to Self-Renew and Generate Multipotential Hematopoietic Colonies. *Experimental Hematology 10*, 166.
- Ogawa,M. (1993). Differentiation and Proliferation of Hematopoietic Stem-Cells. *Blood 81*, 2844-2853.
- Ohteki,T. (2002). Critical role for IL-15 in innate immunity. *Curr. Mol. Med. 2*, 371-380.
- Okano,J., Opitz,O.G., Nakagawa,H., Jenkins,T.D., Friedman,S.L., and Rustgi,A.K. (2000). Physical interaction between the Kruppel-like transcriptional factors Zf9/KLF6 and GKLF/KLF4 and its effect on the human keratin 4 promoter. *Gastroenterology 118*, A865.
- Onodera,K., Takahashi,S., Nishimura,S., Ohta,J., Motohashi,H., Yomogida,K., Hayashi,N., Engel,J.D., and Yamamoto,M. (1997). GATA-1 transcription is controlled by distinct regulatory mechanisms during primitive and definitive erythropoiesis. *Proceedings of the National Academy of Sciences of the United States of America 94*, 4487-4492.
- Orkin,S.H. (2003). Priming the hematopoietic pump. *Immunity 19*, 633-634.
- Pandya,A.Y., Talley,L.I., Frost,A.R., Fitzgerald,T.J., Trivedi,V., Chakravarthy,M., Chhieng,D.C., Grizzle,W.E., Engler,J.A., Krontiras,H., Bland,K.I., LoBuglio,A.F., Lobo-Ruppert,S.M., and Ruppert,J.M. (2004). Nuclear localization of KLF4 is associated with an aggressive phenotype in early-stage breast cancer. *Clinical Cancer Research 10*, 2709-2719.
- Parker,L.C., Whyte,M.K., Dower,S.K., and Sabroe,I. (2005). The expression and roles of Toll-like receptors in the biology of the human neutrophil. *J. Leukoc. Biol. 77*, 886-892.
- Pevny,L., Simon,M.C., Robertson,E., Klein,W.H., Tsai,S.F., DAgati,V., Orkin,S.H., and Costantini,F. (1991). Erythroid-Differentiation in Chimeric Mice Blocked by A Targeted Mutation in the Gene for Transcription Factor Gata-1. *Nature 349*, 257-260.
- Piccinni,S.A., Bolcato-Bellemin,A.L., Klein,A., Yang,V.W., Kedinger,M., Simon-Assmann,P., and Lefebvre,O. (2004). Kruppel-like factors regulate the Lama1 gene encoding the laminin alpha 1 chain. *Journal of Biological Chemistry 279*, 9103-9114.
- Poortinga,G., Hannan,K.M., Snelling,H., Walkley,C.R., Jenkins,A., Sharkey,K., Wall,M., Brandenburger,Y., Palatsides,M., Pearson,R.B., McArthur,G.A., and Hannan,R.D. (2004). MAD1 and c-MYC regulate UBF and rDNA transcription during granulocyte differentiation. *Embo Journal 23*, 3325-3335.
- Qian,Z.J., Fernald,A.A., Godley,L.A., Larson,R.A., and Le Beau,M.M. (2002). Expression profiling of CD34(+) hematopoietic stem/progenitor cells reveals distinct subtypes of therapy-related acute myeloid leukemia. *Proceedings of the National Academy of Sciences of the United States of America 99*, 14925-14930.
- Querfurth,E., Schuster,M., Kulessa,H., Crispino,J.D., Doderlein,G., Orkin,S.H., Graf,T., and Nerlov,C. (2000). Antagonism between C/EBP beta and FOG in eosinophil lineage commitment of multipotent hematopoietic progenitors. *Genes & Development 14*, 2515-2525.
- Quesenberry,P.J., Colvin,G.A., Abedi,M., Dooner,G., Dooner,M., Aliotta,J., Keaney,P., Luo,L.G., Demers,D., Peterson,A., Foster,B., and Greer,D. (2005). The stem cell continuum. *Hematopoietic Stem Cells V 1044*, 228-235.
- Radomska,H.S., Huettner,C.S., Zhang,P., Cheng,T., Scadden,D.T., and Tenen,D.G. (1998). CCAAT enhancer binding protein alpha is a regulatory switch sufficient for induction of granulocytic development from bipotential myeloid progenitors. *Molecular and Cellular Biology 18*, 4301-4314.
- Radtke,F., Wilson,A., Stark,G., Bauer,M., van Meerwijk,J., MacDonald,H.R., and Aguet,M. (1999). Deficient T cell fate specification in mice with an induced inactivation of Notch1. *Immunity 10*, 547-558.
- Ramsfjell,V., Borge,O.P., Cardier,J., Murphy,M.J., Jr., Lyman,S.D., Lok,S., and Jacobsen,S.E. (1996). Thrombopoietin, but not erythropoietin, directly stimulates multilineage growth of primitive murine bone marrow progenitor cells in synergy with early acting cytokines: distinct interactions with the ligands for c-kit and FLT3. *Blood 88*, 4481-4492.
- Rekhtman,N., Radparvar,F., Evans,T., and Skoultchi,A.I. (1999). Direct interaction of hematopoietic transcription factors PU.1 and GATA-1: functional antagonism in erythroid cells. *Genes & Development 13*, 1398-1411.
- Renauld,J.C., Vanderlugt,N., Vink,A., Vanroon,M., Godfraind,C., Warnier,G., Merz,H., Feller,A., Berns,A., and Vansnick,J. (1994). Thymic Lymphomas in Interleukin-9 Transgenic Mice. *Oncogene 9*, 1327-1332.

## References

- Rennick,D.M., Thompsonsnipes,L., Coffman,R.L., Seymour,B.W.P., Jackson,J.D., and Hudak,S. (1990). Invivo Administration of Antibody to Interleukin-5 Inhibits Increased Generation of Eosinophils and Their Progenitors in Bone-Marrow of Parasitized Mice. *Blood* 76, 312-316.
- Robb,L., Drinkwater,C.C., Metcalf,D., Li,R.L., Kontgen,F., Nicola,N.A., and Begley,C.G. (1995). Hematopoietic and Lung Abnormalities in Mice with A Null Mutation of the Common Beta-Subunit of the Receptors for Granulocyte-Macrophage Colony-Stimulating Factor and Interleukin-3 and Interleukin-5. *Proceedings of the National Academy of Sciences of the United States of America* 92, 9565-9569.
- Roberts,R., Gallagher,J., Spooncer,E., Allen,T.D., Bloomfield,F., and Dexter,T.M. (1988). Heparan-Sulfate Bound Growth-Factors - A Mechanism for Stromal Cell-Mediated Hematopoiesis. *Nature* 332, 376-378.
- Rook,G.A.W., Dheda,K., and Zumla,A. (2005). Do successful tuberculosis vaccines need to be immunoregulatory rather than merely Th1-boosting? *Vaccine* 23, 2115-2120.
- Rook,G.A.W., Dheda,K., and Zumla,A. (2005). Immune responses to tuberculosis in developing countries: implications for new vaccines. *Nature Reviews Immunology* 5, 661-667.
- Rosenbauer,F., Wagner,K., Kutok,J.L., Iwasaki,H., Le Beau,M.M., Okuno,Y., Akashi,K., Fiering,S., and Tenen,D.G. (2004). Acute myeloid leukemia induced by graded reduction of a lineage-specific transcription factor, PU.1. *Nature Genetics* 36, 624-630.
- Rosmarin,A.G., Yang,Z., and Ressendes,K.K. (2005). Transcriptional regulation in myelopoiesis: Hematopoietic fate choice, myeloid differentiation, and leukemogenesis. *Exp. Hematol.* 33, 131-143.
- Rothenberg,M.E. and Hogan,S.P. (2006). The eosinophil. *Annual Review of Immunology* 24, 147-174.
- Rowland,B.D., Bernards,R., and Peepoer,D.S. (2005). The KLF4 tumour suppressor is a transcriptional repressor of p53 that acts as a context-dependent oncogene. *Nature Cell Biology* 7, 1074-1082.
- Saifudeen,Z., Dipp,S., Fan,H., and El Dahr,S.S. (2005). Combinatorial control of the bradykinin B2 receptor promoter by p53, CREB, KLF-4, and CBP: implications for terminal nephron differentiation. *American Journal of Physiology-Renal Physiology* 288, F899-F909.
- Sarkar,J., Seshadri,V., Tripoulas,N.A., Ketterer,M.E., and Fox,P.L. (2003). Role of ceruloplasmin in macrophage iron efflux during hypoxia. *Journal of Biological Chemistry* 278, 44018-44024.
- Sauvageau,G., Thorsteinsdottir,U., Hough,M.R., Hugo,P., Lawrence,H.J., Largman,C., and Humphries,R.K. (1997). Overexpression of HOXB3 in hematopoietic cells causes defective lymphoid development and progressive myeloproliferation. *Immunity* 6, 13-22.
- SchartonKersten,T., Contursi,C., Masumi,A., Sher,A., and Ozato,K. (1997). Interferon consensus sequence binding protein-deficient mice display impaired resistance to intracellular infection due to a primary defect in interleukin 12 p40 induction. *Journal of Experimental Medicine* 186, 1523-1534.
- Scheller,M., Foerster,J., Heyworth,C.M., Waring,J.F., Lohler,J., Gilmore,G.L., Shadduck,R.K., Dexter,T.M., and Horak,I. (1999). Altered development and cytokine responses of myeloid progenitors in the absence of transcription factor, interferon consensus sequence binding protein. *Blood* 94, 3764-3771.
- Schepers,H., Geugien,M., Eggen,B.J.L., and Vellenga,E. (2003). Constitutive cytoplasmic localization of p21(Waf1/Cip1) affects the apoptotic process in monocytic leukaemia. *Leukemia* 17, 2113-2121.
- Schmidt,M., Nagel,S., Proba,J., Hochhaus,A., Waring,J.F., Rosenbauer,F., Ehninger,G., Huhn,D., Wittig,B., Hehlmann,R., Horak,I., and Neubauer,A. (1997). Lack of interferon consensus sequence binding protein (ICSBP) transcripts in human myeloid leukemias and induction by interferon-alpha. *Blood* 90, 878.
- Schmidt,M., Nagel,S., Proba,J., Thiede,C., Ritter,M., Waring,J.F., Rosenbauer,F., Huhn,D., Wittig,B., Horak,I., and Neubauer,A. (1998). Lack of interferon consensus sequence binding protein (ICSBP) transcripts in human myeloid leukemias. *Blood* 91, 22-29.
- Schorle,H., Holtschke,T., Hunig,T., Schimpl,A., and Horak,I. (1991). Development and Function of T-Cells in Mice Rendered Interleukin-2 Deficient by Gene Targeting. *Nature* 352, 621-624.
- Scott,E.W., Simon,M.C., Anastasi,J., and Singh,H. (1994). Requirement of Transcription Factor Pu.1 in the Development of Multiple Hematopoietic Lineages. *Science* 265, 1573-1577.
- Scott,E.W., Fisher,R.C., Olson,M.C., Kehrli,E.W., Simon,M.C., and Singh,H. (1997). PU.1 functions in a cell-autonomous manner to control the differentiation of multipotential lymphoid-myeloid progenitors. *Immunity* 6, 437-447.

## References

- Segre,J.A., Bauer,C., and Fuchs,E. (1999). Klf4 is a transcription factor required for establishing the barrier function of the skin. *Nature Genetics* 22, 356-360.
- Shie,J.L., Chen,Z.Y., Fu,M.F., Pestell,R.G., and Tseng,C.C. (2000). Gut-enriched Kruppel-like factor represses cyclin D1 promoter activity through Sp1 motif. *Nucleic Acids Research* 28, 2969-2976.
- Shields,J.M., Christy,R.J., and Yang,V.W. (1996). Identification and characterization of a gene encoding a gut-enriched Kruppel-like factor expressed during growth arrest. *Journal of Biological Chemistry* 271, 20009-20017.
- Shields,J.M. and Yang,V.W. (1996). The gut-enriched Kruppel-like factor (GKLF) is a growth arrest-specific gene expressed in the mouse colonic epithelium. *Gastroenterology* 110, A360.
- Shivdasani,R.A. and Orkin,S.H. (1996). The transcriptional control of hematopoiesis. *Blood* 87, 4025-4039.
- Shivdasani,R.A., Fujiwara,Y., McDevitt,M.A., and Orkin,S.H. (1997). A lineage-selective knockout establishes the critical role of transcription factor GATA-1 in megakaryocyte growth and platelet development. *Embo Journal* 16, 3965-3973.
- Sieweke,M.H., Tekotte,H., Frampton,J., and Graf,T. (1996). MafB is an interaction partner and repressor of Ets-1 that inhibits erythroid differentiation. *Cell* 85, 49-60.
- Sitnicka,E., Lin,N., Priestley,G.V., Fox,N., Brody,V.C., Wolf,N.S., and Kaushansky,K. (1996). The effect of thrombopoietin on the proliferation and differentiation of murine hematopoietic stem cells. *Blood* 87, 4998-5005.
- Skalnik,D.G. (2002). Transcriptional mechanisms regulating myeloid-specific genes. *Gene* 284, 1-21.
- Skurkovich,S. and Skurkovich,B. (2005). Anticytokine therapy, especially anti-interferon-gamma, as a pathogenetic treatment in Th-1 autoimmune diseases. *Autoimmune Diseases and Treatment: Organ-Specific and Systemic Disorders* 1051, 684-700.
- Smith,M.A., Court EL, and Smith,J.G. (2001). Stem cell factor: laboratory and clinical aspects. *Blood Rev.* 15, 191-197.
- Soddu,S., Blandino,G., Scardigli,R., Coen,S., Marchetti,A., Rizzo,M.G., Bossi,G., Cimino,L., Crescenzi,M., and Sacchi,A. (1996). Interference with p53 protein inhibits hematopoietic and muscle differentiation. *Journal of Cell Biology* 134, 193-204.
- Stanley,E., Lieschke,G.J., Grail,D., Metcalf,D., Hodgson,G., Gall,J.A.M., Maher,D.W., Cebon,J., Sinickas,V., and Dunn,A.R. (1994). Granulocyte/Macrophage Colony-Stimulating Factor-Deficient Mice Show No Major Perturbation of Hematopoiesis But Develop A Characteristic Pulmonary Pathology. *Proceedings of the National Academy of Sciences of the United States of America* 91, 5592-5596.
- Steinman,R.A., Huang,J.P., Yaroslavskiy,B., Goff,J.P., Ball,E.D., and Nguyen,A. (1998). Regulation of p21(WAF1) expression during normal myeloid differentiation. *Blood* 91, 4531-4542.
- Steinman,R.A. and Johnson,D.E. (2000). p21WAF1 prevents down-modulation of the apoptotic inhibitor protein c-IAP1 and inhibits leukemic apoptosis. *Mol. Med.* 6, 736-749.
- Stroncek,D. (2004). Granulocyte gene polymorphisms and mutations: effects on immune response and neutrophil proliferation. *Vox Sanguinis* 87, 101-104.
- Sutton,S.A., Assa'ad,A.H., and Rothenberg,M.E. (2005). Anti-IL-5 and hypereosinophilic syndromes. *Clinical Immunology* 115, 51-60.
- Suzuki,A., Tsutomi,Y., Miura,M., and Akahane,K. (1999). Caspase 3 inactivation to suppress Fas-mediated apoptosis: identification of binding domain with p21 and ILP and inactivation machinery by p21. *Oncogene* 18, 1239-1244.
- Suzuki,T., Shen,H.F., Akagi,K., Morse,H.C., Malley,J.D., Naiman,D.Q., Jenkins,N.A., and Copeland,N.G. (2002). New genes involved in cancer identified by retroviral tagging. *Nature Genetics* 32, 166-174.
- Svedberg,H., Chylicki,K., Baldetorp,B., Rauscher,F.J., and Gullberg,U. (1998). Constitutive expression of the Wilms' tumor gene (WT1) in the leukemic cell line U937 blocks parts of the differentiation program. *Oncogene* 16, 925-932.
- Takahashi,K., Naito,M., and Takeya,M. (1996). Development and heterogeneity of macrophages and their related cells through their differentiation pathways. *Pathology International* 46, 473-485.
- Takeda,K., Tsutsui,H., Yoshimoto,T., Adachi,O., Yoshida,N., Kishimoto,T., Okamura,H., Nakanishi,K., and Akira,S. (1998). Defective NK cell activity and Th1 response in IL-18-deficient mice. *Immunity* 8, 383-390.

## References

- Tamura,T., Nagamura-Inoue,T., Shmeltzer,Z., Kuwata,T., and Ozato,K. (2000). ICSBP directs bipotential myeloid progenitor cells to differentiate into mature macrophages. *Immunity* 13, 155-165.
- Taniguchi,T., Ogasawara,K., Takaoka,A., and Tanaka,N. (2001). IRF family of transcription factors as regulators of host defense. *Annual Review of Immunology* 19, 623-655.
- Terada,N., Hamazaki,T., Oka,M., Hoki,M., Mastalerz,D.M., Nakano,Y., Meyer,E.M., Morel,L., Petersen,B.E., and Scott,E.W. (2002). Bone marrow cells adopt the phenotype of other cells by spontaneous cell fusion. *Nature* 416, 542-545.
- Terskikh,A.V., Miyamoto,T., Chang,C., Diatchenko,L., and Weissman,I.L. (2003). Gene expression analysis of purified hematopoietic stem cells and committed progenitors. *Blood* 102, 94-101.
- Terszowski,G., Waskow,C., Conradt,P., Lenze,D., Koenigsmann,J., Carstanjen,D., Horak,I., and Rodewald,H.R. (2005). Prospective isolation and global gene expression analysis of the erythrocyte colony-forming unit (CFU-E). *Blood* 105, 1937-1945.
- Thomas,D., Vadas,M., and Lopez,A. (2004). Regulation of haematopoiesis by growth factors - emerging insights and therapies. *Expert Opinion on Biological Therapy* 4, 869-879.
- Thorsteinsdottir,U., Sauvageau,G., Hough,M.R., Dragowska,W., Lansdorp,P.M., Lawrence,H.J., Largman,C., and Humphries,R.K. (1997). Overexpression of HOXA10 in murine hematopoietic cells perturbs both myeloid and lymphoid differentiation and leads to acute myeloid leukemia. *Molecular and Cellular Biology* 17, 495-505.
- Till,J.E. and Mcculloch,E.A. (1980). Hematopoietic Stem-Cell Differentiation. *Biochimica et Biophysica Acta* 605, 431-459.
- Ton-That,H., Kaestner,K.H., Shields,J.M., Mahatanankoon,C.S., and Yang,V.W. (1997). Expression of the gut-enriched Kruppel-like factor gene during development and intestinal tumorigenesis. *Febs Letters* 419, 239-243.
- Tsai,F.Y., Keller,G., Kuo,F.C., Weiss,M., Chen,J.Z., Rosenblatt,M., Alt,F.W., and Orkin,S.H. (1994). An Early Hematopoietic Defect in Mice Lacking the Transcription Factor Gata-2. *Nature* 371, 221-226.
- Tsai,S.F., Strauss,E., and Orkin,S.H. (1991). Functional-Analysis and Invivo Footprinting Implicate the Erythroid Transcription Factor Gata-1 As A Positive Regulator of Its Own Promoter. *Genes & Development* 5, 919-931.
- Tsujimura,H., Nagamura-Inoue,T., Tamura,T., and Ozato,K. (2002). IFN consensus sequence binding protein/IFN regulatory factor-8 guides bone marrow progenitor cells toward the macrophage lineage. *Journal of Immunology* 169, 1261-1269.
- Tsukamoto,K., Kinoshita,M., Kojima,K., Mikuni,Y., Kudo,M., Mori,M., Fujita,M., Horie,E., Shimazu,N., and Teramoto,T. (2002). Synergically increased expression of CD36, CLA-1 and CD68, but not of SR-A and LOX-1, with the progression to foam cells from macrophages. *J. Atheroscler. Thromb.* 9, 57-64.
- Vanzant,G. (1984). Studies of Hematopoietic Stem-Cells Spared by 5-Fluorouracil. *Journal of Experimental Medicine* 159, 679-690.
- Vlassara,H., Brownlee,M., Manogue,K.R., Dinarello,C.A., and Pasagian,A. (1988). Cachectin Tnf and IL-1 Induced by Glucose-Modified Proteins - Role in Normal Tissue Remodeling. *Science* 240, 1546-1548.
- Vonfreedenjeffry,U., Vieira,P., Lucian,L.A., Mcneil,T., Burdach,S.E.G., and Murray,R. (1995). Lymphopenia in Interleukin (IL)-7 Gene-Deleted Mice Identifies IL-7 As A Nonredundant Cytokine. *Journal of Experimental Medicine* 181, 1519-1526.
- Vyas,P., McDevitt,M.A., Cantor,A.B., Katz,S.G., Fujiwara,Y., and Orkin,S.H. (1999). Different sequence requirements for expression in erythroid and megakaryocytic cells within a regulatory element upstream of the GATA-1 gene. *Development* 126, 2799-2811.
- Wagers,A.J., Sherwood,R.I., Christensen,J.L., and Weissman,I.L. (2002). Little evidence for developmental plasticity of adult hematopoietic stem cells. *Science* 297, 2256-2259.
- Waldmann,T.A. (2006). Effective cancertherapy through immunomodulation. *Annual Review of Medicine* 57, 65-81.
- Waldmann,T.A. (2006). The biology of interleukin-2 and interleukin-15: implications for cancer therapy and vaccine design. *Nat. Rev. Immunol.* 6, 595-601.
- Wang,H., Yang,L., Jamaluddin,M.S., and Boyd,D.D. (2004). The Kruppel-like KLF4 transcription factor, a novel regulator of urokinase receptor expression, drives synthesis of this binding site in colonic crypt luminal surface epithelial cells. *Journal of Biological Chemistry* 279, 22674-22683.

## References

- Watt,F.M. and Hogan,B.L.M. (2000). Out of Eden: Stem cells and their niches. *Science* 287, 1427-1430.
- Watts,T.H. (2005). Tnf/ttnfr family members in costimulation of T cell responses. *Annual Review of Immunology* 23, 23-68.
- Wei,D.Y., Gong,W.D., Kanai,M., Schlunk,C., Wang,L.W., Yao,J.C., Wu,T.T., Huang,S.Y., and Xie,K.P. (2005). Drastic down-regulation of Kruppel-like factor 4 expression is critical in human gastric cancer development and progression. *Cancer Research* 65, 2746-2754.
- Weintraub,H. (1985). Assembly and Propagation of Repressed and Derepressed Chromosomal States. *Cell* 42, 705-711.
- Weiss,M.J. and Orkin,S.H. (1995). Gata Transcription Factors - Key Regulators of Hematopoiesis. *Experimental Hematology* 23, 99-107.
- Weissman,I.L., Anderson,D.J., and Gage,F. (2001). Stem and progenitor cells: Origins, phenotypes, lineage commitments, and transdifferentiations. *Annual Review of Cell and Developmental Biology* 17, 387-403.
- Wodnar-Filipowicz,A. (2003). Flt3 ligand: Role in control of hematopoietic and immune functions of the bone marrow. *News in Physiological Sciences* 18, 247-251.
- Woodcock,J.M., Bagley,C.J., and Lopez,A.F. (1999). The functional basis of granulocyte-macrophage colony stimulating factor, interleukin-3 and interleukin-5 receptor activation, basic and clinical implications. *International Journal of Biochemistry & Cell Biology* 31, 1017-1025.
- Wu,H., Liu,X., Jaenisch,R., and Lodish,H.F. (1995). Generation of Committed Erythroid Bfu-e and Cfu-e Progenitors Does Not Require Erythropoietin Or the Erythropoietin Receptor. *Cell* 83, 59-67.
- Xaus,J., Cardo,M., Valledor,A.F., Soler,C., Lloberas,J., and Celada,A. (1999). Interferon gamma induces the expression of p21(waf-1) and arrests macrophage cell cycle, preventing induction of apoptosis. *Immunity* 11, 103-113.
- Xie,H.F., Ye,M., Feng,R., and Graf,T. (2004). Stepwise reprogramming of B cells into macrophages. *Cell* 117, 663-676.
- Yamaguchi,Y., Suda,T., Suda,J., Eguchi,M., Miura,Y., Harada,N., Tominaga,A., and Takatsu,K. (1988). Purified Interleukin-5 Supports the Terminal Differentiation and Proliferation of Murine Eosinophilic Precursors. *Journal of Experimental Medicine* 167, 43-56.
- Yamaguchi,Y., Ackerman,S.J., and Suda,T. (1998). C/EBP beta and GATA-1 synergistically regulate activity of the eosinophil granule protein promoter: Implication for C/EBP beta activity in eosinophil gene expression. *Blood* 92, 166A.
- Yamaguchi,Y., Nishio,H., Kishi,K., Ackerman,S.J., and Suda,T. (1999). C/EBP beta and GATA-1 synergistically regulate activity of the eosinophil granule major basic protein promoter: Implication for C/EBP beta activity in eosinophil gene expression. *Blood* 94, 1429-1439.
- Yamanaka,R., Barlow,C., LekstromHimes,J., Castilla,L.H., Liu,P.P., Eckhaus,M., Decker,T., WynshawBoris,A., and Xanthopoulos,K.G. (1997). Impaired granulopoiesis, myelodysplasia, and early lethality in CCAAT/enhancer binding protein epsilon-deficient mice. *Proceedings of the National Academy of Sciences of the United States of America* 94, 13187-13192.
- Yamanaka,R., Lekstrom-Himes,J., Barlow,C., Wynshaw-Boris,A., and Xanthopoulos,K.G. (1998). CCAAT enhancer binding proteins are critical components of the transcriptional regulation of hematopoiesis (Review). *International Journal of Molecular Medicine* 1, 213-221.
- Yang,V.W., Katz,J.P., Kaestner,K.H., and Chen,X.M. (2002). Transcriptional profiling of genes regulated by gut-enriched Kruppel-like factor (GKLF or KLF4) during DNA damage. *Gastroenterology* 122, A236.
- Yang,Y.Z., Goldstein,B.G., Chao,H.H., and Katz,J.P. (2005). KLF4 and KLF5 regulate proliferation, apoptosis and invasion in Esophageal cancer cells. *Cancer Biology & Therapy* 4, 1216-1221.
- Yang,Y.Z., Goldstein,B.G., Chao,H.H., and Katz,J.P. (2005). Klf4 and Klf5 play critical and complementary roles in esophageal epithelial proliferation and differentiation. *Gastroenterology* 128, A356.
- Yang,Y.Z., Goldstein,B.G., Chao,H.H., and Katz,J. (2005). KLF4 and KLF5 are regulators of proliferation, differentiation, and apoptosis in esophageal cancer cells. *Gastroenterology* 128, A602.
- Yaroslavskiy,B., Watkins,S., Donnenberg,A.D., Patton,T.J., and Steinman,R.A. (1999). Subcellular and cell-cycle expression profiles of CDK-inhibitors in normal differentiating myeloid cells. *Blood* 93, 2907-2917.

## References

- Yeager,A.M., Levin,J., and Levin,F.C. (1983). The Effects of 5-Fluorouracil on Hematopoiesis - Studies of Murine Megakaryocyte-Cfc, Granulocyte-Macrophage-Cfc, and Peripheral-Blood Cell Levels. *Experimental Hematology* 11, 944-952.
- Yet,S.F., McA'Nulty,M.M., Folta,S.C., Yen,H.W., Yoshizumi,M., Hsieh,C.M., Layne,M.D., Chin,M.T., Wang,H., Perrella,M.A., Jain,M.K., and Lee,M.E. (1998). Human EZF, a Kruppel-like zinc finger protein, is expressed in vascular endothelial cells and contains transcriptional activation and repression domains. *Journal of Biological Chemistry* 273, 1026-1031.
- Yoon,H.S., Chen,X.M., and Yang,V.W. (2003). Kruppel-like factor 4 mediates p53-dependent G(1)/S cell cycle arrest in response to DNA damage. *Journal of Biological Chemistry* 278, 2101-2105.
- Yoon,H.S. and Yang,V.W. (2004). Requirement of Kruppel-like factor 4 in preventing entry into mitosis following DNA damage. *Journal of Biological Chemistry* 279, 5035-5041.
- Yoon,H.S. and Yang,V.W. (2004). Kruppel-like factor 4 (KLF4) prevents entry into mitosis following DNA damage by suppressing cyclin B1 gene expression. *Gastroenterology* 126, A7.
- Yoon,H.S., Ghaleb,A.M., Nandan,M.O., Hisamuddin,I.M., Dalton,W.B., and Yang,V.W. (2005). Kruppel-like factor 4 prevents centrosome amplification following gamma-irradiation-induced DNA damage. *Oncogene* 24, 4017-4025.
- Yoshida,T., Ikuta,K., Sugaya,H., Maki,K., Takagi,M., Kanazawa,H., Sunaga,S., Kinashi,T., Yoshimura,K., Miyazaki,J., Takaki,S., and Takatsu,K. (1996). Defective B-1 cell development and impaired immunity against *Angiostrongylus cantonensis* in IL-5R alpha-deficient mice. *Immunity* 4, 483-494.
- Yu,C.N., Cantor,A.B., Yang,H., Browne,C., Wells,R.A., Fujiwara,Y., and Orkin,S.H. (2002). Targeted deletion of a high-affinity GATA-binding site in the GATA-1 promoter leads to selective loss of the eosinophil lineage in vivo. *Journal of Experimental Medicine* 195, 1387-1395.
- Zafarana,G., Rottier,R., Grosveld,F., and Philipsen,S. (2000). Erythroid overexpression of C/EBP gamma in transgenic mice affects gamma-globin expression and fetal liver erythropoiesis. *Embo Journal* 19, 5856-5863.
- Zhan,Y.F., Lieschke,G.J., Grail,D., Dunn,A.R., and Cheers,C. (1998). Essential roles for granulocyte-macrophage colony-stimulating factor (GM-CSF) and G-CSF in the sustained hematopoietic response of Listeria monocytogenes - Infected mice. *Blood* 91, 863-869.
- Zhang,D.E., Zhang,P., Wang,N.D., Hetherington,C.J., Darlington,G.J., and Tenen,D.G. (1997). Absence of granulocyte colony-stimulating factor signaling and neutrophil development in CCAAT enhancer binding protein alpha-deficient mice. *Proceedings of the National Academy of Sciences of the United States of America* 94, 569-574.
- Zhang,P., Zhang,X.B., Iwama,A., Yu,C.N., Smith,K.A., Mueller,B.U., Narravula,S., Torbett,B.E., Orkin,S.H., and Tenen,D.G. (2000). PU.1 inhibits GATA-1 function and erythroid differentiation by blocking GATA-1 DNA binding. *Blood* 96, 2641-2648.
- Zhang,W., Chen,X., Kato,Y., Evans,P.M., Yuan,S.B., Yang,J., Rychahou,P.G., Yang,V.W., He,X., Evers,B.M., and Liu,C.M. (2006). Novel cross talk of Kruppel-like factor 4 and beta-catenin regulates normal intestinal homeostasis and tumor repression. *Molecular and Cellular Biology* 26, 2055-2064.
- Zhang,W.Q., Shields,J.M., Sogawa,K., Fujii-Kuriyama,Y., and Yang,V.W. (1998). The gut-enriched Kruppel-like factor suppresses the activity of the CYP1A1 promoter in an Sp1-dependent fashion. *Journal of Biological Chemistry* 273, 17917-17925.
- Zhang,W.Q., Geiman,D.E., Shields,J.M., Dang,D.Y.T., Mahatan,C.S., Kaestner,K.H., Biggs,J.R., Kraft,A.S., and Yang,V.W. (2000). The cut-enriched Kruppel-like factor (Kruppel-like factor 4) mediates the transactivating effect of p53 on the p21(WAF1/Cip1) promoter. *Journal of Biological Chemistry* 275, 18391-18398.
- Zhao,W.D., Hisamuddin,I.M., Nandan,M.O., Babbin,B.A., Lamb,N.E., and Yang,V.W. (2004). Identification of Kruppel-like factor 4 as a potential tumor suppressor gene in colorectal cancer. *Oncogene* 23, 395-402.
- Zheng,H., Fletcher,D., Kozak,W., Jiang,M.H., Hofmann,K.J., Conn,C.A., Soszynski,D., Grabiec,C., Trumbauer,M.E., Shaw,A., Kostura,M.J., Stevens,K., Rosen,H., North,R.J., Chen,H.Y., Tocci,M.J., Kluger,M.J., and Vanderploeg,L.H.T. (1995). Resistance to Fever Induction and Impaired Acute-Phase Response in Interleukin-1-Beta-Deficient Mice. *Immunity* 3, 9-19.
- Zhou,X., Schmidtke,P., Zepp,F., and Meyer,C.U. (2005). Boosting interleukin-10 production: therapeutic effects and mechanisms. *Curr. Drug Targets Immune Endocr. Metabol. Disord.* 5, 465-475.

## References

Zhou,Y., McLane,M., and Levitt,R.C. (2001). Th2 cytokines and asthma. Interleukin-9 as a therapeutic target for asthma. *Respir. Res.* 2, 80-84.

Zwerina,J., Redlich,K., Schett,G., and Smolen,J.S. (2005). Pathogenesis of rheumatoid arthritis: Targeting cytokines. *Autoimmune Diseases and Treatment: Organ-Specific and Systemic Disorders 1051*, 716-729.