

7. Literaturverzeichnis

Adler, A.I., Stevens R.J., Manley S.E. et al. on behalf of the UKPDS Group (2003):
Development and progression of nephropathy in type 2 diabetes: The United Kingdom
Prospective Diabetes Study (UKPDS 64).

Kidney Int. 63: 225-232.

Allen, R., Nissenson, M.D., Brian, J.G. et al. (2001):
Prevalence and characteristics of individuals with chronic kidney disease in a large health
maintenance organization.
Am. J. Kidney Dis. 37: 1177-1183.

Andersen, S., Tarnow, L., Rossing, P. et al. (2000):
Renoprotective effects of angiotensin II receptor blockade in type 1 diabetic patients with
diabetic nephropathy.
Kidney Int. 57: 601-606.

Bakris, GL. (1998):
Progression of diabetic nephropathy. A focus on arterial pressure level and methods of
reduction.
Diabetes Res.Clin. Pract. 39: 35-42.

Bakris, G.L., Whelton, P., Weir, M., et al. (2000):
The future of clinical trials in chronic renal disease: outcome of an NIH/FDA/Physician
Specialist Conference. Evaluation of Clinical Trial Endpoints in Chronic Renal Disease Study
Group.
J. Clin. Pharmac. 40 (8): 815-825.

Benecke-Timp, A., Hermann, R., Timp, D. et al. (1995):
Krankheitsspezifische Kontrollüberzeugungen und deren Auswirkungen auf die Belastung bei
Typ II-Diabetikern.
Praxis der Klinischen Verhaltensmedizin und Rehabilitation 29: 55-59.

Berger, M. (1995):
Patientenschulung als Grundlage der Therapie.
Diabetes mellitus 295-308.

Bergström, K., Barany, P., Holm, I. (1999):
An educational programme for persistent life-style changes in patients with chronic renal
disease.
EDTNA ERCA J. 25 (4): 42-44.

Bertram, L. (2002):
K/DOQI. Clinical Practice Guidelines for Chronic Kidney Disease: Evaluation, Classification,
and Stratification.
Am. J. Kidney Dis. 39 (1): 17-31.

Binik, Y.M., Dervis, G.M., Barré, P.E. et al. (1993):
Live and Learn: Patient education delays the need to initiate renal replacement therapy in ESR
diseases.
J. Nerv. Mental Dis. 181: 371-376.

Bleichardt, G. (2000):
Selbstmanagement in der Patientenschulung.
Praxis, Klinische Verhaltensmedizin und Rehabilitation 51: 31-34.

Bommer, J. (2002):
Prevalence and socio-economic aspects of chronic kidney disease.
Nephrol. Dial. Transplant. 17 (11): 8-12.

- Bonnet, F., Deprele, C., Sassolas, A. et al. (2001):
Excessive body weight as a new independent risk factor for clinical and pathological progression in primary IgA nephritis.
Am. J. Kidney Dis. 37 (4): 720-727.
- Brenner, B.M. (1983):
Hemodynamically mediated glomerular injury and the progressive nature of kidney disease.
Kidney Int. 23: 647.
- Brenner, B.M., Cooper, M.E., de Zeeuw, D. et al. (2001):
Effects of Losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy.
N. Engl. J. Med. 345 (12): 861-869.
- Brian, A.J., Walters, Ph.D., Ron, D. et al. (2002):
Health-related quality of life, depressive symptoms, anemia and malnutrition at hemodialysis initiation.
Am. J. Kidney Dis. 40 (6): 1185-1194.
- Bullinger, M. (1996):
Assessment of health related quality of life with the SF-36 Health Survey.
Rehabilitation 35 (3): 17-27.
- Bundesarbeitsgemeinschaft für Rehabilitation (1993):
Arbeitshilfe für die Rehabilitation von chronischen Nierenkranken, Dialysepflichtigen und Nierentransplantierten.
In: Schriftenreihe der Bundesarbeitsgemeinschaft für Rehabilitation, Frankfurt a.M. 10: 19-20.
- Bundesversicherungsanstalt für Angestellte (BfA): Sozialgesetzbuch IX, Berlin 2002.

Cirillo, M., Stellato, D., Laurenzi, M. et al. (2000):
Pulse pressure and isolated systolic hypertension: Association with microalbuminuria.
Kidney Int. 58: 1211-1218.

Cleveland, D.R., Kailash, K.J., Hirsch, D.J. et al. (2002):
Quality of prereferral care in patients with chronic renal insufficiency.
Am. J. Kidney Dis. 40 (1): 30-36.

Cockcroft, D.W., Gault, M.H. (1976):
Prediction of creatinine-clearance from serum creatinine.
Nephron 16: 31-34.

Daugirdas, J.T. (1993):
Second generation logarithmic estimates of single-pool variable volume Kt/V: an analysis of error.
J. Am. Soc. Nephrol. 4: 1205-1213.

Daul, A.E., Völker, K., Alberty, A. et al. (1990):
Dialysesportgruppe: Eine Möglichkeit zur Verbesserung der körperlichen Leistungsfähigkeit und der psycho-sozialen Rehabilitation chronischer Dialysepatienten.
Nieren- und Hochdruckkrankheiten 6: 275-284.

The Diabetes Control and Complications (DCCT) Research Group (1995):
Effect of intensive therapy on the development and progression of diabetic nephropathy in the Diabetes Control and Complications Trial.
Kidney Int. 47: 1703-1720.

Decimus Junius Juvenalis (140 n. Christi), (Übersetzung, 1984):
Satire X: De Ijdelheid van het menselijk streven.
In: d'Hane-Skeltema, M., De Satiren. Amsterdam:
Ambo BV, Baaren, Athenaeum-Polak & Van Gennep. 142.

Deutsche Hochdruckliga, Deutsche Hypertonie Gesellschaft
Empfehlungen zur Hochdruckbehandlung
17. Auflage – Heidelberg, November 2002

Devins, G.M., Hollomby, D.J., Barre, P.E. et al. (2000):
Long-term knowledge retention following predialyses psychoeducational intervention.
Nephron 86 (2): 129-134.

Dwyer, J.T., Larive, B., Leung, J. et al. (2002):
Nutritional status affects quality of life in hemodialysis study patients at baseline.
J. Ren. Nutr. 12 (4): 213-223.

Erben, K. (1997):
Trend zu längerer Dialysezeit. Nierenersatztherapie in Amerika.
Der Dialysepatient 1: 80-81.

Evans, R.U. (1990):
Quality of life assessment and the treatment of end-stage renal diseases.
Transplantation 4 (1): 28-51.

Faltermaier, T. (1996):
Nicht nur Compliance, sondern gleichberechtigte Interaktion.
Der Dialysepatient 2: 79-80.

Falvo, R. (1995).
Educational evaluation: what are the outcomes?
Adv. Renal Repl. Ther. 2 (3): 227-233.

Franklin, S.S. et al. (2001):
Does the relation of blood pressure to coronary heart disease risk change with aging?
Circulation 103: 1245-1249.

Frei, U., Schober-Halstenberg, H.J. (2002):

Nierenersatztherapie in Deutschland. Bericht über Dialysebehandlung und Nierentransplantation in Deutschland.

Quasi-Niere

Friend, et al. (1989):

Survival advantage for hemodialysis patients.

Am. J. Publ. Health. 76; 670-672.

Fried, L.F., Orchard, T.J., Kasiske, B.L. for the lipids and renal disease progression meta-analysis study group (2001):

Effect of lipid reduction on the progression of renal disease: A meta-analysis.

Kidney Int. 59: 260-269.

Fritschka, E., Bennis, M., Fangmann, C., et al. (1991):

Profiles of blood pressure and endothelin after kidney transplantation.

J. Vasc. Res. 29: 115.

Fritschka, E., Samtleben, W. (1998):

Rehabilitation bei nephrologischen Erkrankungen.

In: Delbrück H, Haupt E. Rehabilitationsmedizin. München, Wien, Baltimore:
Urban und Schwarzenberg. 535-561.

Fritschka, E., Mahlmeister, J., Liebscher-Steinecke, R. et al. (2000):

Rehabilitation bei Patienten mit chronischer Niereninsuffizienz, Dialysepatienten und nach Nierentransplantation.

Nieren- und Hochdruckkrankheiten 29 (11): 555-565.

Garg, A.X., Clark, W.F., Haynes, R.B. et al. (2002):

Moderate renal insufficiency and the risk of cardiovascular mortality: Results from the NHANES I.

Kidney Int. 61: 1486-1494.

Gerdes, N., Jäckel, W.H. (1992):

„Indikatoren des Reha-Status (IRES)“ – ein Fragebogen zur Beurteilung von Rehabilitationsbedürftigkeit und –erfolg.

Rehabilitation. 31; 73-79.

Ghossein, C., Serrano, A., Rammohan, M. et al. (2002):

The role of comprehensive renal clinic in chronic kidney disease stabilization and management: The Northwestern experience.

Semin. Nephrol. 22 (6): 526-532.

The Gisen Group. (1997):

Randomisierte, placebokontrollierte Untersuchung zur Wirkung von Ramipril (Delix) auf den Rückgang der glomerulären Filtrationsrate und das Risiko eines terminalen Nierenversagens bei Patienten mit nicht-diabetischer Nephropathie (REIN-Studie).

The Lancet 349: 1857-1863.

Goldberg, A.P., Geltman, E.M., Gavin, J.R. et al. (1986):

Exercise training reduces coronary risk and effectively rehabilitates hemodialysis patients.

Nephron 42: 311-316.

Golper, T. (2001):

Patient education: can it maximaize the succes of therapy?

Nephrol. Dial. Transplant. 16 (Suppl. 7): 20-24.

Grüßer, M., Jörgens, V. (1996):

Hypertonie- Behandlungs- und Schulungsprogramm.

Deutscher Ärzte-Verlag, Kirchheim, Mainz.

Hall, J.E., Brandy, M.W., Henegar, J.R. et al. (1998):

Abnormal kidney function as a cause and a consequence of obesity hypertension.

Clin. Exp. Pharmacol. Physiol. 25: 58-64.

Hannedouche, T., Chauveau, P., Kalou, F. et al. (1993):
Factors affecting progression in advanced chronic renal failure.
Clin. Nephrol. 39: 312-320.

Hansen, H.P., Tauber-Lassen, E., Jensen, B.R. et al. (2002):
Effect of dietary protein restriction on prognosis in patients with diabetic nephropathy.
Kidney Int. 62: 220-228.

Hariprasad, S. et al. (2002):
Slowing the progression of chronic renal failure: economic benefits and patient's
perspectives.
Am. J. Kidney Dis. 39 (4): 721-729.

Harris, L.E., Luft, F.C., Rudy, D.W. et al. (1993):
Clinical correlates of functional status in patients with chronic renal insufficiency.
Am. J. Kidney Dis. 21: (2): 161-166.

Hayslip, D.M., Suttle, C.D. (1995):
Pre-ESRD patient education: A review of the literature.
Adv. Renal Repl. Ther. 2 (3): 217-226.

Heise, T., Jennen, E., Hochlenert, D. et al. (2002):
Results of a structured hypertension teaching and treatment programme.
Z. Gastroenterol. 40 (Suppl. 1): 27-32.

Higashi, Y., Oshima, T., Sasaki, S. et al. (1998):
Angiotensin-converting-enzyme inhibition, but not calcium antagonism, improves a response
of the renal vasculature to L-arginine in patients with essential hypertension.
Hypertension 32: 16-24.

Hostetter, T.H., Lising, M. (2002):
National kidney disease education program.
J. Natl. Med. Assoc. 94(8): 72-75.

Hovind, P., Rossing, P., Tarnow, L. et al. (2001):
Progression of diabetic nephropathy.
Kidney Int. 59: 702-709.

Huber, W., Kettner, A., Höffken, B. et al. (1981):
Berufliche Rehabilitation und Begutachtung von Niereninsuffizienten.
Medizinische Welt 32 (22): 880-883.

Huber, W., Kettner, A., Ritz, E. (1982):
Rehabilitation bei chronischer Niereninsuffizienz.
In: Losse, H., Renner, E.: Klinische Nephrologie 2
Thieme, Stuttgart. 518-528.

Huber, W., Roth, G., Göbel, M. (1997):
Soziale und berufliche Wiedereingliederung und Erwerbsfähigkeit bei chronischer
Niereninsuffizienz.
In: Franz, H.E., Hörl, W.H.: Blutreinigungsverfahren. 31; 373-389.
Thieme, Stuttgart, New York.

Ignacio-Garcia, J.M., Pinto-Tenorio, M., Chocron-Giraldez, M.J. et al. (2002):
Benefits after 3yrs of an asthma education programme coupled with regular reinforcement.
Eur. Respir. J. 20 (5): 1095-10101.

Iseki, K., Ikemiya, Y., Fukiyama, K. (1997):
Predictors of end-stage renal disease and body mass index in a screened cohort.
Kidney Int. 52: 169-170.

Ismail, N., Neyra, R., Hakim, R. (1998):

The medical and economical advantages of early referral of chronic renal failure patients to renal specialists.

Nephrol. Dial. Transplant. 13: 246-250.

Jäckel, W.H., Gerdes, N., Herdt, J. et al. (2002):

Wissensmanagement in der Rehabilitation-Vorschlag zu einer systematischen Entwicklung von Leitlinien.

Rehabilitation 41: 217-225.

Jafar, Th., Schmid, C.H., Landa, M. et al. for the ACE-inhibition in progressive renal disease study group (2001):

Angiotensin-converting enzyme inhibitors and progression of non-diabetic renal disease. A meta-analysis of patient-level data.

Ann. Intern. Med. 135: 73-87.

Jones, C.A., McQuilan, G.M., Kusek, J.W. et al. (1998):

Serum creatinine levels in the US population: Third National health and Nutrition Examination Survey.

Am. J. Kidney Dis. 32: 992-999.

Jungers, P., Skhiri, H., Zingraff, J. et al. (1997):

Benefits of early nephrological management in chronic renal failure.

Presse Med. 26: 1325-1329.

Jungers, P., Massy, Z.A., Nguyen, T. et al. (2001):

Longer duration of predialysis nephrological care is associated with improved long-term survival of dialysis patients.

Nephrol. Dial. Transplant. 16: 2357-2364.

Kambham, N., Markowitz, G., Valeri, A.M. et al. (2001):
Obesity-related glomerulopathy: an emerging epidemic.
Kidney Int. 59: 1498-1509.

Kaplan DeNour, A., Czaczkes, W. (1976):
The influence of patients personality on adjustment to chronic dialysis.
J. Nerv. Ment. Dis. 162 (5): 323-333.

Kaplan DeNour, A., Brickman, A. (1996):
Determining quality of life and pharmacoconomics.
Clinical Trials 2: 953-960.

Kaplan DeNour, A., Brickman, A. (1997):
Depression bei terminaler Niereninsuffizienz.
In: Franz, H.E., Hörl, W.H.: Blutreinigungsverfahren.
Thieme, Stuttgart, New York. 366-372.

Karley, M.L. (1998):
The use of group process in the education of renal patients and families: A multi-centred collaborative approach.
CANNT 22: 8-4.

Khan, S.S., Kazmi, W.H., Abichandani, R., et al. (2002):
Health care utilization among patients with chronic kidney disease.
Kidney Int. 62: 229-236.

Klag, M.J., Whelton, P.K., Randall, B.L. et al. (1996):
Blood pressure and end-stage renal disease in men.
N. Engl. J. Med. 334: 13-18.

Klahr, S., Levey, A.S., Beck, G.J. et al. (1994):
The effects of dietary protein restriction and blood-pressure control on the progression of chronic renal disease (MDRD).
N. Engl. J. Med. 330 (29): 877-884.

Klang, B., Björvell, H., Berglund, J. et al. (1998):
Predialysis patient education: effects on functioning and well-being in uremic patients.
J. Adv. Nurs. 28 (1): 36-44.

Knäuper, B., Schwarzer, R. (2000):
Selbstwirksamkeitserwartungen in der Patientenschulung.
Praxis klinische Verhaltensmedizin und Rehabilitation 51: 5-10.

Konstantinidou, E., Koukouvou, G., Kouidi, E. et al, (2002):
Exercise training in patients with end-stage renal disease on hemodialysis: comparison of three rehabilitation programs.
J. Rehabilitation Medicin. 34 (1): 40-45.

Kopple, J.D. (2001):
National Kidney Foundation K/DOQI clinical practice guidelines for nutrition in chronic renal failure.
Am. J. Kidney Dis. 37 (1): 66-70.

Kotchen, T.A., Piering, A.W., Cowley, A.W. et al. (2000):
Glomerular hyperfiltration in hypertensive African-Americans.
Hypertension 35 (3): 822-826.

Krämer, B.K., Wiecek, A., Ritz, E. (1997):
Womit und wie intensiv soll der Blutdruck bei diabetischer Nephropathie gesenkt werden.
DMW. 22: 829-832.

Kutner, N.G., Cardenas, D.D. (1982):

Assessment of rehabilitation outcomes among chronic dialysis patients.

Am. J. Nephrol. 2: 128-132.

Kutner, N.G., Lin, L.S., Fielding, B. et al. (1994):

Continued survival of older hemodialysis patients: Investigation of psychosocial predictors.

Am. J. Kidney Dis. 24 (1): 42-49.

Kurtner, G.K., Curtin, R.B., Oberley, E. et al. (1997):

Fulfilling the promise: Linking rehabilitation interventions with ESRD patients outcomes.

Dial. Transplant. 26 (5): 282-292.

Ladwig, K.H. (1995):

Verhaltensmedizinische Grundsätze und Motivationsstrategien für die Prävention.

Bayerisches Ärzteblatt 12: 530-533.

Lam, K.S.L., Cheng, I.K.P. et al. (1995):

Cholesterol-lowering therapy may retard the progression of diabetic nephropathy.

Diabetologica 38: 604-609.

Latham, C.E. (1998):

Is there data to support the concept that educated, empowered patients have better outcomes?

J. Am. Soc. Nephrol. 9 (12): 141-144.

Ledermann, H., Schaffner, O. (1995):

Erhebung über den Schulungsstand stationärer Diabetespatienten.

Präv.-Rehab. 7 (4): 206-208.

Lemozy-Cadroy, S., Cognier, S., Gourdy, P. et al. (2002):

Intensified treatment of type 1 diabetes: prospective evaluation at one year of a therapeutic patient education programme.

Diabetes Metab. 28 (4): 287-294.

Levey, A.S., Adler, S., Caggiula, A.W. et al. (1996):
Effects of dietary protein restriction on the progression of advanced renal disease in the
modification of diet in Renal Disease Study.
Am. J. Kidney Dis. 27: 652-663.

Levin, A., Lewis, M., Mortiboy, P. et al. (1997):
Multidisciplinary predialysis programs: quantification and limitations of their impact on
patient outcomes in two Canadian settings.
Am. J. Kidney Dis. 29 (4): 533-540.

Lewis, E.J., Hunsicker, L.G., Bain, et al. (1993):
The effect of angiotensin-converting- enzyme inhibition on diabetic nephropathy.
N. Engl. J. Med. 329 (20): 1456-1462.

Lewis, E.J., Hunsicker, L.G., Clarke, W.R., et al. (2001):
Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with
nephropathy due to type 2 diabetes.
N. Engl. J. Med. 345: 851-860.

Ligttenberg, G. Blankestijn, P.J., Oey, P.L. et al. (1999):
Reduction of sympathetic hyperactivity by enalapril in patients with chronic renal failure.
N. Engl. J. Med. 340: 1321-1328.

Locatelli, F., Carbarns, I.R.I., Maschio, G. et al. (1997):
Long-term progression of chronic renal insufficiency in the AIPRI Extension Study.
Kidney Int. 52: 63-66.

Locatelli, F., Del Vecchio, L. (1999):
How long can dialysis be postponed by low protein diet and ACE-inhibitors?
Nephrol. Dial. Transplant. 14: 1360-1364.

Locatelli, F., Del Vecchio, L., Andrulli, S. et al. (2000):
The role of underlying nephropathy in the progression of renal disease.
Kidney Int. 57: 49-55.

Locatelli, F., Del Vecchio, L., Pozzoni, P. (2002):
The importance of early detection of chronic kidney disease.
Nephrol. Dial. Transplant. 17: 2-7.

London, G.M., Marchais, S.J., Guérin, A.P. et al. (2002):
Impairment of arterial function in chronic renal disease: prognostic impact and therapeutic approach.
Nephrol. Dial. Transplant. 17: 13-15.

Maddox, D.A., Alavi, F.K., Santella, R.N. et al. (2002):
Prevention of obesity-linked renal disease: Age-dependent effects of dietary food restriction.
Kidney Int. 62: 208-219.

Massy, Z.A., Jennie, Z.Ma., Louis, T.A. et al. (1995):
Lipid-lowering therapy in patients with renal disease.
Kidney Int. 48: 188-198.

Muthny, F.A., Koch, U., Köster, W. et al. (1984):
Psychosoziale Reaktionen auf chronische Niereninsuffizienz und Dialyse – eine empirische Untersuchung.
Nieren- und Hochdruckkrankheiten 13; 9: 349-355.

Muthny, F.A., Koch, U., Kösters, W. et al. (1984):
Psychosoziale Reaktion auf chronische Niereninsuffizienz und Dialyse- eine empirische Untersuchung.
Nieren- und Hochdruckkrankheiten 13 (9): 349-355.

Muthny, F.A., Koch, U., Jontofsohn, R. et al. (1986):
Medizinisches Wissen, Compliance und Behandlungszufriedenheit von Dialysepatienten.
Nieren- und Hochdruckkrankheiten 15 (5): 187-196.

Muthny, F.A. (1994):
Psychosoziale Probleme bei der chronischen Niereninsuffizienz und Behandlungsmöglichkeiten.
In: Brass, H., Philipp, Th., Schulz, W.: Manuale nephrologicum.
Dustri, Deisenhofen. IX-4: 1-12.

Nebel, M. (2002):
Behandlungskosten der Nierenersatztherapie in Deutschland 1999.
Nieren- und Hochdruckkrankheiten 31 (3): 85-92.

Nissenson, A.R., Pereira, B.J., Collins, A.J. et al. (2001):
Prevalence and characteristics of individuals with chronic kidney disease in a large health maintenance organization.
Am. J. Kidney Dis. 37 (6): 1177-1183.

Nitz, J., Shayman, D. (1986):
A model for patient education.
ANNA J. 13 (5): 253-255.

Painter, P. (1994):
The importance of exercise training in rehabilitation of patients with end-stage renal disease.
Am. J. Kidney Dis. 24 (1): 52-59.

Parsons, D.S., Harris, D.C. (1997):
A review of quality of life in chronic renal failure.
Pharmacoeconomics 12 (2): 140-160.

Parving, H., Schmidt, U.M., Hommel, E. et al. (1993):
Effective antihypertensive treatment postpones renal insufficiency in diabetic nephropathy.
Am. J. Kidney Dis. 22 (1): 188-195.

Patel, S.S., Shah, V.S., Peterson, R.A. et al. (2002):
Psychosocial variables, quality of life, and religious beliefs in ESRD patients treated with
hemodialysis.
Am. J. Kidney Dis. 40 (5): 1013-1022.

Pedrini, M.T., Levey, A.S., Lau, J. et al. (1996):
The effect of dietary protein restriction on the progression of diabetic and nondiabetic renal
disease.
Ann. Intern. Med. 124: 627-632.

Pereira, BJ. (2002):
Overcoming barriers to the early detection and treatment of chronic kidney diseases and
improving outcomes for end-stage renal disease.
Am. J. Management Care. 8 (4): 122-135.

Posavac, E.J. (1980):
Evaluations of Patient Education Programms.
Evaluation & The Health Professions. 3; 1, 47-62.

Praga, M., Hernandez, E., Andres, A. et al. (1995):
Effects of body-weight loss and captopril treatment on proteinuria associated with obesity.
Nephron 70: 35-41.

Praga, M., Hernandez, E., Morales, E. et al. (2001):
Clinical features and long-term outcome of obesity-associated focal segmental
glomerulosclerosis.
Nephrol. Dial. Transplant. 16: 1790-1798.

Praga, M. (2002):

Obesity- a neglected culprit in renal disease.

Nephrol. Dial. Transplant. 17: 1157-1159.

Price, V., Archbold, J. (1995):

Development and application of social learning theory.

Br. J. Nurs. 4 (21): 1263-1268.

Rasgon, S., Schwankovsky, L., James-Rogers, A. et al. (1993):

An intervention for employment maintenance among blue-collar workers with end-stage renal disease.

Am. J. Kidney Dis. 22 (3): 403-412.

Rachmani, R., Levi, Z., Slavachevski, I. et al. (2002):

Teaching patients to monitor their risk factors retards the progression of vascular complications in high-risk patients with type 2 diabetes mellitus – a randomized prospective study.

Diabet. Med. 19 (5): 385-392.

Ravid, M., Neumann, L., Lishner, M. (1995):

Plasma lipids and the progression of nephropathy in diabetes mellitus type II: Effect of ACE-inhibitors.

Kidney Int. 47: 907-910.

Röseler, E., Aurisch, R., Strangfeld, D. et al. (1987):

Diagnostik leistungsvermindernder Veränderungen bei chronisch Nierenkranken.

Nieren- und Hochdruckkrankheiten 16 (1): 7-16.

Röseler, E., Strangfeld, D., Scholz, D. et al. (1997):

Die körperliche Leistungsfähigkeit chronisch hämodialysierter Männer und Frauen.

Deutsches Gesundheitswesen 32: 1408-1412.

- Röseler, E., Strangfeld, D., Aurisch, R. et al. (1998):
Das Verhalten der körperlichen Leistungsfähigkeit chronisch nierenkranker Männer in verschiedenen Stadien der chronischen Niereninsuffizienz.
Deutsches Gesundheitswesen 33: 977-982.
- Ruggenenti, P., Perna, A., Remuzzi, G. (2001):
ACE-inhibitors to prevent end-stage renal disease: when to start and why possibly never to stop: a post hoc analysis of the REIN trial results. Ramipril Efficacy in Nephropathy.
J. Am. Soc. Nephrol. 12 (12): 2832-2837.
- Ruilope, L.M., Rodicio, J.L. (2001):
The kidney in arterial hypertension.
Nephrol. Dial. Transplant. 16 (Suppl 1): 50-52.
- Sacks, C.R., Peterson, R.A., Kimmel, P.L. (1990):
Perception of illness and depression in chronic renal disease.
Am. J. Kidney Dis. 15 (1): 31-39.
- Salahudeen, A.K. (2002):
Halting progression of renal failure: consideration beyond angiotensin II inhibition.
Nephrol. Dial. Transplant. 17 (11): 1871-1875.
- Samtleben, W., Fritschka, E. (1998):
Rehabilitation bei nephrologischen und urologischen Erkrankungen.
In: Delbrück und Haupt: Rehabilitationsmedizin
Urban & Schwarzenberg. 535-561.
- Schmidt, J., Lamprecht, F., Wittmann, W.W. (1989):
Zufriedenheit mit der stationären Versorgung. Entwicklung eines Fragebogens und erste Validitätsuntersuchungen.
Psychother. med. Psychol. 39: 248-255.

Schumann, V., Fritschka, E., Goepel, M. et al. (1994):
Long-term renal function after unilateral nephrectomy.
Nieren- und Hochdruckkrankheiten 23 (3): 125-131.

Slowik, M.M. (2001):
Early education of patients with chronic renal insufficiency: the healthy start program. Case study of the anemic patient.
J. Nephrol. Nurs. 28 (6): 643-646.

Stephenson, R.N., Villano, R. (1993):
Results of a predialysis education program.
Dial. Transplant. 566-571.

Taal, M.W., Brenner, B.M. (2001):
Achieving maximal renal protection in nondiabetic chronic renal disease.
Am. J. Kidney Dis. 38: 1365-1372.

Tazeen, H., Stark, P.C., Schmid, C.H. (2001):
Proteinuria as a modifiable risk faktor for the progression of non-diabetic renal disease.
Kidney Int. 60: 1131-1140.

Tews, H.P., Schreiber, W.K., Huber, W. et al. (1980):
Vocational rehabilitation in dialysed patients.
Nephron 26: 130-136.

Thurns, S., Schliehe, F. (2000):
Struktur und Organisation der Rehabilitation.
In: Gostomzyk, J.G.: Angewandte Sozialmedizin. Handbuch für Weiterbildung und Praxis.
Ecomed, Kaufbeuren.

Toto, R.D. (2002):

Hypertension and kidney literature review 2000.

Clin. Nephrol. 58 (4): 253-259.

Trivedi., H.S., Pang, M.M., Campbell, A. et al. (2002):

Slowing the progression of chronic renal failure: economic benefits and patients' perspectives.

Am. J. Kidney Dis. 39 (4): 721-729.

van Hout, B.A., Simeon, G.P., McDonell, J. et al. (1997):

Economic evaluation of benazepril in chronic renal insufficiency.

Kidney Int. 52 (63): 159-162.

Wassermann, K. et al. (1994):

Principles of exercise testing and interpretation.

Lea & Febiger, Philadelphia.

White, C.A., Pilkey, R.M., Lam, M. et al. (2002):

Predialysis clinic attendance improves quality of life among hemodialysis patients.

BMC Nephrol. 5 (3): 3.

Yilmaz, A., Akkaya, E. (2002):

Evaluation of long-term efficacy of an asthma education programme in an out-patient clinic.

Respir. Med. 96 (7): 519-524.