

11 Literature cited

11.1 Accepted and published papers of the author

DIANE N, HH HILGER & M GOTTSCHLING (2002b): Transfer cells in the seeds of Boraginales. *Bot J Linn Soc* **140**: 155–164.

GOTTSCHLING M (2001): Evolutionäre Interpretationen molekularbiologischer Ergebnisse am Beispiel der Ehretiaceae (Boraginales). *Sitzungsber Ges Naturf Freunde Berlin* **40**: 59–73.

GOTTSCHLING M, N DIANE, HH HILGER & M WEIGEND (2002): Von Pflanzen und Geologie. – In: BÖSE M, H KEUPP (eds.): *Der belebte Planet*. Berlin: FU Berlin, FB Geowissenschaften.

GOTTSCHLING M & HH HILGER (2001): Phylogenetic analysis and character evolution of *Ehretia* and *Bourreria* (Ehretiaceae, Boraginales) based on ITS1 sequences. *Bot Jahrb Syst* **123**: 249–268.

GOTTSCHLING M & HH HILGER (in press): First fossil record of transfer cells in angiosperms. *Amer J Bot*

GOTTSCHLING M, HH HILGER, M WOLF & N DIANE (2001): Secondary structure of the ITS1 transcript and its application in a reconstruction of the phylogeny of Boraginales. *Pl Biol* **3**: 629–636.

GOTTSCHLING M, DH MAI & HH HILGER (2002): The systematic position of *Ehretia* fossils (Ehretiaceae, Boraginales) from the European Tertiary and implications for character evolution. *Rev Paleobot Palynol* **121**: 149–156.

HILGER HH & M GOTTSCHLING (in press): Phylogenetic analysis and character evolution of *Ehretia* and *Bourreria* (Ehretiaceae, Boraginales) – correction. *Bot Jahrb Syst*

11.2 Submitted manuscripts of the author

GOTTSCHLING M, HH HILGER, N DIANE & M WEIGEND (subm.): Time estimates and historical biogeography of the Primarily Woody Boraginales: Ehretiaceae, Cordiaceae, and Heliotropiaceae. *Int J Pl Sci*

11.3 Manuscripts of the author in preparation

GOTTSCHLING M & HH HILGER (in prep.): Secondary structure of the *trnL_{UAA}* (group I) intron and the molecular delimitation of Ehretiaceae (Boraginales).

GOTTSCHLING M, HH HILGER, JS MILLER & M WEIGEND (in prep. a): Phylogenetic analysis of Cordiaceae (Boraginales) inferred from ITS1 sequence data.

GOTTSCHLING M, HH HILGER & M WEIGEND (in prep. b): Cordiaceae R.BR. ex DUMORT. – In: KUBITZKI K (ed.): *Families of Flowering Plants*. Berlin: Springer.

11.4 Symposium abstracts of the author

- GOTTSCHLING M & HH HILGER** (2001): Phylogenetic systematics of Ehretiaceae (Boraginales). Biodiversität und Evolutionsbiologie, Bochum (Germany).
- GOTTSCHLING M & HH HILGER** (2001): Fossil fruit morphology and DNA systematics: phylogeny and current distribution of *Ehretia* (Ehretiaceae, Boraginales). Deep Morphology, Vienna (Austria).
- GOTTSCHLING M, N DIANE, M WEIGEND & HH HILGER** (2002): Distribution patterns in Cordiaceae, Ehretiaceae, and Heliotropiaceae (Boraginales). Botany 2002, Madison (Wisconsin, USA).
- GOTTSCHLING M, HH HILGER & N DIANE** (2002): The secondary structure of the ITS1 transcript and its application for reconstruction phylogeny. Botany 2002, Madison (Wisconsin, USA).

11.5 Papers of other authors

- AGARDH JG (1858): *Theoria systematis plantarum*. Lund etc.: Gleerup.
- ALBACH DC, PS SOLTIS, DE SOLTIS & RG OLMSTEAD (2001): Phylogenetic analysis of asterids based on sequences of four genes. *Ann Missouri Bot Gard* **88**: 163–212.
- AL-SHEHBAZ IA (1991): The genera of Boraginaceae in the Southeastern United States. *J Arnold Arbor, Suppl Ser* **1**: 1–65.
- ALTSCHUL SF, TL MADDEN, AA SCHÄFFER, J ZHANG, Z ZHANG, W MILLER & DJ LIPMAN (1997): Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**: 3389–3402.
- AN SS, T FRIEDL & E HEGEWALD (1999): Phylogenetic relationships of *Scenedesmus* and *Scenedesmus*-like coccoid green algae as inferred from ITS-2 rDNA sequence comparison. *Pl Biol* **1**: 418–428.
- ANDREASEN K & BG BALDWIN (2001): Unequal evolutionary rates between annual and perennial lineages of Checker Mallows (*Sidalcea*, Malvaceae): evidence from 18S–26S rDNA Internal and External Transcribed Spacers. *Mol Biol Evol* **18**: 936–944.
- ASKINS RA, KM ERCOLINO & JD WALTER (1987): Flower destruction and nectar depletion by avian nectar robbers on a tropical tree, *Cordia sebestena*. *J Field Ornithol* **58**: 345–349.
- BACON JD (1987): Systematics of *Nama* (Hydrophyllaceae): seed coat morphology of *Lemmonia californica* and *Nama* species allied with *Nama demissum*. *Aliso* **11**: 441–450.
- BACON JD & LH BRAGG (1986): Systematics of *Nama* (Hydrophyllaceae): comparison of seed morphology of sects. *Arachnoidea* and *Cinerascentia* with five species of *Eriodictyon* and *Turricula parryi*. *Sida* **11**: 271–281.
- BALDWIN BG (1992): Phylogenetic utility of the Internal Transcribed Spacers of nuclear ribosomal DNA in plants: an example from the Compositae. *Mol Phylogenet Evol* **1**: 3–16.
- BÖHLE U-R & HH HILGER (1997): Chloroplast DNA systematics of “Boraginaceae” and related families – a goodbye to the old and familiar concept of five subfamilies (Abstract). *Scripta Bot Belg* **15**: 30.
- BÖHLE U-R, HH HILGER & WF MARTIN (1996): Island colonization and evolution of the insular woody habit in *Echium* L. (Boraginaceae). *Proc Natl Acad Sci USA* **93**: 11740–11745.
- BONNEMAIN JL, S BOURQUIN, S RENAULT, C OFFLER & DG FISHER (1991): Transfer cells: structure and physiology. — In: BONNEMAIN JL, S DELROT, WL LUCAS & L DAINTY (eds.): *Recent advances in phloem transport and assimilate compartmentation*. Paris, Nantes: Ouest Edition, Presses Académiques.

- BORHIDI AE, E GONDÁR & ZS OROSZ-KOVÁCS (1988): The re-consideration of the genus *Cordia* L. Acta Bot Hung **34**: 375–423.
- BREMER B, K BREMER, N HEIDARI, P ERIXON, RG OLMSTEAD, AA ANDERBERG, M KÄLLERSJÖ & E BARKHORDARIAN (2002): Phylogenetics of asterids based on 3 coding and 3 non-coding chloroplast DNA markers and the utility of non-coding DNA at higher taxonomical levels. Mol Phylogenet Evol **24**: 274–301.
- BRENNER GJ (1996): Evidence of the earliest stage of angiosperm pollen evolution: a paleoequatorial section from Israel. — In: TAYLOR DW & LJ HICKEY (eds.): Flowering plant origin, evolution and phylogeny. New York: Chapman & Hall.
- BRITTON NL & P WILSON (1925): Scientific Survey of Porto Rico and the Virgin Islands **6**. New York: New York Acadamy of Sciences.
- BROWNE P (1756): The civil and natural history of Jamaica in three parts. London: author.
- CANDOLLE AP de (1845): Boragineæ. — In: CANDOLLE A de (ed.): Prodromus systematis naturalis regni vegetabilis **9**. Paris etc.: Treuttel & Würtz.
- CHANCE GD & JD BACON (1984): Systematic implications of seed coat morphology in *Nama* (Hydrophyllaceae). Amer J Bot **71**: 829–842.
- CHANDLER MEJ (1961): The Lower Tertiary floras of Southern England **1**. London: British Museum.
- CHANDLER MEJ (1962): The Lower Tertiary floras of Southern England **2**. London: British Museum.
- CHANDLER MEJ (1964): The Lower Tertiary floras of Southern England **4**. London: British Museum.
- CHASE MW, DE SOLTIS, RG OLMSTEAD, D MORGAN, DH LES, BD MISHLER, MR DUVALL, RA PRICE, HG HILLS, Y-L QIU, KA KRON, JH RETTIG, E CONTI, J PALMER, JR MANHART, KJ SYTSMA, HJ MICHAELS, WJ KRESS, KG KAROL, WD CLARK, M HEDRÉN, BS GAUT, RK JANSEN, K-J KIM, CF WIMPEE, JF SMITH, GR FURNIER, SH STRAUSS, Q-Y XIANG, GM PLUNKETT, PS SOLTIS, SM SWENSEN, SE WILLIAMS, PA GADEK, JC QUINN, LE EGUIARTE, E GOLENBERG, GH LEARN Jr., SW GRAHAM, SCH BARRETT, S DAYANANDAN & VA ALBERT (1993): Phylogenetics of seed plants: an analysis of nucleotide sequences from the plastid gene *rbcL*. Ann Missouri Bot Gard **80**: 528–580.
- CECH TR (1988): Conserved sequences and structures of group I introns: building an active site for RNA catalysis – a review. Gene **73**: 259–271.
- CECH TR, SH DAMBERGER & RR GUTELL (1994): Representation of the secondary and tertiary structure of group I introns. Nat Struct Biol **1**: 273–280.
- CHELEBAJEVA AI (1984): The genus *Cordia* (Boraginaceae) from the Palaeogene of Kamchatka (Russian SFSR, USSR) and adjacent territories. Bot Zhurn (Moscow & Leningrad) **69**: 605–615.
- CHILTON NB & RB GASSER (1999): Sequence differences in the Internal Transcribed Spacers of DNA among four species of hookworm (Ancylostomatoidea: *Ancylostoma*). Int J Parasitol **29**: 1971–1977.
- CHUANG TI & L CONSTANCE (1992): Seeds and systematics in Hydrophyllaceae: tribe Hydrophylleae. Amer J Bot **79**: 257–264.
- COLEMAN AW & JC MAI (1997): Ribosomal ITS-1 and ITS-2 sequence comparisons as a tool for predicting genetic relatedness. J Mol Evol **45**: 168–177.
- COLEMAN AW, RM PREPARATA, B MEHROTRA & JC MAI (1998): Derivation of the secondary structure of the ITS-1 transcript in Volvocales and its taxonomical correlations. Protist **149**: 135–146.

- COLLINSON ME (1983): Fossil plants of the London Clay. London: Palaeontological Association.
- COSNER ME, RK JANSEN & TG LAMMERS (1994): Phylogenetic relationships in the Campanulales based on *rbcL* sequences. *Pl Syst Evol* **190**: 79–95.
- COSTA GM da, TLG de LEMOS, ODL PESSOA, FJQ MONTE & R BRAZ-FILHO (1999): Glaziovianol, a new terpenoid hydroquinone from *Auxemma glazioviana*. *J Nat Prod (Lloydia)* **62**: 1044–1045.
- CRAVEN LA (1996): A taxonomic revision of *Heliotropium* (Boraginaceae) in Australia. *Austral Syst Bot* **9**: 521–657.
- CRONQUIST A (1988): The evolution and classification of flowering plants (2nd ed.). Lawrence, Kansas: Allen Press.
- DAHLGREN R (1983): General aspects of angiosperm evolution and macrosystematics. *Nord J Bot* **3**: 119–149.
- DAVIS CC, CD BELL, S MATHEWS & MJ DONOGHUE (2002): Laurasian migration explains Gondwanan disjunctions: evidence from Malpighiaceae. *Proc Natl Acad Sci USA* **99**: 6833–6837.
- DEHGAN B & CKKH YUEN (1983): Seed morphology in relation to dispersal, evolution, and propagation of *Cycas* L. *Bot Gaz* **144**: 412–418.
- DIANE N, H FÖRTHER & HH HILGER (2002a): A systematic analysis of *Heliotropium*, *Tournefortia*, and allied taxa of the Heliotropiaceae (Boraginales) based on ITS1 sequences and morphological data. *Amer J Bot* **89**: 287–295.
- DONG M, Y ODA & M HIROTA (2000): (10E,12Z,15Z)-9-Hydroxy-10,12,15-octadecatrienoic acid methyl ester as an anti-inflammatory compound from *Ehretia dicksonii*. *Biosci Biotechnol Biochem* **64**: 882–886.
- EGERTON-WARBURTON LM (1998): A smoke-induced alteration of the sub-testa cuticle in seeds of the post-fire recruiter, *Emmenanthe penduliflora* BENTH. (Hydrophyllaceae). *J Exp Bot* **49**: 1317–1327.
- FACCIOLA S (1990): Cornucopia. A source book of edible plants. Vista: Kampong Publications.
- FELSENSTEIN J (1981): Evolutionary trees from DNA sequences: A maximum likelihood approach. *J Mol Evol* **17**: 368–376.
- FELSENSTEIN J (1985): Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**: 783–791.
- FERGUSON DM (1999): Phylogenetic analysis and relationships in Hydrophyllaceae s.str. based on *ndhF* sequence data. *Syst Bot* **23**: 253–268.
- FÖRTHER H (1998): Die infragenerische Gliederung der Gattung *Heliotropium* L. und ihre Stellung innerhalb der subfam. Heliotropioideae (SCHRAD.) ARN. (Boraginaceae). *Sendtnera* **5**: 35–241.
- FREY W, M HOFMANN & HH HILGER (2001): The gametophyte-sporophyte junction: unequivocal hints for two evolutionary lines of archegoniate land plants. *Flora* **196**: 431–445.
- FRIESEN FCV (1933): Les caractères essentiels de la famille des Sebestenaceae et revision du genre *Varronia*. *Bull Soc Bot Genève* **24**: 117–201.
- FULVIO TE di (1965): Sobre la embriología de *Cortesia cuneifolia* (Boraginaceae). I. – Anatomia floral, esporogenesis y gametogenesis. *Kurtziana* **2**: 7–25.
- FULVIO TE di (1978): Sobre la vasculatura floral, embryología y chromosomas de *Ixorhea tschudiana* (Heliotropiaceae). *Kurtziana* **11**: 75–105.
- GAVIRIA JC (1987): Die Gattung *Cordia* in Venezuela. *Mitt Bot Staatssamml München* **23**: 1–279.

- GERSTBERGER P & P LEINS (1978): Rasterelektronenmikroskopische Untersuchungen an Blütenknospen von *Physalis philadelphica* (Solanaceae). Anwendung einer neuen Präparationsmethode. Ber Deutsch Bot Ges **91**: 381–387.
- GOLBOFF PA (1993): Nona Vers. 2.0. Tucumán.
- GOTTWALD H (1982): First description of the wood anatomy of *Antrophora*, *Lepidocordia* and *Pteleocarpa* (Boraginaceae). IAWA Bull ns **3**: 161–165.
- GRAHAM A & DM JARZEN (1969): Studies in neotropical palaeobotany. I. The Oligocene communities of Puerto Rico. Ann Missouri Bot Gard **56**: 308–357.
- GRAY A (1862): Notes upon a portion of Dr. SEEMAN's recent collection of dried plants gathered in the Feegee Islands. Proc Amer Acad Arts **5**: 340–341.
- GRAY A (1878): Forest geography and archaeology. A lecture delivered before the Harvard University Natural History Society. Amer J Sci Arts, Ser 3 **16**: 85–94, 183–196.
- GÜRKE M (1893): Boraginaceae. — In: ENGLER, A & K PRANTL (eds.): Die natürlichen Pflanzenfamilien **4** (3a). Leipzig: Engelmann.
- GUIGNARD NL (1893): Recherches sur le développement de la graine et en particulier du tégument séminal – Boraginées. J Bot **7**: 205–214.
- GUNNING BES & JS PATE (1969): ‘Transfer cells’ plant cells with wall ingrowth, specialized in relation to short distance transport of solutes – their occurrence, structure, and development. Protoplasma **68**: 107–133.
- GUSULEAC M (1937): Über die Orientierung des Ovulums bei den Boraginaceen und Labiaten, nebst Ausblick auf das System dieser Familien. — In: NISTOR II (ed.): Omagin lui Ion I. NISTOR. Cernauti: Glasul Bucovinei.
- HALLIER H (1923): Über die Lennoeen, eine zu LINNÉ's Bicornes verirrte Sippe der Boraginaceen. Beih Bot Centralbl (2. Abt) **40**: 1–19.
- HEGNAUER R (1989): Chemotaxonomie der Pflanzen. Nachträge zu Band 3 und 4 (Acanthaceae bis Lythraceae). Basel etc.: Birkhäuser Verlag.
- HEUBL GR, JC GAVIRIA & G WANNER (1990): A contribution to the taxonomy and evolution of *Cordia* (Boraginaceae) and allied genera. Chromosome numbers, pollen morphology and crystal pattern in wood. Bot Jahrb Syst **112**: 129–165.
- HILGER HH (1985): Ontogenie, Morphologie und Systematische Bedeutung geflügelter und glochidientragender Cynoglosseae- und Eritricheae-Früchte (Boraginaceae). Bot Jahrb Syst **105**: 323–378.
- HILGER HH (1987): Fruchtbiologische Untersuchungen an Heliotropioideae (Boraginaceae). I. Die Ontogenie der monospermen Früchte von *Heliotropium supinum* L. Flora **179**: 291–303.
- HILGER HH (1992): Morphology of *Heliotropium* (Boraginaceae) dispersal units. Bot Acta **105**: 387–393.
- HILGER HH, JR HOPPE & M HOFMANN (1993): Energiedispersiv Röntgenmikroanalyse (EDX) von Boraginaceae subfam. Boraginoideae – Klausenoberflächen (Sind Silicium- und Calcium-Einlagerungen in die Fruchtwand systematisch verwertbare Merkmale?). Flora **188**: 387–398.
- HOFMANN M (1994): Untersuchungen zur Gynoecialentwicklung in der Tribus Eritrichieae (Boraginaceae – Boraginoideae). Berlin (PhD Thesis): Freie Universität.
- HOFMANN M (1999): Flower and fruit development in the genus *Phacelia* (Phacelieae, Hydrophyllaceae): characters of systematic value. Syst Geogr Pl **68**: 203–212.

- HUTCHINSON J (1959): The families of flowering plants. Oxford: Clarendon Press.
- HUTCHINSON J (1969): Evolution and phylogeny of flowering plants. London etc.: Academic Press.
- IOSET J-R, A MARSTON, MP GUPTA & K HOSTETTMANN (2000): Antifungal and larvicidal compounds from the root bark of *Cordia alliodora*. *J Nat Prod (Lloydia)* **63**: 424–426.
- ITURRALDE-VINENT MA, RDE MACPHEE, S DÍAZ-FRANCO, R ROJAS-CONSUEGRA, W SUÁREZ & A LOMBA (2000): Las breas de San Felipe, a Quaternary fossiliferous asphalt seep near Mar't (Matanzas Province, Cuba). *Caribbean J Sci* **36**: 300–313.
- JAEGER JA, DH TURNER & M ZUCKER (1989): Improved predictions of secondary structures for RNA. *Proc Natl Acad Sci USA* **86**: 7706–7710.
- JAEGER JA, DH TURNER & M ZUCKER (1990) Predicting optimal and suboptimal secondary structure for RNA. *Meth Enzymol* **183**: 281–306.
- JOHNSTON IM (1930): Studies in the Boraginaceae, – VIII. *Contr Gray Herb* **92**: 3–95.
- JOHNSTON IM (1949): Studies in the Boraginaceae, XVII. A. *Cordia* section *Varronia* in Mexico and Central America. *J Arnold Arbor* **30**: 85–110.
- JOHNSTON IM (1950): Studies in the Boraginaceae XIX. A. Noteworthy species from tropical America. B. *Cordia* sect. *Gerascanthus* in Mexico and Central America. *J Arnold Arbor* **31**: 172–187.
- JOHNSTON IM (1951): Studies in the Boraginaceae, XX. Representatives of three subfamilies in Eastern Asia. *J Arnold Arbor* **32**: 1–26, 99–109.
- JOHRI BM & IK VASIL (1956): The embryology of *Ehretia laevis* ROXB. *Phytomorphology* **6**: 134–143.
- KANEHIRA R (1936): Formosan trees indigenous to the island. Tokyo: Department of Forestry, Government Research Institute.
- KHALEEL TF (1974): Embryology of *Cynoglossum denticulatum* DC. *Bot Not* **127**: 193–210.
- KHALEEL TF (1977): Embryology of *Ehretia acuminata*. *Proc Montana Acad Sci* **37**: 35–53.
- KHALEEL TF (1982): Embryology of *Cordia sebestena* (Boraginaceae). *Pl Syst Evol* **139**: 303–311.
- KHALEEL TF (1985): A review of endosperm and the taxonomical position of Boraginaceae. *J Pl Sci Res* **1**: 117–133.
- KIMURA M (1980): A simple method for estimating evolutionary rate of base substitutions though comparative studies of nucleotide sequences. *J Mol Evol* **16**: 111–120.
- KRAUSE K (1906): Boraginaceae andinae. *Bot Jahrb Syst* **37**: 627–636.
- KUNTZE O (1891): *Revisio generum plantarum* **2**. Leipzig etc.: Felix etc.
- KUHSEL MG, R STRICKLAND & J PALMER (1990): An ancient group I intron shared by Eubacteria and Chloroplasts. *Science* **250**: 1570–1573.
- LÄNGSTRÖM E & MW CHASE (2002): Tribes of Boraginoideae (Boraginaceae) and placement of *Antiphytum*, *Echiochilon*, *Ogastemma* and *Sericostoma*: A phylogenetic analysis based on *atpB* plastid DNA sequence data. *Pl Syst Evol* **234**: 137–153.
- LAWRENCE JR (1937): A correlation of the taxonomy and the floral anatomy of certain of the Boraginaceae. *Amer J Bot* **24**: 433–444.

- LEWIS WH (1991): Leaves of *Ehretia cymosa* (Boraginaceae) used to heal fractures in Ghana increase bone remodeling. *Econ Bot* **45**: 281–282.
- LIU J-S & CL SCHARDL (1994): A conserved sequence in Internal Transcribed Spacer 1 of plant nuclear rRNA genes. *Pl Mol Biol* **26**: 775–778.
- LLOYD DG (1979): Evolution towards dioecy in heterostylous populations. *Pl Syst Evol* **131**: 71–80.
- MAI DH (1991): Ehretiaceae. — In: MAI DH & H WALTHER (eds.): Die oligozänen und untermiozänen Floren NW-Sachsens und des Bitterfelder Raumes. Abh Staatl Mus Mineral Geol, Dresden **38**: 124, 220–221.
- MAI DH (1995): Tertiäre Vegetationsgeschichte Europas. Jena etc.: Fischer.
- MALCOMBER ST (2002): Phylogeny of *Gaertnera* LAM. (Rubiaceae) based on multiple DNA markers: evidence of a rapid radiation in a widespread, morphologically diverse genus. *Evolution* **56**: 42–57.
- MANOS P, JJ DOYLE & KC NIXON (1999): Phylogeny, biogeography, and processes of molecular differentiation in *Quercus* subgenus *Quercus* (Fagaceae). *Mol Phylogenetic Evol* **12**: 333–349.
- MAYOL M & JA ROSELLÓ (2001): Why nuclear ribosomal DNA spacers (ITS) tell different stories in *Quercus*. *Mol Phylogenetic Evol* **19**: 167–176.
- METCALFE CR & L CHALK (1950): Anatomy of the dicotyledons. Oxford: Clarendon Press.
- MEZ C (1890): Morphologische und anatomische Studien über die Gruppe der Cordieae. *Bot Jahrb Syst* **12**: 526–586.
- MIERS J (1875): On the Auxemmeæ, a new tribe of the Cordiaceæ. *Trans Linn Soc* **12**: 23–36, tab. 5–8.
- MILLER JS (1985): Systematics of the genus *Cordia* (Boraginaceae) in Mexico and Central America. London (PhD Thesis): Univ. Microfilms Internat.
- MILLER JS (1989): A revision of the New World species of *Ehretia* (Boraginaceae). *Ann Missouri Bot Gard* **76**: 1050–1076.
- MILLER JS (2001): Two new species of *Cordia* L. (Boraginaceae) from Madagascar. *Adansonia (Sér. 3)* **23**: 289–295.
- MILLER JS & JW NOWICKE (1989): Sectional placement of some problematic *Cordia* species (Boraginaceae). *Syst Bot* **14**: 271–280.
- MILLER JS & JW NOWICKE (1990): Dioecy and a reevaluation of *Lepidocordia* and *Antrophora* (Boraginaceae: Ehretioideae). *Amer J Bot* **77**: 543–551.
- MILLER RB (1977): Vestured pits in Boraginaceae. *IAWA Bull ns* **3**: 43–48.
- MULLER J (1981): Fossil pollen records of extant angiosperms. *Bot Rev* **47**: 1–142.
- NASIR YJ (1989): No. 191 Boraginaceae. *Fl Pakistan* **191**: 1–200.
- NAYAR TS, ES ANIL KUMAR & P PUSHPANGADAN (1999): *Rotula aquatica*, Boraginaceae – First record on its psychoactive property. *Econ Bot* **53**: 115–117.
- NOWICKE JW (1969): Flora of Panama Part IX. Family 167. Boraginaceae. *Ann Missouri Bot Gard* **56**: 33–69.
- NOWICKE JW & JS MILLER (1990): Pollen morphology of the Cordioideae (Boraginaceae): *Auxemma*, *Cordia*, and *Patagonula*. *Pl Syst Evol Suppl* **5**: 103–121.
- NOWICKE JW & JE RIDGWAY (1973): Pollen studies in the genus *Cordia* (Boraginaceae). *Amer J Bot* **60**: 584–591.

- NUES RW van, JMJ RIENTJES, CAFM van der SANDE, SF ZERP, C SLUITER, J VENEMA, RJ PLANTA & HA RAUÉ (1994): Separate structural elements within Internal Transcribed Spacer 1 of *Saccharomyces cerevisiae* precursor ribosomal RNA direct the formation of 17S and 26S rRNA. Nucleic Acids Res **22**: 912–919.
- OLMSTEAD RG, B BREMER, KM SCOTT & JD PALMER (1993): A parsimony analysis of the asteridae *sensu lato* based on *rbcL* sequences. Ann Missouri Bot Gard **80**: 700–722.
- OLMSTEAD RG & DM FERGUSON (2001). A molecular phylogeny of the Boraginaceae/Hydrophyllaceae s.str. (Abstract). <http://www.2001.botanyconference.org/section12/abstracts/167.shtml>.
- OPLER PA, HG BAKER & GH FRANKIE (1975): Reproductive biology of some Costa Rican *Cordia* species (Boraginaceae). Biotropica **7**: 234–247.
- PARMAR PJ (1991): 88. Ehretiaceae. — In: SHETTY BV & V SINGH (eds.): Flora of Rajasthan. Calcutta: Botanical Survey of India.
- PARRISH JT (1993): The palaeogeography of the opening South Atlantic. — In: GEORGE W & R LAVOCAT (eds.): The Africa-South America connection. Oxford: Clarendon Press.
- PATE JS & BES GUNNING (1972): Transfer Cells. Annual Rev Pl Physiol **23**: 173–196.
- PATE JS, BES GUNNING & FF MILLIKEN (1970): Function of transfer cells in the nodal regions of stems, particularly in relation to the nutrition of young seedlings. Protoplasma **71**: 313–334.
- PEER Y van de & R de WACHTER (1993): TREECON. A software package for the construction and drawing of evolutionary trees. Comput Appl Biosci **9**: 177–182.
- PERCIVAL M (1974): Floral ecology of coastal scrub in Southeastern Jamaica. Biotropica **6**: 104–129.
- PÉREZ-MOREAU RL & ÁL CABRERA (1983): Boraginaceae. — In: CABRERA ÁL (ed.): Flora de la provincia de Jujuy **8**. Buenos Aires: Colección Científica del INTA.
- PITMAN III WC, S CANDE, J LABREQUE & J PINDELL (1993): Fragmentation of Gondwana: the separation of Africa from South America. — In: GOLDBLATT P (ed.): Biological relationships between Africa and South America. New Haven, London: Yale University Press.
- PITOT A (1939a): Le fruit d' „*Ehretia serrata*“ ROXB. Bull Soc Bot France **86**: 310–314.
- PITOT A (1939b): Le fruit de quelques Ehrétiées. Bull Soc Bot France **86**: 325–332.
- POSADA D & KA CRANDALL (1998): Modeltest: testing the model of DNA substitution. Bioinformatics **14**: 817–818.
- QUISUMBING E (1951): Medical Plants of the Philippines. Manila: Bureau of Printing.
- RAMBAUT A (2001): Se-Al. Sequence alignment program v2.0a72. Oxford.
- RANDELL BR (1993): *Ehretia*. J Adelaide Bot Gard **15**: 93–94.
- RAO BH & SS RANI (1987): Floral anatomy of *Ehretia*. Acta Bot Indica **15**: 50–54.
- RAO BH & PSP RAO (1984): Embryology of three species of *Ehretia*. Proc Indian Acad Sci Pl Sci **93**: 57–65.
- RAVEN PH & DI AXELROD (1974): Angiosperm biogeography and past continental movements. Ann Missouri Bot Gard **61**: 539–673.
- REID EM (1923): Nouvelles recherches sur les graines du Pliocène inférieur du Pont-de-Gail (Cantal). Bull Soc Geol France, 4. Ser **23**: 308–355.
- REID EM & MEJ CHANDLER (1933): Flora of the London Clay. London: British Museum.

- RENNER SS, DB FOREMAN & D MURRAY (2000): Timing transantarctic disjunctions in the Atherospermataceae (Laurales): evidence from coding and noncoding chloroplast sequences. *Syst Biol* **49**: 579–591.
- RENNER SS & K MEYER (2001): Melastome come full circle: biogeographic reconstruction and molecular clock dating. *Evolution* **55**: 1315–1324.
- RICHARDSON AT (1977): Monograph of the genus *Tiquilia* (*Coldenia*, sensu lato), Boraginaceae: Ehretioideae. *Rhodora* **79**: 467–572.
- RIDLEY NH (1930): The dispersal of plants throughout the world. Ashford, Kent: Reeve.
- RIEDL H (1997): Boraginaceae. Fl Malesiana, Ser 1, Spermatophyt **13**: 43–144.
- RITLAND CE, K RITLAND & NA STRAUS (1993): Variation in the ribosomal internal transcribed spacers (ITS1 and ITS2) among eight taxa of the *Mimulus guttatus* species complex. *Mol Biol Evol* **10**: 1273–1288.
- ROHWER JG (1996): Die Frucht- und Samenstrukturen der Oleaceae. Eine vergleichend-anatomische Untersuchung. *Biblioth Bot* **148**: 1–177.
- SAHAY SK (1979): On the pollen morphology of Ehretiaceae with reference to taxonomy. IV Int Palynol Conf Lucknow 1976-77 **1**: 471–479.
- SARGENT CS, CE FAXON & MW GILL (1933): Manual of the trees of North America. Boston, New York: Houghton Mifflin Company.
- SARGENT CS, CE FAXON, P PICART & E PICART (1947): The silva of North America. New York: Smith.
- SANDERSON MJ (1998) Estimating rate and time in molecular phylogenies: beyond the molecular clock? — In: SOLTIS DE, PS SOLTIS & JJ DOYLE (eds.): Molecular systematics of plants II: DNA sequencing. Norwell: Kluwer Academic Publishers.
- SCHULENBURG JHG von der, U ENGLISCH & J-W WÄGELE (1999): Evolution of ITS1 rDNA in the Digenea (Platyhelminthes: Trematoda): 3' end sequence conservation and its phylogenetic utility. *J Mol Evol* **48**: 2–12.
- SCHULZ OE (1911): III. *Beureria* JACQ. – In: URBAN I (ed.): *Symbolae Antillanae* **7** (1). Leipzig etc.: Bornträger etc.
- SCOTESE CR (1998): Quicktime computer animations, PALEOMAP Project. Arlington: University of Texas.
- SCOTESE CR, LM GAHAGAN & RL LARSON (1988): Plate tectonic reconstructions of the Cretaceous and Cenozoic ocean basins. *Tectonophysics* **155**: 27–48.
- SCOTT PE & RF MARTIN (1984): Avian consumers of *Bursea*, *Ficus*, and *Ehretia* fruit in Yucatán. *Biotropica* **16**: 319–323.
- SEIBERT J (1978): Fruchtanatomische Untersuchungen an Lithospermeae (Boraginaceae). *Diss Bot* **44**: 1–207.
- SELVANAYAGAM ZE, SG GNAVAVENDHAND, K BALAKRISHNA, R RAO & SU BHIMA ALI (1995): Survey of medicinal plants with antisnake venom activity in Chengalpattu District, Tamilnadu, India. *Fitoterapia* **66**: 488–494.
- SERRATO-VALENTI G, L CORNARA, P MODENESI, M PIANA & MG MARIOTTI (2000): Structure and histochemistry of embryo envelope tissues in the mature dry seed and early germination in *Phacelia tanacetifolia*. *Ann Bot (Oxford)* **85**: 625–634.
- SLOTTA TAB (2000): Phylogenetic analysis of *Iliamna* (Malvaceae) using the Internal Transcribed Spacer region. Blacksburg (Thesis): Virginia Polytechnic Institute and State University.

- SMITH RA & CW DEPAMPHILIS (1998): Phylogenetic placement of the holoparasitic family Lennoaceae: preliminary molecular evidence (Abstract). Amer J Bot Suppl **85**: 157.
- SMITH RA, DM FERGUSON, TJ BARKMAN & CW DEPAMPHILIS (2000): Molecular phylogenetic evidence for the origin of Lennoaceae: a case of adelphoparasitism in the angiosperms (Abstract). Amer J Bot Suppl **87**: 158.
- SNOW DW (1981): Tropical frugivorous birds and their food plants: a world survey. Biotropica **13**: 1–14.
- SOLTIS PS, DE SOLTIS, V SAVOLAINEN, PR CRANE & TG BARRACLOUGH (2002): Rate heterogeneity among lineages of tracheophytes: integration of molecular and fossil data and evidence for molecular living fossils. Proc Natl Acad Sci USA **99**: 4430–4435.
- SPURR AR (1969): A low-viscosity epoxy resin embedding medium for electron microscopy. J Ultrastruct Res **26**: 31–43.
- STEGELMEIER BL, JA EDGAR, SM COLEGATE, DR GARDNER, TK SCHOCH, RA COULOMBE & RJ MOLYNEUX (1999): Pyrrolizidine alkaloid plants, metabolism and toxicity. J Nat Toxins **8**: 95–116.
- STRUWE L, M THIV, JW KADEREIT, AS-R PEPPER, TJ MOTLEY, PJ WHITE, JHE ROVA, K POTGIETER & VA ALBERT (1998): *Saccifolium* (Saccifoliaceae), an endemic of Sierra de la Neblina on the Brazilian-Venezuelan border, is related to a temperate-alpine lineage of Gentianaceae. Harvard Pap Bot **3**: 199–214.
- SUH Y, LB THIEN, HE REEVE & EA ZIMMER (1993): Molecular evolution and phylogenetic implications of Internal Transcribed Spacer sequences of ribosomal DNA in Winteraceae. Amer J Bot **80**: 1042–1055.
- SUN G, DL DILCHER, S ZHENG & Z ZHOU (1998): In search of the first flower: a Jurassic angiosperm, *Archaefructus*, from Northeast China. Science **282**: 1692–1695.
- SVENSSON HG (1925): Zur Embryologie der Hydrophyllaceen, Boraginaceen und Heliotropiaceen mit besonderer Rücksicht auf die Endospermmbildung. Uppsala (PhD Thesis): Universitets Årsskrift.
- SWOFFORD DL (1998): PAUP*. Phylogenetic Analysis Using Parsimony *and other methods. Version 4. Sunderland: Sinauer Associates.
- TABERLET P, L GIELLY, G PATOU & J BOUVET (1991): University primers for amplification of three non-coding regions of chloroplast DNA. Pl Mol Biol **17**: 1105–1109.
- TAKHTAJAN A (1986): Floristic regions of the world. Berkley: University of California Press.
- TARODA N & P GIBBS (1986): Studies on the genus *Cordia* L. (Boraginaceae) in Brazil. 1. A new infrageneric classification and conspectus. Revista Brasil Bot **9**: 31–42.
- THOMPSON RD, G HUEROS, HA BECKER & M MAITZ (2001): Development and functions of seed transfer cells. Pl Sci (Elsevier) **160**: 775–783.
- THORNE RF (1972): Major disjunctions in the geographic ranges of seed plants. Quart Rev Biol **47**: 365–411.
- THORNE RF (1983): Proposed new realignments in the angiosperms. Nord J Bot **3**: 85–117.
- TIFFNEY BH (1985a): Perspectives on the origin of the floristic similarity between Eastern Asia and Eastern North America. J Arnold Arbor **66**: 73–94.
- TIFFNEY BH (1985b): The Eocene North Atlantic land bridge: its importance in Tertiary and modern phytogeography of the Northern Hemisphere. J Arnold Arbor **66**: 243–273.
- TOELKEN HR & LA CRAVEN (1986): Family – Boraginaceae. — In: JESSOP JP & HP TOELKEN (eds.): Flora of South Australia **3**. Adelaide: Woolman.

- TOMLINSON PB & P FAWCETT (1980): The biology of trees native to Tropical Florida. Cambridge: Harvard University Press.
- UHLARZ H & F WEBERLING (1977): Ontogenetische Untersuchungen an *Cordia verbenaceae* DC. (Boraginaceae), ein Beitrag zur Kenntnis der Syndesmien. Ber Deutsch Bot Ges **90**: 127–134.
- USHER G (1972): A dictionary of plants used by man. London: Constable.
- VELDKAMP JF (1988): Notes on *Pteleocarpa, incertae sedis*. Fl Males Bull **10**: 47–50.
- VERDCOURT B (1991): Boraginaceae. — In: POLHILL RM (ed.): Flora of tropical East Africa. Rotterdam, Brookfield: Balkema.
- WÄGELE J-W (2000): Grundlagen der Phylogenetischen Systematik. München: Pfeil.
- WARFA AM (1988): *Cordia* (Boraginaceae) in NE Tropical Africa and Tropical Arabia. Acta Univ Uppsala **174**: 1–80.
- WATT JM & MG BREYER-BRANDWIJK (1962): The medical and poisonous plants of Southern and Eastern Africa. Edinburgh, London: Livingstone.
- WENDEL JF, A SCHNABEL & T SEELANAN (1995): An unusual ribosomal sequence from *Gossypium gossypoides* reveals ancient, cryptic, intergenicomic introgression. Mol Phylogenetic Evol **4**: 298–313.
- WIKSTRÖM N, V SAVOLAINEN V & MW CHASE (2001): Evolution of the angiosperms: calibrating the family tree. Proc Roy Soc London, Ser B, Biol Sci **268**: 2211–2220.
- WOLF M, M BUCHHEIM, E HEGEWALD, L KRIENITZ & D HEPPERLE (2002): Phylogenetic Position of the Sphaeropleaceae (Chlorophyta). Pl Syst Evol **230**: 161–171.
- WOLFE, JA (1981): Vicariance biogeography of angiosperms in relation to paleobotanical data. – In: NELSON G, EE ROSEN (eds.): Vicariance biogeography, a critique. New York: Columbia University Press.
- XIANG Q-Y, DE SOLTIS, PS SOLTIS, SR MANCHESTER & DJ CRAWFORD (2000): Timing the Eastern Asian-Eastern North American floristic disjunction: molecular clock corroborates paleontological estimates. Mol Phylogenetic Evol **15**: 462–472.
- XU MQ, SD KATHE, H GOODRICH-BLAIR, SA NIERZWICKI-BAUER & DA SHUB (1990): Bacterial origin of a chloroplast intron: conserved self-splicing group I introns in Cyanobacteria. Science **250**: 1566–1570.
- YATSKIEVYCH G, J MASON Jr. & T CHARLES (1986): A revision of the Lennoaceae. Syst Bot **11**: 531–548.
- YATSKIEVYCH G & M ZAVADA (1984). Pollen Morphology of Lennoaceae. Pollen & Spores **26**: 19–30.