

8. Literaturverzeichnis

- (1) Abel SM, Back DJ, Maggs JL, Park BK. Cortisol Metabolism by human liver in *in vitro* - IV. Metabolism of 9 α -Fluorocortisol by human liver Microsomes and Cytosol. J.Steroid Biochem.Mol.Biol. 46[6], 833-839. 1993.
- (2) Ackermann D, Vogt B, Escher G, Dick B, Reichen J, Frey BM et al. Inhibition of 11 β -hydroxysteroid dehydrogenase by bile acids in rats with cirrhosis. Hepatology 1999; 30:623-629.
- (3) Addison RS, Maguire DJ, Mortimer RH, Roberts MS, Cannell GR. Pathway and kinetics of prednisolone metabolism in the human placenta. J Steroid Biochem Mol Biol 1993; 44:315-320.
- (4) Agarwal AK, Mune T, Monder C, White PC. Mutations in putative glycosylation sites of rat 11 beta-hydroxysteroid dehydrogenase affect enzymatic activity. Biochim Biophys Acta 1995; 1248(1):70-74.
- (5) Agarwal AK, Rogerson FM, Mune T, White PC. Analysis of the human gene encoding the kidney isozyme of 11 beta- hydroxysteroid dehydrogenase. J Steroid Biochem Mol Biol 1995; 55(5-6):473-479.
- (6) Agarwal AK, Rogerson FM, Mune T, White PC. Gene structure and chromosomal localization of the human HSD11K gene encoding the kidney (type 2) isozyme of 11 beta-hydroxysteroid dehydrogenase. Genomics 1995; 29(1):195-199.
- (7) Albiston AL, Obeyesekere VR, Smith RE, Krozowski ZS. Cloning and tissue distribution of the human 11 beta-hydroxysteroid dehydrogenase type 2 enzyme. Mol Cell Endocrinol 1994; 105(2):R11-R17.
- (8) Albiston AL, Smith RE, Obeyesekere VR, Krozowski ZS. Cloning of the 11 beta HSD type II enzyme from human kidney. Endocr Res 1995; 21(1-2):399-409.
- (9) Ananthanarayanan M, Balasubramanian N, Makishima M, Mangelsdorf DJ, Suchy FJ. Human bile salt export pump promoter is transactivated by the farnesiod X receptor/bile acid receptor. J Biol Chem 2001; 276(31):28857-28865.
- (10) Andrews R, Walker BR. Glucocorticoids and Insulin resistance: old hormones, new targets. Clin Sci (Lond) 1999; 96:513-523.
- (11) Arcuri F, Monder C, Lockwood CJ, Schatz F. Expression of 11 beta-hydroxysteroid dehydrogenase during decidualization of human endometrial stromal cells. Endocrinology 1996; 137(2):595-600.
- (12) Armanini D, Karbowiak I, Funder JW. Affinity of liquorice derivates for mineralocorticoid and glucocorticoid receptors. Clin Endocrinol (Oxf) 1983; 19:609-612.
- (13) Arriza JL, Weinberger C, Cerelli G, Glaser TM, Handelin BL, Housman DE. Cloning of human mineralocorticoid receptor complementary DNA : structural and functional kinship with the glucocorticoid receptor. Science 1987; 237:268-275.

- (14) Assandri A, Buniva G, Martinelli E, Perazzi A, Zerilli L. Pharmacokinetics and Metabolism of Deflazacort in the rat, dog, monkey and man. *Adv Exp Med Biol* 1984; 17:9-23.
- (15) Baenkler HW, et.al. Innere Medizin., Kap. Laborwerte und Referenzbereiche Hippokrates Verlag Stuttgart, 1999.
- (16) Bahr V, Pfeiffer AF, Diederich S. The metabolic syndrome X and peripheral cortisol synthesis. *Exp Clin Endocrinol Diabetes* 2002; 110(7):313-318.
- (17) Ballard PL, Carter JP, Graham BS, Baxter, JD. A radioreceptor assay for evaluation of the plasma glucocorticoid activity of natural and synthetic steroids in man. *J Clin Endocrinol Metab* 1975; 41:290-304.
- (18) Bamberger CM, Schulte HM, Chrousos GP. Molecular determinants of glucocorticoid receptor function and tissue sensitivity to glucocorticoids. *Endocr Rev* 1996; 17(3):245-261.
- (19) Barany E. In vitro uptake of bile acids by choroid plexus, kidney cortex and anteroir uvea. The iodipamide-sensitive transport systems in the rabbit. *Acta Physiol Scand* 1975; 93:250-268.
- (20) Barf T, Vallgarda J, Emond R, Haggstrom C, Kurz G, Nygren A et al. Arylsulfonamidothiazoles as a new class of potential antidiabetic drugs. Discovery of potent and selective inhibitors of the 11beta-hydroxysteroid dehydrogenase type 1. *J Med Chem* 2002; 45(18):3813-3815.
- (21) Benediktsson R, Calder AA, Edwards CR, Seckl JR. Placental 11 beta-hydroxysteroid dehydrogenase: a key regulator of fetal glucocorticoid exposure. *Clin Endocrinol (Oxf)* 1997; 46(2):161-166.
- (22) Benediktsson R, Magnusdottir EM, Smith JC, Seckl JR. The effects of ethanol on 11 β -hydroxysteroid dehydrogenase type-2 activity. *J Endocrinol* [15th Joint Meeting], P51. 1996.
- (23) Beretta-Piccoli C, Davis DL, Boddy K, Brown JJ, Cumming AM, East BW. The relation of arterial pressure with plasma and body electrolytes is similar in Conn's syndrome and essential hypertension. *Clin Sci (Lond)* 1982; 63:89-92.
- (24) Berger J, Tanen M, Elbrecht A, Hermanowski-Vosatka A, Moller DE, Wright SD et al. Peroxisome proliferator-activated-receptor gamma ligands inhibit adipocyte 11beta-hydroxysteroid dehydrogenase expression and activity. *J Biol Chem* 2001; 276:12629-12635.
- (25) Best R, Nelson SM, Walker BR. Dexamethasone and 11-dehydrodexamethasone as tools to investigate the isozymes of 11 beta-hydroxysteroid dehydrogenase in vitro and in vivo. *J Endocrinol* 1997; 133(1):41-48.
- (26) Best R, Walker BR. Additional value of measurement of urinary cortisone and unconjugated cortisol metabolites in assessing the activity of 11 beta-hydroxysteroid dehydrogenase in vivo. *Clin Endocrinol (Oxf)* 1997; 47(2):231-236.

- (27) Blum A, Martin HJ, Maser E. Human 11beta-hydroxysteroid dehydrogenase type 1 is enzymatically active in its nonglycosylated form. *Biochem Biophys Res Commun* 2000; 276(2):428-434.
- (28) Bowers A, Halsall T, Jones E, Lemin A. *J Chem Soc* 1953;2548.
- (29) Brand E, Kato N, Chatelain N, Krozowski ZS, Jeunemaitre X, Corvol P et al. Structural analysis and evaluation of the 11beta-hydroxysteroid dehydrogenase type 2 (11beta-HSD2) gene in human essential hypertension. *J Hypertens* 1998; 16(11):1627-1633.
- (30) Brown RW, Chapman KE, Kotelevtsev Y, Yau JL, Lindsay RS, Brett L et al. Cloning and production of antisera to human placental 11 beta-hydroxysteroid dehydrogenase type 2. *Biochem J* 1996; 313 (Pt 3):1007-1017.
- (31) Brown RW, Chapman KE, Murad P, Edwards CR, Seckl JR. Purification of 11 beta-hydroxysteroid dehydrogenase type 2 from human placenta utilizing a novel affinity labelling technique. *Biochem J* 1996; 313 (Pt 3):997-1005.
- (32) Broyer M, Terzi F, Lehnert A, Gagnadoux MF, Guest G, Niaudet P. A controlled study of deflazacort in the treatment of idiopathic nephrotic syndrome. *Pediatr Nephrol* 1997; 11(4):418-422.
- (33) Buhler H, Wang D, Hundertmark S, Blum S. Induction of the glucocorticoid metabolizing enzyme 11 β -HSD in MDCK cells by preincubation with Dexamethasone, Agonistic effects of the "pure" Antiglucocorticoid RU 486. *Wissenschaftswoche Universitätsklinikum Benjamin Franklin* 1993;50-51.
- (34) Bujalska I, Shimojo M, Howie A, Stewart PM. Human 11 beta-hydroxysteroid dehydrogenase: studies on the stably transfected isoforms and localization of the type 2 isozyme within renal tissue. *Steroids* 1997; 62(1):77-82.
- (35) Bujalska IJ, Kumar S, Hewison M, Stewart PM. Differentiation of adipose stromal cells: the roles of glucocorticoids and 11beta-hydroxysteroid dehydrogenase. *Endocrinology* 1999; 140(7):3188-3196.
- (36) Bujalska IJ, Kumar S, Stewart PM. Does central obesity reflect "Cushing's disease of the omentum"? *Lancet* 1997; 349(9060):1210-1213.
- (37) Burkhardt P, Hanke B, Baehr V, Oelkers W, Diederich S. Human 11 β -hydroxysteroid-dehydrogenases and their metabolism of synthetic glucocorticoids. *Jahrbuch 1998 - Fachbereich Humanmedizin, UKBF der Freien Universität Berlin*, 382. 1998.
- (38) Burton PJ, Waddell BJ. 11 beta-Hydroxysteroid dehydrogenase in the rat placenta: developmental changes and the effects of altered glucocorticoid exposure. *J Endocrinol* 1994; 143(3):505-513.
- (39) Bush IE, Hunter S. The Reduction of 9 α -Chloro-11-Oxo Steroids by Preparations of rat liver. *Biochem J* 2001; 71:16p-16p.
- (40) Bush IE, Hunter S, Meigs RA. Metabolism of 11-Oxygenated Steroids Metabolism *in vitro* by preparations of liver. *Biochem J* 107, 239-258. 1968.

- (41) Bush IE, Mahesh VB. Metabolism of 11-oxygenated Steroids - 2-Methyl-Steroids. *Biochem J* 71, 718-742. 1959.
- (42) Bush IE, Mahesh VB. Metabolism of 11-oxygenated steroids. 3. Some 1-dehydro and 9 alpha- fluoro steroids. *Biochem J* 1964; 93(2):236-255.
- (43) Chanoine F, Grenot C, Heidmann P, Junien J. Pharmacokinetics of Butixocort 21-Dipropionate, Budesonide and Beclomethasone Dipropionate in the rat after intratracheal, intravenous and oral treatments. *Drug Metab Dispos* 1991; 19(2):546-553.
- (44) Coghl JP, Butkus A, Denton DA, McDougall JG, Scoggins BA, Whitworth JA. Blood pressure and metabolic effects of 9 α -Fluorocortisone in sheep. *Clin Exp Hypertens* 1, 629-648. 1979.
- (45) Darnel AD, Archer TK, Yang K. Regulation of 11beta-hydroxysteroid dehydrogenase type 2 by steroid hormones and epidermal growth factor in the Ishikawa human endometrial cell line. *J Steroid Biochem Mol Biol* 1999; 70(4-6):203-210.
- (46) Davis EA, Morris DJ. Medicinal uses of licorice through the millenia: The good and plenty of it. *Mol Cell Endocrinol* 1991; 78:1-6.
- (47) Diederich S, Eigendorff E, Burkhardt P, Quinkler M, Bumke-Vogt C, Rochel M et al. 11beta-Hydroxysteroid Dehydrogenase Types 1 and 2: An Important Pharmacokinetic Determinant for the Activity of Synthetic Mineralo- and Glucocorticoids. *J Clin Endocrinol Metab* 2002; 87(12):5695-5701.
- (48) Diederich S, Grossmann C, Hanke B, Quinkler M, Herrmann M, Bahr V et al. In the search for specific inhibitors of human 11beta-hydroxysteroid- dehydrogenases (11beta-HSDs): chenodeoxycholic acid selectively inhibits 11beta-HSD-I. *Eur J Endocrinol* 2000; 142(2):200-207.
- (49) Diederich S, Hanke B, Bahr V, Oelkers W. The metabolism of 9 alpha-fluorinated steroids in the human kidney. *Endocr Res* 1996; 22(4):803-810.
- (50) Diederich S, Hanke B, Burkhardt P, Muller M, Schoneshofer M, Bahr V et al. Metabolism of synthetic corticosteroids by 11 beta-hydroxysteroid-dehydrogenases in man. *Steroids* 1998; 63(5-6):271-277.
- (51) Diederich S, Hanke B, Oelkers W, Bahr V. Metabolism of dexamethasone in the human kidney: nicotinamide adenine dinucleotide-dependent 11beta-reduction. *J Clin Endocrinol Metab* 1997; 82(5):1598-1602.
- (52) Diederich S, Quinkler M, Burkhardt P, Grossmann C, Bahr V, Oelkers W. 11Beta-hydroxysteroid-dehydrogenase isoforms: tissue distribution and implications for clinical medicine. *Eur J Clin Invest* 2000; 30 Suppl 3:21-27.
- (53) Dotsch J, Dorr HG, Stalla GK, Sippell WG. Effect of glucocorticoid excess on the cortisol/cortisone ratio. *Steroids* 2001; 66(11):817-820.

- (54) Dötsch J, Rascher W. The role of 11 β -hydroxysteroid dehydrogenase activity in the metabolic syndrome: lessons learned from the animal model. *Eur J Endocrinol* 2002; 146:603-605.
- (55) Draper N, Walker EA, Bujalska IJ, Tomlinson JW, Chalder SM, Arlt W et al. Mutations in the genes encoding 11 β -hydroxysteroid dehydrogenase type 1 and hexose-6-phosphate dehydrogenase interact to cause cortisone reductase deficiency. *Nat Genet* 2003; 34(4):434-439.
- (56) Edsbacker S, Andersson KE, Ryrfeldt A. Nasal bioavailability and systemic effects of the glucocorticoid budesonide in man. *Eur J Clin Pharmacol* 1985; 29:477-481.
- (57) Edwards CR, Walker BR, Benediktsson R, Seckl JR. Congenital and acquired syndromes of apparent mineralocorticoid excess. *J Steroid Biochem Mol Biol* 1993; 45(1-3):1-5.
- (58) Escher G, et.al. Down-regulation of hepatic and renal 11 β -hydroxysteroid dehydrogenase in rats with liver cirrhosis. *Gastroenterology* 1998; 114:175-184.
- (59) Escher G, Frey FJ, Frey BM. 11beta-Hydroxysteroid dehydrogenase accounts for low prednisolone/prednisone ratios in the kidney. *Endocrinology* 1994; 135(1):101-106.
- (60) Feinstein M, Schleimer R. Regulation of the action of hydrocortisone in airway epithelial cells by 11beta-hydroxysteroid dehydrogenase. *Am J Respir Cell Mol Biol* 1999; 21(3):403-408.
- (61) Felgner PL, Gadek TR, Holm M, Roman R, Chan HW, Wenz M et al. Lipofectin: A highly efficient, lipid mediated DNA - transfection procedure. *Proc Natl Acad Sci U S A* 1987; 34:7413-7417.
- (62) Ferrari P, Lovati E, Frey FJ. The role of the 11beta-hydroxysteroid dehydrogenase type 2 in human hypertension. *J Hypertens* 2000; 18(3):241-248.
- (63) Ferrari P, Smith RE, Funder JW, Krozowski ZS. Substrate and inhibitor specificity of the cloned human 11 beta-hydroxysteroid dehydrogenase type 2 isoform. *Am J Physiol* 1996; 270(5 Pt 1):E900-E904.
- (64) Ferrari P, Smith RE, Funder JW, Krozowski ZS. Substrate and inhibitor specificity of the cloned human 11 beta-hydroxysteroid dehydrogenase type 2 isoform. *Am J Physiol* 1996; 270(5 Pt 1):E900-E904.
- (65) Frey FJ, Escher G, Frey BM. Pharmacology of 11 β -hydroxysteroid dehydrogenase. *Steroids* 1994; 59:74-79.
- (66) Fried J, Sabo E. J.Amer.chem.Soc. 76, 1455. 1953.
- (67) Fukushima K, Bradlow HL, Hellmann L, Gallagher TF. Metabolic transformation of hydrocortisone-4-C14 in normal men. *J Biol Chem* 1960; 235:2246-2253.
- (68) Funder JW, Pearce PT, Smith R, Smit AJ. Mineralocorticoid action: Target tissue specificity is enzyme, not receptor, mediated. *Science* 1988; 242:583-585.

- (69) Gaignage P, Lognay G, Marlier M, Severin M, Dreze P. Applications of chromatographic and spectrometric techniques to the study of dexamethasone related compounds. *Chromatographia* 1989; 28:623-630.
- (70) Ge RS, Gao HB, Nacharaju VL, Gunsalus, Hardy MP. Identification of a kinetically distinct activity of 11 β -hydroxysteroid dehydrogenase in rat Leydig cells. *Endocrinology* 1997; 138:2435-2442.
- (71) Gerhards E, Nieuweboer G, Schulz G, Gibian H. Stoffwechsel von Fluocortolon beim Menschen. *Acta Endocrinologica* 1971; 68:98-126.
- (72) Ghazarian JG, Demura H. 25-Hydroxycholecalciferol-1-hydroxylase: a specific requirement for NADPH and a hemoprotein component in chick kidney mitochondria. *Arch Biochem Biophys* 1974; 160:63-72.
- (73) Gomez-Sanchez EP, Ganjam V, Chen YJ, Cox DL, Zhou MY, Thanigaraj S et al. The sheep kidney contains a novel unidirectional, high affinity NADP(+) -dependent 11 beta-hydroxysteroid dehydrogenase (11 beta-HSD-3). *Steroids* 1997; 62(5):444-450.
- (74) Gomez-Sanchez EP, Ganjam V, Chen YJ, Liu Y, Clark SA, Gomez-Sanchez CE. The 11beta hydroxysteroid dehydrogenase 2 exists as an inactive dimer. *Steroids* 2001; 66(11):845-848.
- (75) Grundy WN, Bailey, TL, Elkan CP, Baker ME. Hidden Markov model analysis of motifs in steroid dehydrogenases and their homologs. *Biochem Biophys Res Commun* 1997; 231:760-766.
- (76) Hague N, Thrasher K, Werk E, Knowles H, Sholiton L. Studies on Dexamethasone metabolism in man, effect of diphenylhydantoin. *J Clin Endocrinol Metab* 1972; 34:44-50.
- (77) Hammami MM, Siiteri PK. Regulation of 11 beta-hydroxysteroid dehydrogenase activity in human skin fibroblasts: enzymatic modulation of glucocorticoid action. *J Clin Endocrinol Metab* 1991; 73(2):326-334.
- (78) Handoko K, Yang K, Strutt BJ, Khalil MW, Killinger DW. Insulin attenuates the stimulatory effects of tumor necrosis factor α on 11 β -hydroxysteroid dehydrogenase 1 in human adipose stromal cells. *J Steroid Biochem Mol Biol* 2000; 72:163-168.
- (79) Harris HJ, Kotelevtsev Y, Mullins JJ, Seckl JR, Holmes MC. Intracellular regeneration of glucocorticoids by 11beta-hydroxysteroid dehydrogenase (11beta-HSD)-1 plays a key role in regulation of the hypothalamic-pituitary-adrenal axis: analysis of 11beta-HSD-1-deficient mice. *Endocrinology* 2001; 142(1):114-120.
- (80) Heilmann P, Heide J, Hundertmark S, Schoneshofer M. Administration of glycyrrhetic acid: significant correlation between serum levels and the cortisol/cortisone-ratio in serum and urine. *Exp Clin Endocrinol Diabetes* 1999; 107(6):370-378.
- (81) Hirasawa G, Sasano H, Takahashi K, Fukushima K, Suzuki K, Hiwatashi N et al. Colocalisation of 11 β -hydroxysteroid dehydrogenase type II and mineralocorticoid receptor in human epithelia. *J Clin Endocrinol Metab* 1997; 82:3859-3863.

- (82) Holmes MC, Kotelevtsev Y, Mullins JJ, Seckl JR. Phenotypic analysis of mice bearing targeted deletions of 11beta-hydroxysteroid dehydrogenases 1 and 2 genes. *Mol Cell Endocrinol* 2001; 171(1-2):15-20.
- (83) Hult M, Jörnvall H, Oppermann UC. Selective inhibition of human type1 11 β -hydroxysteroid dehydrogenase by synthetic steroids and xenobiotics. *FEBS Letters* 441, 25-28. 1998.
- (84) Hundertmark S, Buhler H, Ragosch V, Dinkelborg L, Arabin B, Weitzel HK. Correlation of surfactant phosphatidylcholine synthesis and 11 beta-hydroxysteroid dehydrogenase in the fetal lung. *Endocrinology* 1995; 136(6):2573-2578.
- (85) Hundertmark S, Buhler H, Rudolf M, Weitzel HK, Ragosch V. Inhibition of 11 beta-hydroxysteroid dehydrogenase activity enhances the antiproliferative effect of glucocorticosteroids on MCF-7 and ZR-75-1 breast cancer cells. *J Endocrinol* 1997; 155(1):171-180.
- (86) Iki K, Miyamori I, Hatakeyama H, Yoneda T, Takeda Y, Takeda R et al. The activities of 5 beta-reductase and 11 beta-hydroxysteroid dehydrogenase in essential hypertension. *Steroids* 1994; 59(11):656-660.
- (87) Imbasciati E, Gusmano R, Edefonti A, Zucchelli P, Pozzi C, Grassi C et al. Controlled trial of methylprednisolone pulses and low dose oral prednisone for the minimal change nephrotic syndrome. *Br Med J (Clin Res Ed)* 1985; 291(6505):1305-1308.
- (88) Jamieson PM, Chapman KE, Edwards CR, Seckl JR. 11 beta-hydroxysteroid dehydrogenase is an exclusive 11 β -reductase in primary cultures of rat hepatocytes: Effect of physicochemical and hormonal manipulations. *Endocrinology* 1995; 136(11):4754-4761.
- (89) Jamieson PM, Walker BR, Chapman KE, Andrew R, Rossiter S, Seckl JR. 11 beta-hydroxysteroid dehydrogenase type 1 is a predominant 11 beta-reductase in the intact perfused rat liver. *J Endocrinol* 2000; 165(3):685-692.
- (90) Jellinck PH, Dhabhar FS, Sakai RR, McEwen BS. Long-term corticosteroid treatment but not chronic stress affects 11beta-hydroxysteroid dehydrogenase type I activity in rat brain and peripheral tissues. *J Steroid Biochem Mol Biol* 1997; 60(5-6):319-323.
- (91) Jenkins J, Sampson P. Conversion of Cortisol and Prednisolone to Prednison. *Br Med J* 1967; 2:205-207.
- (92) Kaiser H, Kley H. Cortisontherapie - Corticoide in Klinik und Praxis. 1992.
- (93) Katz JR, Mohamed-Ali V, Wood PJ, Yudkin JS, Coppock SW. An in vivo study of the cortisol-cortisone shuttle in subcutaneous abdominal adipose tissue. *Clin Endocrinol (Oxf)* 1999; 50(1):63-68.
- (94) Kenouch S, Lombes M, Delahaye F, Eugene E, Bonvalet JP, Farman N. Human skin as target for aldosterone: Coexpression of mineralocorticoid receptors and 11 β -hydroxysteroid dehydrogenase. *J Clin Endocrinol Metab* 1994; 79:1334-1341.

- (95) Korbmacher C, Schulz W, Konig M, Siebe H, Lichtenstein I, Hierholzer K. Renal epithelial cell lines (BSC-1, MDCK, LLC-PK1) express 11 beta-hydroxysteroid dehydrogenase activity. *Biochim Biophys Acta* 1989; 1010(3):311-317.
- (96) Kotelevtsev Y, Holmes MC, Burchell A, Houston PM, Schmoll D, Jamieson P et al. 11beta-hydroxysteroid dehydrogenase type 1 knockout mice show attenuated glucocorticoid-inducible responses and resist hyperglycemia on obesity or stress. *Proc Natl Acad Sci U S A* 1997; 94(26):14924-14929.
- (97) Krozowski Z. 11 β -Hydroxysteroid dehydrogenase and the short-chain alcohol dehydrogenase (SCAD)superfamily. *Mol Cell Endocrinol* 1992; 84:C25-C31.
- (98) Kyossev ZN, Reeves WB. N-glycosylation is not essential for enzyme activity of 11beta-hydroxysteroid dehydrogenase type 2. *Kidney Int* 1997; 52(3):682-686.
- (99) Lakshmi V, Monder C. Extraction of 11 beta-hydroxysteroid dehydrogenase from rat liver microsomes by detergents. *J Steroid Biochem* 1985; 22(3):331-340.
- (100) Lakshmi V, Monder C. Purification and characterisation of the corticosteroid 11 β -dehydrogenase component of the rat liver 11 β -hydroxysteroid dehydrogenase complex. *Endocrinology* 1988; 123:2390-2398.
- (101) Lan NC, Graham BS, Bartter FC, Baxter JD. Binding of steroids to mineralocorticoid receptors: implications for *in vivo* occupancy by glucocorticoids. *J Clin Endocrinol Metab* 1982; 54:332-342.
- (102) Levitz M, Jansen V, Dancis J. The transfer and the metabolism of corticosteroids in the perfused human placenta. *Am J Obstet Gynecol* 1978; 132:363-366.
- (103) Li KX, Obeyesekere VR, Krozowski ZS, Ferrari P. Oxoreductase and dehydrogenase activities of the human and rat 11beta- hydroxysteroid dehydrogenase type 2 enzyme. *Endocrinology* 1997; 138(7):2948-2952.
- (104) Liu YJ, Nakagawa Y, Nasuda K, Saegusa H, Igarashi Y. Effect of growth hormone, insulin and dexamethasone on 11 beta-hydroxysteroid dehydrogenase activity on a primary culture of rat hepatocytes. *Life Sci* 1996; 59(3):227-234.
- (105) Lovati E, Ferrari P, Dick B, Jostarndt K, Frey BM, Frey FJ et al. Molecular basis of human salt sensitivity: the role of the 11beta-hydroxysteroid dehydrogenase type 2. *J Clin Endocrinol Metab* 1999; 84(10):3745-3749.
- (106) Low SC, Chapman KE, Edwards CR, Seckl JR. 'Liver-type' 11 beta-hydroxysteroid dehydrogenase cDNA encodes reductase but not dehydrogenase activity in intact mammalian COS-7 cells. *J Mol Endocrinol* 1994; 13(2):167-174.
- (107) Low SC, Moisan MP, Noble JM, Edwards CR, Seckl JR. Glucocorticoids regulate hippocampal 11 beta-hydroxysteroid dehydrogenase activity and gene expression in vivo in the rat. *J Neuroendocrinol* 1994; 6(3):285-290.
- (108) Lowy MT, Meltzer HY. Dexamethasone bioavailability: implications for DST research. *Biol Psychiatry* 1987; 22(3):373-385.

- (109) Mantero F, Palermo M, Petrelli MD, Tedde R, Stewart PM, Shackleton CH. Apparent mineralocorticoid excess: type I and type II. *Steroids* 1996; 61(4):193-196.
- (110) Maser E, Volker B, Friebertshauser J. 11 Beta-hydroxysteroid dehydrogenase type 1 from human liver: dimerization and enzyme cooperativity support its postulated role as glucocorticoid reductase. *Biochemistry* 2002; 41(7):2459-2465.
- (111) Masuzaki H, Paterson J, Shinya H, Morton NM, Mullins JJ, Seckl JR et al. A Transgenic Model of Visceral Obesity and the Metabolic Syndrome. *Science* 2001; 294(5549):2166-2170.
- (112) Michael AE, Evangelatou M, Norgate DP, Clarke RJ, Antoniw JW, Stedman BA et al. Isoforms of 11 β -hydroxysteroid dehydrogenase in human granulosa-lutein cells. *Mol Cell Endocrinol* 1997; 132:43-52.
- (113) Minagawa K, Kasuya Y, Baba S, Knapp G, Skelly J. Identifikation und quantifikation of 6 β -hydroxy-dexamethasone as major urine metabolite of Dexamethasone in man. *Steroids* 1986; 47:175-188.
- (114) Monder C, Hardy MP, Blanchard RJ, Bland R. Comparative aspects of 11 β -hydroxysteroid dehydrogenase. Testicular 11 β -hydroxysteroid dehydrogenase: development of a model for the mediation of Leydig cell function by corticosteroids. *Steroids* 1994; 59:69-73.
- (115) Monder C, Lakshmi V, Miroff Y. Kinetic studies on rat liver 11 beta-hydroxysteroid dehydrogenase. *Biochim Biophys Acta* 1991; 1115(1):23-29.
- (116) Monder C, Stewart PM, Lakshmi V, Valentino R, Burt D, Edwards CR. Licorice inhibits corticosteroid 11 beta-dehydrogenase of rat kidney and liver: in vivo and in vitro studies. *Endocrinology* 1989; 125(2):1046-1053.
- (117) Morita H, Zhou M, Foecking MF, Gomez-Sanchez EP, Cozza EN, Gomez-Sanchez CE. 11 beta-Hydroxysteroid dehydrogenase type 2 complementary deoxyribonucleic acid stably transfected into Chinese hamster ovary cells: specific inhibition by 11 alpha-hydroxyprogesterone. *Endocrinology* 1996; 137(6):2308-2314.
- (118) Morris DJ, Souness GW. Endogenous 11 beta-hydroxysteroid dehydrogenase inhibitors and their role in glucocorticoid Na⁺ retention and hypertension. *Endocr Res* 1996; 22(4):793-801.
- (119) Morton NM, Holmes MC, Fievet C, Staels B, Tailleux A, Mullins JJ et al. Improved lipid and lipoprotein profile, hepatic insulin sensitivity, and glucose tolerance in 11beta-hydroxysteroid dehydrogenase type 1 null mice. *J Biol Chem* 2001; 276(44):41293-41300.
- (120) Mune T, Rogerson FM, Nikkila H, Agarwal AK, White PC. Human hypertension caused by mutations in the kidney isozyme of 11 beta-hydroxysteroid dehydrogenase. *Nat Genet* 1995; 10(4):394-399.
- (121) Murphy D, West H, Bethel A. Some aspects of the metabolism and disposition of betamethason. *Acta Endocrinologica* 1964; 45:498-508.

- (122) Naray-Fejes-Toth A, Fejes-Toth G. Subcellular localisation of the type 2 11 β -hydroxysteroid dehydrogenase - A green fluorescent protein study. *J Biol Chem* 1996; 271:15436-15442.
- (123) New MI, Levine LS, Biglieri EG, Pareira J, Ulick S. Evidence for an unidentified Steroid in a Child with Apparent Mineralocorticoid Hypertension. *J Clin Endocrinol Metab* 1977; 44:924-933.
- (124) Obeyesekere V, Li KX, Ferrari P, Krozowski Z. Truncation of the N- and C-terminal regions of the human 11 β -hydroxysteroid dehydrogenase type 2 enzyme and effects on solubility and bidirectional enzyme activity. *Mol Cell Endocrinol* 1997; 131:173-182.
- (125) Odermatt A, Arnold P, Frey FJ. The intracellular localization of the mineralocorticoid receptor is regulated by 11beta-hydroxysteroid dehydrogenase type 2. *J Biol Chem* 2001; 276(30):28484-28492.
- (126) Oelkers W, Buchen S, Diederich S, Krain J, Muhme S, Schoneshofer M. Impaired renal 11 beta-oxidation of 9 alpha-fluorocortisol: an explanation for its mineralocorticoid potency. *J Clin Endocrinol Metab* 1994; 78(4):928-932.
- (127) Oppermann UC, Netter KJ, Maser E. Cloning and primary structure of murine 11 beta-hydroxysteroid dehydrogenase/microsomal carbonyl reductase. *Eur J Biochem* 1995; 227(1-2):202-208.
- (128) Oppermann UC, Persson B, Johanssen S. The 11 β -hydroxysteroid dehydrogenase system, a determinant of glucocorticoid and mineralocorticoid action. *Eur J Biochem* 1997; 249:355-360.
- (129) Palermo M, Shackleton CH, Mantero F, Stewart PM. Urinary free cortisone and the assessment of 11 beta-hydroxysteroid dehydrogenase activity in man. *Clin Endocrinol (Oxf)* 1996; 45(5):605-611.
- (130) Perschel FH, Buhler H, Hierholzer K. Bile acids and their amides inhibit 11 beta-hydroxysteroid dehydrogenase obtained from rat kidney. *Pflugers Arch* 1991; 418(6):538-543.
- (131) Petrelli MD, Lim-Tio SS, Condon J, Hewison M, Stewart PM. Differential expression of nuclear 11 β -hydroxysteroid dehydrogenase type 2 in mineralocorticoid receptor positive and negative tissues. *Endocrinology* 1997; 138:3077-3080.
- (132) Phillipov G, Palermo M, Shackleton CH. Apparent cortisone reductase deficiency: a unique form of hypercortisolism. *J Clin Endocrinol Metab* 1996; 81(11):3855-3860.
- (133) Pinder R, Brogden R, Sawyer P, Speight T, Spencer R, Avery GS. Carbenoxolone: A review of its pharmacological properties and therapeutic efficacy in peptic ulcer disease. *Drugs* 1976; 11:245-307.
- (134) Ponz de Leon M, Loria P, Carulli N, Murphy G, Dowling R. Intestinal solubilization, absorption, pharmacokinetics and bioavailability of chenodesoxycholic acid. *Eur J Clin Invest* 1980; 10:261-271.

- (135) Quinkler M, Johanssen S, Grossmann C, Bahr V, Muller M, Oelkers W et al. Progesterone metabolism in the human kidney and inhibition of 11beta-hydroxysteroid dehydrogenase type 2 by progesterone and its metabolites. *J Clin Endocrinol Metab* 1999; 84(11):4165-4171.
- (136) Quinkler M, Kosmale B, Bahr V, Oelkers W, Diederich S. Evidence for isoforms of 11 beta-hydroxysteroid dehydrogenase in the liver and kidney of the guinea pig. *J Endocrinol* 1997; 153(2):291-298.
- (137) Quinkler M, Oelkers W, Diederich S. In vivo measurement of renal 11beta-hydroxysteroid dehydrogenase type 2 activity. *J Clin Endocrinol Metab* 2000; 85(12):4921-4923.
- (138) Quinkler M, Troeger H, Eigendorff E, Maser-Gluth C, Stiglic A, Oelkers W et al. Enhanced 11beta-hydroxysteroid dehydrogenase type 1 activity in stress adaptation in the guinea pig. *J Endocrinol* 2003; 176(2):185-192.
- (139) Rajan V, Edwards CR, Seckl JR. 11 beta-Hydroxysteroid dehydrogenase in cultured hippocampal cells reactivates inert 11-dehydrocorticosterone, potentiating neurotoxicity. *J Neurosci* 1996; 16(1):65-70.
- (140) Ricketts M, Shoesmith KJ, Hewison M, Strain A, Stewart PM. Immunolocalisation of type 1 11 β -hydroxysteroid dehydrogenase in human tissues. *J Endocrinol* 1998; 156S:279.
- (141) Ricketts ML, Shoesmith KJ, Hewison M, Strain A, Eggo MC, Stewart PM. Regulation of 11 beta-hydroxysteroid dehydrogenase type 1 in primary cultures of rat and human hepatocytes. *J Endocrinol* 1998; 156(1):159-168.
- (142) Rogerson FM, Dimopoulos N, Sluka P, Chu S, Curtis AJ, Fuller PJ. Structural determinants of Aldosterone Binding Selectivity in the mineralocorticoid receptor. *J Biol Chem* 1999; 274(51):36305-36311.
- (143) Roland BL, Li KX, Funder JW. Hybridization histochemical localization of 11 beta-hydroxysteroid dehydrogenase type 2 in rat brain. *Endocrinology* 1995; 136(10):4697-4700.
- (144) Rupprecht R, Arriza JL, Spengler D, Reul JM, Evans R, Holboer F et al. Transactivation and synergistic properties of the mineralocorticoid receptor:relationship to the glucocorticoid receptor. *Mol Endocrinol* 1993; 7:597-603.
- (145) Sasano N, Frost AR, Saito S, Matsunaga G, Nagura H, Krozowski Z et al. Localisation of mineralocorticoid receptor and 11 β -hydroxysteroid dehydrogenase type 2 in human breast and its disorders. *Anticancer Res* 1997; 17:2001-2007.
- (146) Schoneshofer M, Fenner A. A convenient and efficient method for the extraction and fractionation of steroid hormones from serum or urine. *J Clin Chem Clin Biochem* 1981; 19:71-74.
- (147) Seckl JR, Walker BR. Minireview: 11beta-hydroxysteroid dehydrogenase type 1 - a tissue-specific amplifier of glucocorticoid action. *Endocrinology* 2001; 142(4):1371-1376.

- (148) Shackleton CH, Whitney JO. Use of Sep-pak Cartridges for urinary steroid extraction: evaluation of the method for use prior to gas chromatographic analysis. *Clin Chim Acta* 1980; 107:231-243.
- (149) Shimojo M, Stewart PM. Apparent mineralocorticoid excess syndromes. *J Endocrinol Invest* 1995; 18(7):518-532.
- (150) Shimojo M, Whorwood CB, Stewart PM. 11 beta-Hydroxysteroid dehydrogenase in the rat adrenal. *J Mol Endocrinol* 1996; 17(2):121-130.
- (151) Siebe H, Baude G, Lichtenstein I, Wang D, Buhler H, Hoyer GA et al. Metabolism of dexamethasone: sites and activity in mammalian tissues. *Ren Physiol Biochem* 1993; 16(1-2):79-88.
- (152) Siegenthaler W. *Klinische Pathophysiologie*. 1994.
- (153) Slight SH, Ganjam VK, Gomez-Sanchez CE, Zhou MY, Weber KT. High affinity NAD(+)-dependent 11 beta-hydroxysteroid dehydrogenase in the human heart. *J Mol Cell Cardiol* 1996; 28(4):781-787.
- (154) Souness GW, Brem AS, Morris DJ. 11 beta-Hydroxysteroid dehydrogenase antisense affects vascular contractile response and glucocorticoid metabolism. *Steroids* 2002; 67(3-4):195-201.
- (155) Steppan CM, Bailey ST, Bhat S, Brown EJ, Banerjee RR, Wright CM et al. The hormone resistin links obesity to diabetes. *Nature* 2001; 409(6818):307-312.
- (156) Stewart PM, Krozowski ZS. 11 beta-Hydroxysteroid dehydrogenase. *Vitam Horm* 1999; 57:249-324.
- (157) Stewart PM, Mason JI. Cortisol to cortisone: glucocorticoid to mineralocorticoid. *Steroids* 1995; 60(1):143-146.
- (158) Stewart PM, Murry BA, Mason JI. Human kidney 11 beta-hydroxysteroid dehydrogenase is a high affinity nicotinamide adenine dinucleotide-dependent enzyme and differs from the cloned type I isoform. *J Clin Endocrinol Metab* 1994; 79(2):480-484.
- (159) Stewart PM, Rogerson FM, Mason JI. Type 2 11 beta-hydroxysteroid dehydrogenase messenger ribonucleic acid and activity in human placenta and fetal membranes: its relationship to birth weight and putative role in fetal adrenal steroidogenesis. *J Clin Endocrinol Metab* 1995; 80(3):885-890.
- (160) Stewart PM, Wallace AM, Atherden SM, Shearing CH, Edwards CR. Mineralocorticoid activity of carbenoxolone: contrasting effects of carbenoxolone and liquorice on 11 beta-hydroxysteroid dehydrogenase activity in man. *Clin Sci (Colch)* 1990; 78(1):49-54.
- (161) Stewart PM, Whorwood CB, Mason JI. Type 2 11 beta-hydroxysteroid dehydrogenase in foetal and adult life. *J Steroid Biochem Mol Biol* 1995; 55(5-6):465-471.

- (162) Stokes J, Noble J, Brett L, Phillips C, Seckl JR, O'Brien C et al. Distribution of glucocorticoid and mineralocorticoid receptors and 11beta-hydroxysteroid dehydrogenases in human and rat ocular tissues. *Invest Ophthalmol Vis Sci* 2000; 41(7):1629-1638.
- (163) Suzuki T, Sasano H, Suzuki S, Hirasawa G, Takayama K, Muramatu Y et al. 11 β -Hydroxysteroid dehydrogenase type 2 in human lung: Possible regulator of mineralocorticoid action. *J Clin Endocrinol Metab* 1998; 83:4022-4025.
- (164) Szefler SJ. Pharmakokinetcs of intranasal corticosteroids. *J Allergy Clin Immunol* 108, S26-31. 2001.
- (165) Szefler SJ, Ebling WF, Georgitis JW, Jusko WJ. Methylprednisolone versus prednisolone pharmacokinetics in relation to dose in adults. *Eur J Clin Pharmacol* 1986; 30(3):323-329.
- (166) Tannin GM, Agarwal AK, Monder C, New MI, White PC. The human gene for 11 beta-hydroxysteroid dehydrogenase. Structure, tissue distribution, and chromosomal localization. *J Biol Chem* 1991; 266(25):16653-16658.
- (167) Tröger H et.al. *Exp Clin Endocrinol Diabetes* 2000;91-91.
- (168) Ulick S, Chan CK, Rao KN, Edassery J, Mantero F. A new form of the syndrome of apparent mineralocorticoid excess. *J Steroid Biochem* 1989; 32(1B):209-212.
- (169) Ulick S, Levine LS, Gunczler P, Zanconato G, Ramirez LC, Rauh W et al. A syndrome of apparent mineralocorticoid excess associated with defects in the peripheral metabolism of cortisol. *J Clin Endocrinol Metab* 1979; 49(5):757-764.
- (170) Ulick S, Wang JZ, Blumenfeld JD, Pickering TG. Cortisol inactivation overload: a mechanism of mineralocorticoid hypertension in the ectopic adrenocorticotropin syndrome. *J Clin Endocrinol Metab* 1992; 74(5):963-967.
- (171) Ulick S, Wang JZ, Hanukoglu A, Rosler A. The effect of carbenoxolone on the peripheral metabolism of cortisol in human patients. *J Lab Clin Med* 1993; 122(6):673-676.
- (172) Valentino R, Tommaselli AP, Savastano S, Stewart PM, Ghiggi MR, Galletti F et al. Alcohol inhibits 11-beta-hydroxysteroid dehydrogenase activity in rat kidney and liver. *Horm Res* 1995; 43(5):176-180.
- (173) Velazquez H, Naray-Fejes-Toth A, Silva T, Andujar E, Reilly RF, Desir GV et al. Rabbit distal convoluted tubule coexpresses NaCl cotransporter and 11 beta-hydroxysteroid dehydrogenase II mRNA. *Kidney Int* 1998; 54(2):464-472.
- (174) Vogeser M, Zachoval R, Jacob K. Serum cortisol/cortisone ratio after Synacthen stimulation. *Clin Biochem* 2001; 34(5):421-425.
- (175) Voice MW, Seckl JR, Edwards CR, Chapman KE. 11 beta-hydroxysteroid dehydrogenase type 1 expression in 2S FAZA hepatoma cells is hormonally regulated: a model system for the study of hepatic glucocorticoid metabolism. *Biochem J* 1996; 317 (Pt 2):621-625.

- (176) Walker B, Phillips ID, Noon JP, Panarelli M, Padfield PL, Edwards C. Increased vasoconstrictor sensitivity to glucocorticoid in essential hypertension. *Hypertension* 1998; (31):891-895.
- (177) Walker BR, Connacher AA, Lindsay RM, Webb DJ, Edwards CR. Carbenoxolone increases hepatic insulin sensitivity in man: a novel role for 11-oxosteroid reductase in enhancing glucocorticoid receptor activation. *J Clin Endocrinol Metab* 1995; 80(11):3155-3159.
- (178) Walker BR, Stewart PM, Shackleton CH, Padfield PL, Edwards CR. Deficient inactivation of cortisol by 11 beta-hydroxysteroid dehydrogenase in essential hypertension. *Clin Endocrinol (Oxf)* 1993; 39(2):221-227.
- (179) Watson B, Jr., Bergman SM, Myracle A, Callen DF, Acton RT, Warnock DG. Genetic association of 11 beta-hydroxysteroid dehydrogenase type 2 (HSD11B2) flanking microsatellites with essential hypertension in blacks. *Hypertension* 1996; 28(3):478-482.
- (180) White PC, Mune T, Rogerson FM, Kayes KM, Agarwal AK. 11 beta-Hydroxysteroid dehydrogenase and its role in the syndrome of apparent mineralocorticoid excess. *Pediatr Res* 1997; 41(1):25-29.
- (181) White PC, Mune T, Rogerson FM, Kayes KM, Agarwal AK. Molecular analysis of 11 beta-hydroxysteroid dehydrogenase and its role in the syndrome of apparent mineralocorticoid excess. *Steroids* 1997; 62(1):83-88.
- (182) Whorwood CB, Franklyn JA, Sheppard MC, Stewart PM. Tissue localization of 11 beta-hydroxysteroid dehydrogenase and its relationship to the glucocorticoid receptor. *J Steroid Biochem Mol Biol* 1992; 41(1):21-28.
- (183) Whorwood CB, Mason JI, Ricketts M, Howie A, Stewart PM. Detection of human 11 β -hydroxysteroid dehydrogenase isoforms using reverse transcriptas-polymerase chain reaction and localisation of the type 2 isoform to renal collecting ducts. *Mol Cell Endocrinol* 1995; 110:R7-R12.
- (184) Whorwood CB, Ricketts ML, Stewart PM. Epithelial cell localization of type 2 11 beta-hydroxysteroid dehydrogenase in rat and human colon. *Endocrinology* 1994; 135(6):2533-2541.
- (185) Whorwood CB, Sheppard MC, Stewart PM. Licorice inhibits 11 beta-hydroxysteroid dehydrogenase messenger ribonucleic acid levels and potentiates glucocorticoid hormone action. *Endocrinology* 1993; 132(6):2287-2292.
- (186) Wilson RC, Nimkarn S, New MI. Apparent mineralocorticoid excess. *Trends Endocrinol Metab* 2001; 12(3):104-111.
- (187) Yau JL, Noble J, Kenyon CJ, Hibberd C, Kotelevtsev Y, Mullins JJ et al. Lack of tissue glucocorticoid reactivation in 11beta -hydroxysteroid dehydrogenase type 1 knockout mice ameliorates age-related learning impairments. *Proc Natl Acad Sci U S A* 2001; 98(8):4716-4721.