

BIBLIOGRAPHY

- 1 Petit, M.C., Benkirane, N., Guichard, G., Du, A.P., Marraud, M., Cung, M.T., *et al.* Solution structure of a retro-inverso peptide analogue mimicking the foot-and-mouth disease virus major antigenic site. Structural basis for its antigenic cross-reactivity with the parent peptide. *Journal of Biological Chemistry* 1999, **274**, 3686-3692
- 2 Nielsen, P.E. Applications of peptide nucleic acids. *Current Opinion in Biotechnology* 1999, **10**, 71-75
- 3 Houghten, R.A. General method for the rapid solid-phase synthesis of large numbers of peptides: specificity of antigen-antibody interaction at the level of individual amino acids. *Proc. Natl. Acad. Sci. USA* 1985, **82**, 5131-5135
- 4 Geysen, H.M., Meloen, R.H. and Barteling, S.J. Use of peptide synthesis to probe viral antigens for epitopes to a resolution of a single amino acid. *Proc. Natl. Acad. Sci. (U. S. A.)* 1984, **81**, 3998-4002
- 5 Adler, S., Frank, R., Lanzavecchia, A. and Weiss, S. T cell epitope analysis with peptides simultaneously synthesized on cellulose membranes: fine mapping of two DQ dependent epitopes. *FEBS Letters* 1994, **352**, 167-170
- 6 Jung, C., Kalbus, M., Fleckenstein, B., Melms, A., Jung, G. and Wiesmuller, K.H. New ligands for HLA DRB1 * 0301 by random selection of favourable amino acids ranked by competition studies with undecapeptide amide sublibraries. *Journal of Immunological Methods* 1998, **219**, 139-149
- 7 Tarbit, M.H. and Berman, J. High-throughput approaches for evaluating absorption, distribution, metabolism and excretion properties of lead compounds. *Current Opinion in Chemical Biology* 1998, **2**, 411-416
- 8 Nicolaou, K.C., Trujillo, J.I., Jandeleit, B., Chibale, K., Rosenfeld, M., Diefenbach, B., *et al.* Design, synthesis and biological evaluation of nonpeptide integrin antagonists. *Bioorganic & Medicinal Chemistry* 1998, **6**, 1185-1208
- 9 Kempf, D.J., Marsh, K.C., Denissen, J.F., McDonald, E., Vasavanonda, Flentge, C.A., *et al.* ABT-538 is a potent inhibitor of human immunodeficiency virus protease and has high oral bioavailability in humans. *Proceedings of the National Academy of Sciences of the United States of America* 1995, **92**, 2484-2488
- 10 Rose, K. Facile synthesis of homogeneous artificial proteins. *Journal of the American Chemical Society* 1994, **116**, 30-33
- 11 Tuchscherer, G. and Mutter, M. Templates in protein de novo design. *Journal of Biotechnology* 1995, **41**, 197-210
- 12 Dawson, P.E., Muir, T.W., Clark-Lewis, I. and Kent, S.B.H. Synthesis of proteins by native chemical ligation. *Science* 1994, **266**, 776-779
- 13 Ferencik, M. *Handbook of Immunochemistry*. Chapman & Hall, London, 1993

- 14 Celis, J.E., Leffers, H., Rasmussen, H.H., Madsen, P., Honore, B., Gesser, B., *et al.* The master two-dimensional gel database of human AMA cell proteins: towards linking protein and genome sequence and mapping information (update 1991). *Electrophoresis* 1991, **12**, 765-801
- 15 Adams, M.D., Kerlavage, A.R., Fleischmann, R.D., Fuldner, R.A., Bult, C.J., Lee, N.H., *et al.* Initial assessment of human gene diversity and expression patterns based upon 83 million nucleotides of cDNA sequence. *Nature* 1995, **377**, 3-174
- 16 Hudson, T.J., Stein, L.D., Gerety, S.S., Ma, J.L., Castle, A.B., Silva, *et al.* An STS-based map of the human genome. *Science* 1995, **270**, 1945-1954
- 17 Berna, A., Godefroy, C.T., Stussi and Garaud, C. Preparation of an antiserum against an *in-vitro* translation product of Alfalfa Mosaic Virus RNA 3. *Journal of General Virology* 1985, **66**, 1669-1678
- 18 Schoumacher, F., Erny, C., Berna, A., Godefroy-Colburn, T. and Stussi-Garaud, C. Nucleic acid-binding properties of the alfalfa mosaic virus movement protein produced in yeast. *Virology* 1992, **188**, 896-899
- 19 Walter, G. Production and use of antibodies against synthetic peptides. *J. Immunol. Methods* 1986, **88**, 149-162
- 20 Walter, G., Scheidtmann, K.H., Carbone, A., Laudano, A.P. and Doolittle, R.F. Antibodies specific for the carboxy- and amino-terminal regions of simian virus 40 large tumor antigen. *Proceedings of the National Academy of Sciences of the United States of America* 1980, **77**, 5197-5200
- 21 Green, N., Alexander, H., Olson, A., Alexander, S., Shinnick, T.M., Sutcliffe, J.G., *et al.* Immunogenic structure of the influenza virus hemagglutinin. *Cell* 1982, **28**, 477-487
- 22 DiMarchi, R., Brooke, G., Gale, C., Cracknell, V., Doel, T. and Mowat, N. Protection of cattle against foot-and-mouth disease by a synthetic peptide. *Science* 1986, **232**, 639-641
- 23 Roop, D.R., Cheng, C.K., Titterington, L., Meyers, C.A., Stanley, JR, *et al.* Synthetic peptides corresponding to keratin subunits elicit highly specific antibodies. *Journal of Biological Chemistry* 1984, **259**, 8037-8040
- 24 Brown, C.D., Barnes, K. and Turner, A.J. Anti-peptide antibodies specific to rat endothelin-converting enzyme-1 isoforms reveal isoform localisation and expression. *FEBS Letters* 1998, **424**, 183-187
- 25 Grazioli, L., Germain, V., Weiss, A. and Acuto, O. Anti-peptide antibodies detect conformational changes of the inter-SH2 domain of ZAP-70 due to binding to the zeta chain and to intramolecular interactions. *Journal of Biological Chemistry* 1998, **273**, 8916-8921
- 26 Wallukat, G., Fu, M.L.X., Magnusson, Y., Hjalmarson, A., Hoebeke, J. and Wollenberger. Agonistic effects of anti-peptide antibodies and autoantibodies directed against adrenergic and cholinergic receptors: Absence of desensitization. *Blood Pressure* 1996, **5**, 31-36
- 27 Lerner, R.A. Tapping the immunological repertoire to produce antibodies of predetermined specificity. *Nature* 1982, **299**, 593-596

- 28 Baron, M.H. and Baltimore, D. Antibodies against a synthetic peptide of the poliovirus replicase protein: reaction with native, virus-encoded proteins and inhibition of virus-specific polymerase activities in vitro. *Journal of Virology* 1982, **43**, 969-978
- 29 Semler, B.L., Anderson, C.W., Hanecak, R., Dorner, L.F. and Wimmer, E. A membrane-associated precursor to poliovirus VPg identified by immunoprecipitation with antibodies directed against a synthetic heptapeptide. *Cell* 1982, **28**, 405-412
- 30 Murakami, T., Simonds, W.F. and Spiegel, A.M. Site-specific antibodies directed against G protein beta and gamma subunits: effects on alpha and beta gamma subunit interaction. *Biochemistry* 1992, **31**, 2905-2911
- 31 Tamura, R.N., Cooper, H.M., Collo, G. and Quaranta, V. Cell type-specific integrin variants with alternative alpha chain cytoplasmic domains. *Proceedings of the National Academy of Sciences of the United States of America* 1991, **88**, 10183-10187
- 32 Hanecak, R., Semler, B.L., Ariga, H., Anderson, C.W. and Wimmer, E. Expression of a cloned gene segment of poliovirus in *E. coli*: evidence for autocatalytic production of the viral proteinase. *Cell* 1984, **37**, 1063-1073
- 33 Baron, M.H. and Baltimore, D. Anti-VPg antibody inhibition of the poliovirus replicase reaction and production of covalent complexes of VPg-related proteins and RNA. *Cell* 1982, **30**, 745-752
- 34 Neurath, A.R., Kent, S.B., Strick, N., Taylor, P. and Stevens, C.E. Hepatitis B virus contains pre-S gene-encoded domains. *Nature* 1985, **315**, 154-156
- 35 Woulfe, J., Lafourche, L., de Nadai, F., Kitabgi, P. and Beaudet, A. Post-translational processing of the neuropeptidyl/peptidomodulin N precursor in the central nervous system of the rat--II. Immunohistochemical localization of maturation products. *Neuroscience* 1994, **60**, 167-181
- 36 Loffler, J., Langui, D., Probst, A. and Huber, G. Accumulation of a 50 kDa N-terminal fragment of beta-APP695 in Alzheimer's disease hippocampus and neocortex. *Neurochemistry International* 1994, **24**, 281-288
- 37 Stein, P., Worrad, D.M., Belyaev, N.D., Turner, B.M. and Schultz, R.M. Stage-dependent redistributions of acetylated histones in nuclei of the early preimplantation mouse embryo. *Molecular Reproduction & Development* 1997, **47**, 421-429
- 38 Fleckenstein, B. Eberhard-Karls-Universität, Institut für organische Chemie, Auf der Morgenstelle 18, 72076 Tübingen, Germany (personal communication)
- 39 Fruton, J.S. From peptones to peptides. A historical retrospect. Peptides, 1986; 19th European Symposium, Chalkidiki, Greece. 1987, 25-34
- 40 du Vigenaud, V., Ressler, C., Swann, J.M., Roberts, C.W. and Katsoyannis, P.G. The synthesis of oxytocin. *Journal of the American Chemical Society* 1954, **76**, 3115-3124
- 41 Meienhofer, J., Schnable, E., Bremer, H., Brinkhoff, O., Zabel, R. and Sroka, W. Synthese der Insulinketten und ihre Kombination zu insulinaktiven Präparaten. *Zeitung Naturforschung* 1963, **18b**, 1120-1131

- 42 Merrifield, R.B. Peptide synthesis on a solid polymer. *Fed. Proc. Fed. Amer. Soc. Exp. Biol.* 1962, **21**, 412-419
- 43 Merrifield, R.B. Solid phase peptide synthesis. I. The synthesis of a tetrapeptide. *J. Amer. Chem. Soc.* 1963, **85**, 2149-2154
- 44 Merrifield, R.B. Solid phase peptide synthesis. 3. Improved synthesis of Bradykinin. *Biochemistry* 1964, **3**, 1385-1390
- 45 Carpino, L.A. and Han, G.Y. 9-fluorenylmethyloxycarbonyl amino-protecting group. *Journal of Organic Chemistry* 1972, **37**, 3404-3406
- 46 Chang, C.D. and Meienhofer, J. Solid-phase peptide synthesis using mild base cleavage of N alpha-fluorenylmethyloxycarbonylamino acids, exemplified by a synthesis of dihydrosomatostatin. *International Journal of Peptide & Protein Research* 1978, **11**, 246-249
- 47 Meienhofer, J., Waki, M., Heimer, E.P., Lambros, T.J., Makofske, RC, et al. Solid phase synthesis without repetitive acidolysis. Preparation of leucyl-alanyl-glycyl-valine using 9-fluorenylmethyloxycarbonylamino acids. *International Journal of Peptide & Protein Research* 1979, **13**, 35-42
- 48 Gait, M.J. and Sheppard, R.C. Rapid synthesis of oligodeoxyribonucleotides: a new solid-phase method. *Nucleic Acids Research* 1977, **4**, 1135-1158
- 49 Young, C.R. and Atassi, M.Z. T-cell dependency of the antibody response to free small synthetic peptides of a protein: demonstration with an antigenic site of myoglobin. *Immunological Investigations* 1985, **14**, 1-5
- 50 Young, C.R., Schmitz, H.E. and Atassi, M.Z. Genetic control of the immune response to myoglobin. XVI. Control of antibodies with preselected specificities following immunization with free synthetic peptides representing the antigenic sites or surface non-immunogenic locations in the protein. *Journal of Immunogenetics* 1983, **10**, 453-464
- 51 Antoni, G., Mariani, M., Presentini, R., Lafata, M., Neri, P., Bracci, L., et al. Production and characterization of monoclonal antibodies to anti-human chorionic somatomammotropin by immunization with two free synthetic peptides. *Molecular Immunology* 1985, **22**, 1237-1241
- 52 Atassi, M.Z. and Young, C.R. Discovery and implications of the immunogenicity of free small synthetic peptides: powerful tools for manipulating the immune system and for production of antibodies and T cells of preselected submolecular specificities. *Critical Reviews in Immunology* 1985, **5**, 387-409
- 53 Srinivasan, M., Domanico, S.Z., Kaumaya, P.T.P. and Pierce, S.K. Peptides of 23 residues or greater are required to stimulate a high affinity class II-restricted T cell response. *Eur. J. Immunol.* 1993, **23**, 1011-1016
- 54 Van Regenmortel, M.H.V., Briand, J.P., Muller, S. and Plauè, S. Synthetic polypeptides as antigens. Burdon, R. H. and Van Knippenberg, P. H. (Eds.). Elsevier science publishers, Amsterdam, 1988
- 55 Schulze-Gahmen, U., Klenk, H.D. and Beyreuther, K. Immunogenicity of loop-structured short synthetic peptides mimicking the antigenic site A of influenza virus hemagglutinin. *European Journal of Biochemistry* 1986, **159**, 283-289

- 56 Choppin, J., Metzger, J.J., Bouillot, M., Briand, J.P., Connan, F., Van Regenmortel, M.H., *et al.* Recognition of HLA class I molecules by antisera directed to synthetic peptides corresponding to different regions of the HLA-B7 heavy chain. *Journal of Immunology* 1986, **136**, 1738-1744
- 57 Delmas, A., Milhaud, G., Raulais, D. and Rivaille, P. Epitopes of the 44-68 human parathyroid hormone fragments: the importance of specific hydrophilic peptide sequences. *Molecular Immunology* 1985, **22**, 675-679
- 58 Gendloff, E.H., Casale, W.L., Ram, B.P., Tai, J.H., Pestka, J.J., Hart, *et al.* Hapten-protein conjugates prepared by the mixed anhydride method. Cross-reactive antibodies in heterologous antisera. *Journal of Immunological Methods* 1986, **92**, 15-20
- 59 Modrow, S., Hoflacher, B., Mertz, R. and Wolf, H. Carrier-bound synthetic peptides. Use as antigen in HIV-1 ELISA tests and in antiserum production. *Journal of Immunological Methods* 1989, **118**, 1-7
- 60 Nayak, B.P., Tuteja, R., Manivel, V., Roy, R.P., Vishwakarma, R.A. and Rao, K.V. B cell responses to a peptide epitope. V. Kinetic regulation of repertoire discrimination and antibody optimization for epitope. *Journal of Immunology* 1998, **161**, 3510-3519
- 61 Tam, J.D. Synthetic peptide vaccine design: Synthesis and properties of a high-density multiple antigenic peptide system. *Proc. Natl. Acad. Sci. (U. S. A.)* 1988, **85**, 5409-5413
- 62 Francis, M.J., Hastings, G.Z., Brown, F., McDermed, J., Lu, Y.A. and Tam, J.P. Immunological evaluation of the multiple antigen peptide (MAP) system using the major immunogenic site of foot-and-mouth disease virus. *Immunology* 1991, **73**, 249-254
- 63 Kanda, P., Kennedy, R.C. and Sparrow, J.T. Synthesis of polyamide supports for use in peptide synthesis and as peptide-resin conjugates for antibody production. *International Journal of Peptide & Protein Research* 1991, **38**, 385-391
- 64 Hopp, T.P. Immunogenicity of a synthetic HBsAg peptide: enhancement by conjugation to a fatty acid carrier. *Molecular Immunology* 1984, **21**, 13-16
- 65 Huang, W., Nardelli, B. and Tam, J.P. Lipophilic multiple antigen peptide system for peptide immunogen and synthetic vaccine. *Mol. Immunol.* 1994, **31**, 1191-1199
- 66 Nardelli, B., Haser, P.B. and Tam, J.P. Oral administration of an antigenic synthetic lipopeptide (MAP-P3C) evokes salivary antibodies and systemic humoral and cellular responses. *Vaccine* 1994, **12**, 1335-1339
- 67 Ahlers, J.D., Takeshita, T., Pendleton, C.D. and Berzofsky, J.A. Enhanced immunogenicity of HIV-1 vaccine construct by modification of the native peptide sequence. *Proceedings of the National Academy of Sciences of the United States of America* 1997, **94**, 10856-10861
- 68 Rizo, J. and Giersch, L.M. Constrained peptides: models of bioactive peptides and protein substructures. *Annu. Rev. Biochem.* 1992, **61**, 387-418
- 69 Chin, L.-T., Hinkula, J., Levi, M., Ohlin, M., Wahren, B. and Borrebaeck, C.A.K. Site-directed primary *in vitro* immunization: Production of HIV-1

- neutralizing human monoclonal antibodies from lymphocytes obtained from seronegative donors. *Immunol.* 1994, **81**, 428-434
- 70 Kanda, P., Fritz, D.A., Gage, D.A. and Shuler, K.R. Dependence of the murine antibody response to an anti-CD4 CDR2 V_H peptide on immunogen formulation. *Mol. Immunol.* 1995, **32**, 1319-1328
- 71 Shen, Z., Reznikoff, G., Dranoff, G. and Rock, K.L. Cloned dendritic cells can present exogenous antigens on both MHC class I and class II molecules. *Journal of Immunology* 1997, **158**, 2723-2730
- 72 Groettrup, M., Soza, A., Kuckelkorn, U. and Kloetzel, P.M. Peptide antigen production by the proteasome: complexity provides efficiency. *Immunology Today* 1996, **17**, 429-435
- 73 Nardin, E.H., Calvocalle, J.M., Oliveira, G.A., Clavijo, P., Nussenzweig, R., Simon, R., et al. *Plasmodium falciparum* polyoximes - highly immunogenic synthetic vaccines constructed by chemoselective ligation of repeat B-cell epitopes and a universal T-cell epitope of CS protein. *Vaccine* 1998, **16**, 590-600
- 74 Ho, P.C., Mutch, D.A., Winkel, K.D., Saul, A.J., Jones, G.L., Doran, T.J., et al. Identification of two promiscuous T cell epitopes from tetanus toxin. *European Journal of Immunology* 1990, **20**, 477-483
- 75 Wisdom, G.B. Peptide Antigens. Oxford University Press, Oxford, 1995
- 76 Samokhin, G.P. and Filimonov, I.N. Coupling of peptides to protein carriers by mixed anhydride procedure. *Analytical Biochemistry* 1985, **145**, 311-314
- 77 Houen, G. and Jensen, O.M. Conjugation to preactivated proteins using divinylsulfone and iodoacetic acid. *J. Immunol. Methods* 1995, **181**, 187-200
- 78 Hansen, P.R., Holm, A. and Houen, G. Solid-phase peptide synthesis on proteins. *Int. J. Peptide Protein Res.* 1992, **41**, 237-245
- 79 Brumeau, T.D., Dehazya, P., Wolf, I. and Bona, C.A. Enzymatically mediated, glycosidic conjugation of immunoglobulins with viral epitopes. *J. Immunol. Meth.* 1995, **183**, 185-197
- 80 Brumeau, T.-D., Casares, S., Harris, P.E., Dehazya, P., Wolf, I., von Boehmer, H., et al. Immunopotency of a viral peptide assembled to the carbohydrate moieties of self immunoglobulins. *Nature Biotechnology* 1996, **14**, 722-725
- 81 Roman, E. and Moreno, C. Synthetic peptides non-covalently bound to bacterial hsp 70 elicit peptide-specific T-cell responses in vivo. *Immunology* 1996, **88**, 487-492
- 82 Udon, H. and Srivastava, P.K. Heat shock protein 70-associated peptides elicit specific cancer immunity. *Journal of Experimental Medicine* 1993, **178**, 1391-1396
- 83 Young, D., Roman, E., Moreno, C., O'Brien, R. and Born, W. Molecular chaperones and the immune response. *Philosophical Transactions of the Royal Society of London - Series B: Biological Sciences* 1993, **339**, 363-367

- 84 Tsao, J., Lin, X., Lackalndn, H., Tous, G., Wu, Y. and Stein, S. Internally standardized amino acid analysis for determining peptide-carrier protein coupling ratio. *Analytical Biochemistry* 1991, **197**, 137-142
- 85 Kolodny, N. and Robey, F.A. Conjugation of synthetic peptides to proteins: quantitation from S-carboxymethylcysteine released upon acid hydrolysis. *Analytical Biochemistry* 1990, **187**, 136-140
- 86 Jackson, D.C., O'Brien-Simpson, N., Ede, N.J. and Brown, L.E. Free radical induced polymerization of synthetic peptides into polymeric immunogens. *Vaccine* 1997, **15**, 1697-1705
- 87 Jackson, D.C., Fitzmaurice, C., Sheppard, R.C., McMurray, J. and Brown, L.E. Antigenic and immunogenic properties of synthetic peptide-based T-cell determinant polymers. *Biomedical Peptides, Proteins & Nucleic Acids* 1995, **1**, 171-176
- 88 Kohn, J., Niemi, S.M., Albert, E.C., Murphy, J.C., Langer, R., Fox, *et al.* Single-step immunization using a controlled release, biodegradable polymer with sustained adjuvant activity. *Journal of Immunological Methods* 1986, **95**, 31-38
- 89 Eldridge, J.H., Meulbroek, J.A., Staas, J.K., Tice, T.R. and Gilley, R.M. Vaccine-containing biodegradable microspheres specifically enter the gut-associated lymphoid tissue following oral administration and induce a disseminated mucosal immune response. *Advances in Experimental Medicine & Biology* 1989, **251**, 191-202
- 90 Wiesmuller, K.H., Jung, G. and Hess, G. Novel low-molecular-weight synthetic vaccine against foot-and-mouth disease containing a potent B-cell and macrophage activator. *Vaccine* 1989, **7**, 29-33
- 91 Allison, A.C. and Gregoriadis, G. Liposomes as immunological adjuvants. *Recent Results in Cancer Research* 1976, 58-64
- 92 Gregoriadis, G. Engineering liposomes for drug delivery: progress and problems. *Trends In Biotechnology* 1995, **13**, 527-537
- 93 Cevc, G., Gebauer, D., Stieber, J., Schatzlein, A. and Blume, G. Ultraflexible vesicles, Transfersomes, have an extremely low pore penetration resistance and transport therapeutic amounts of insulin across the intact mammalian skin. *Biochimica et Biophysica Acta* 1998, **1368**, 201-215
- 94 Paul, A., Cevc, G. and Bachhawat, B.K. Transdermal immunisation with an integral membrane component, gap junction protein, by means of ultradeformable drug carriers, transfersomes. *Vaccine* 1998, **16**, 188-195
- 95 Gregoriadis, G. Immunological adjuvants: a role for liposomes. *Immunol. Today* 1990, **11**, 89-96
- 96 Alving, C.R., Richards and R.L. *Liposomes*. Marcel Dekker, New York, 1983, p. 209
- 97 Brynestad, K., Babbit, B., Huang, L. and Rouse, B.T. Influence of peptide acylation, liposome incorporation, and synthetic immunomodulators on the immunogenicity of a 1-23 peptide of glycoprotein D of herpes simplex virus: implications for subunit vaccines. *Journal of Virology* 1990, **64**, 680-685

- 98 Neurath, A.R., Kent, S.B. and Strick, N. Antibodies to hepatitis B surface antigen (HBsAg) elicited by immunization with a synthetic peptide covalently linked to liposomes. *Journal of General Virology* 1984, **65**, 1009-1014
- 99 Friede, M., Muller, S., Briand, J.-P., Van Regenmortel, M.H.V. and Schuber, F. Induction of immune response against a short synthetic peptide antigen coupled to small neutral liposomes containing monophosphoryl lipid A. *Mol. Immunol.* 1993, **30**, 539-547
- 100 Morein, B., Sundquist, B., Hoglund, S., Dalsgaard, K. and Osterhaus. Iscom, a novel structure for antigenic presentation of membrane proteins from enveloped viruses. *Nature* 1984, **308**, 457-460
- 101 Takahashi, H., Takeshita, T., Morein, B., Putney, S., Germain, RN, *et al.* Induction of CD8+ cytotoxic T cells by immunization with purified HIV-1 envelope protein in ISCOMs. *Nature* 1990, **344**, 873-875
- 102 Taussig, M.J. Studies on the induction of immunological tolerance. The inhibition of tolerance-induction by antiserum: split tolerance and the time-course of tolerance induction. *European Journal of Immunology* 1971, **1**, 367-371
- 103 Sarnesto, A. Monoclonal antibodies reacting with isotypic and/or allotypic determinants of human IgG. *Med. Biol.* 1983, **61**, 126-132
- 104 Herzenberg, L.A. and Tokuhisa, T. Epitope-specific regulation. III. Induction of allotype-restricted suppression for IgG antibody responses to individual epitopes on complex antigens. *European Journal of Immunology* 1982, **12**, 814-818
- 105 Ladner, R.C. Constrained peptides as binding entities. *Trends Biotechnol.* 1995, **13**, 426-430
- 106 Marqusee, S. and Baldwin, R.L. Helix stabilization by Glu-...Lys+ salt bridges in short peptides of de novo design. *Proceedings of the National Academy of Sciences of the United States of America* 1987, **84**, 8898-8902
- 107 Gurunath, R., Beena, T.K., Adiga, P.R. and Balaram, P. Enhancing peptide antigenicity by helix stabilization. *FEBS Letters* 1995, **361**, 176-178
- 108 Posnett, D.N., McGrath, H. and Tam, J.P. A novel method for producing anti-peptide antibodies. *J. Biol. Chem.* 1988, **263**, 1719-1725
- 109 Mutter, M. and Vuilleumier, S. A chemical approach to protein design - template-assembled synthetic proteins (TASP). *Angew. Chem. Int. Ed. Engl.* 1995, **28**, 535-554
- 110 Viamontes, G.I., Audhya, T. and Goldstein, G. Antibodies to thymopoietin following implantation of paper disks derivatized with synthetic Cys-thymopoietin. *Journal of Immunological Methods* 1986, **94**, 13-17
- 111 Smith, R.C., Neff, A.W. and Malacinski, G.M. Accumulation, organization and deployment of oogenetically derived Xenopus yolk/nonyolk proteins. *Journal of Embryology & Experimental Morphology* 1986, **97**, 45-64
- 112 Ramon, G. Sur l'augmentation anormale de l'antitoxine chez les chevaux producteurs de serum antidiphérique. *Bull. Soc. Centr. Med. Vet.* 1925, **101**, 227-234

- 113 Ramon, G. Procedres pour acroitre la production des antitoxins. *Ann. Institut Pasteur* 1926, **40**, 1-10
- 114 Goding, J.W. Monoclonal antibodies: Principles and practice. Academic Press, London, 1986
- 115 Gupta, R.K. and Siber, G.R. Adjuvants for human vaccines--current status, problems and future prospects. *Vaccine* 1995, **13**, 1263-1276
- 116 Lin, R., Tarr, P.E. and Jones, T.C. Present status of the use of cytokines as adjuvants with vaccines to protect against infectious diseases. *Clin. Infect. Dis.* 1995, **21**, 1439-1449
- 117 Soltysik, S., Bedore, D.A. and Kensil, C.R. Adjuvant activity of QS-21 isomers. *Annals of the New York Academy of Sciences* 1993, **690**, 392-395
- 118 Newman, M.J., Wu, J.Y., Gardner, B.H., Anderson, C.A., Kensil, C.R., Recchia, J., et al. Induction of cross-reactive cytotoxic T-lymphocyte responses specific for HIV-1 gp120 using saponin adjuvant (QS-21) supplemented subunit vaccine formulations. *Vaccine* 1997, **15**, 1001-1007
- 119 Audibert, F.M. and Lise, L.D. Adjuvants: current status, clinical perspectives and future prospects. *Immunology Today* 1993, **14**, 281-284
- 120 Gupta, R.K., Relyveld, E.H., Lindblad, E.B., Bizzini, B., Ben-Efraim, S. and Gupta, C.K. Adjuvants--a balance between toxicity and adjuvanticity. *Vaccine* 1993, **11**, 293-306
- 121 Taylor, C.E. Cytokines as adjuvants for vaccines: Antigen-specific responses differ from polyclonal responses. *Infect. Immun.* 1995, **63**, 3241-3244
- 122 Liles, W.C. and Van Voorhis, W.C. Review: nomenclature and biologic significance of cytokines involved in inflammation and the host immune response. *Journal of Infectious Diseases* 1995, **172**, 1573-1580
- 123 Coutelier, J.P., van der Logt, J.T. and Heessen, F.W. IgG subclass distribution of primary and secondary immune responses concomitant with viral infection. *Journal of Immunology* 1991, **147**, 1383-1386
- 124 Gupta, R.K. and Siber, G.R. Comparison of adjuvant activities of aluminium phosphate, calcium phosphate and stearyl tyrosine for tetanus toxoid. *Biologicals* 1994, **22**, 53-63
- 125 Forsthuber, T., Yip, H.C. and Lehmann, P.V. Induction of TH1 and TH2 immunity in neonatal mice. *Science* 1996, **271**, 1728-1730
- 126 Brewer, J.M., Conacher, M., Satoskar, A., Bluethmann, H. and Alexander, J. In interleukin-4-deficient mice, alum not only generates T helper 1 responses equivalent to freund's complete adjuvant, but continues to induce T helper 2 cytokine production. *European Journal of Immunology* 1996, **26**, 2062-2066
- 127 Gasson, J.C. Molecular physiology of granulocyte-macrophage colony-stimulating factor. *Blood* 1991, **77**, 1131-1145
- 128 Fischer, H.G., Frosch, S., Reske, K. and Reske-Kunz, A.B. Granulocyte-macrophage colony-stimulating factor activates macrophages derived from bone marrow cultures to synthesis of MHC class II molecules and to

- augmented antigen presentation function. *Journal of Immunology* 1988, **141**, 3882-3888
- 129 Fischer, H.G., Opel, B., Reske, K. and Reske-Kunz, A.B. Granulocyte-macrophage colony-stimulating factor-cultured bone marrow-derived macrophages reveal accessory cell function and synthesis of MHC class II determinants in the absence of external stimuli. *European Journal of Immunology* 1988, **18**, 1151-1158
- 130 Gerhartz, H.H., Engelhard, M., Meusers, P., Brittinger, G., Wilmanns, W., Schlimok, G., et al. Randomized, double-blind, placebo-controlled, phase III study of recombinant human granulocyte-macrophage colony-stimulating factor as adjunct to induction treatment of high-grade malignant non-Hodgkin's lymphomas. *Blood* 1993, **82**, 2329-2339
- 131 Roilides, E. and Pizzo, P.A. Modulation of host defenses by cytokines: evolving adjuncts in prevention and treatment of serious infections in immunocompromised hosts. *Clinical Infectious Diseases* 1992, **15**, 508-524
- 132 Larsen, C.P., Ritchie, S.C., Hendrix, R., Linsley, P.S., Hathcock, KS, et al. Regulation of immunostimulatory function and costimulatory molecule (B7-1 and B7-2) expression on murine dendritic cells. *Journal of Immunology* 1994, **152**, 5208-5219
- 133 Jones, T.C. Future uses of granulocyte-macrophage colony-stimulating factor (GM-CSF). *Stem Cells* 1994, **12 Suppl 1**, 229-239
- 134 Tao, M.H. and Levy, R. Idiotype/granulocyte-macrophage colony-stimulating factor fusion protein as a vaccine for B-cell lymphoma. *Nature* 1993, **362**, 755-758
- 135 Taylor-Papadimitriou, J. and Balkwill, F. Cancer. Fusion potential for vaccines. *Nature* 1993, **362**, 695
- 136 Dranoff, G., Jaffee, E., Lazenby, A., Columbek, P., Levitsky, H., Brose, K., et al. Vaccination with irradiated tumor cells engineered to secrete murine granulocyte-macrophage colony-stimulating factor stimulates potent, specific, and long-lasting anti-tumor immunity. *Proceedings of the National Academy of Sciences of the United States of America* 1993, **90**, 3539-3543
- 137 Coffman, R.L. and Mosmann, T.R. CD4+ T-cell subsets: regulation of differentiation and function. *Research in Immunology* 1991, **142**, 7-9
- 138 Arai, K.I., Lee, F., Miyajima, A., Miyatake, S., Arai, N. and Yokota. Cytokines: coordinators of immune and inflammatory responses. *Annual Review of Biochemistry* 1990, **59**, 783-836
- 139 Paul, W.E. Interleukin-4: a prototypic immunoregulatory lymphokine. *Blood* 1991, **77**, 1859-1870
- 140 Carter, K.C., Gallagher, G., Baillie, A.J. and Alexander, J. The induction of protective immunity to Leishmania major in the BALB/c mouse by interleukin 4 treatment. *European Journal of Immunology* 1989, **19**, 779-782
- 141 Cameron, M.J., Arreaza, G.A., Zucker, P., Chensue, S.W., Strieter, RM, et al. IL-4 prevents insulitis and insulin-dependent diabetes mellitus in nonobese diabetic mice by potentiation of regulatory T helper-2 cell function. *Journal of Immunology* 1997, **159**, 4686-4692

- 142 elGhazali, G.E., Paulie, S., Andersson, G., Hansson, Y., Holmquist, G., Sun, J.B., *et al.* Number of interleukin-4- and interferon-gamma-secreting human T cells reactive with tetanus toxoid and the mycobacterial antigen PPD or phytohemagglutinin: distinct response profiles depending on the type of antigen used for activation. *European Journal of Immunology* 1993, **23**, 2740-2745
- 143 Palfreyman, J.W., Aitcheson, T.C. and Taylor, P. Guidelines for the production of polypeptide specific antisera using small synthetic oligopeptides as immunogens. *Journal of Immunological Methods* 1984, **75**, 383-393
- 144 Houen, G., Jakobsen, M.H., Svaerke, C., Koch, C. and Barkholt, V. Conjugation to preadsorbed preactivated proteins and efficient generation of anti peptide antibodies. *J. Immunol. Methods* 1997, **206**, 125-134
- 145 Peeters, J.M., Hazendonk, T.G., Beuvery, E.C. and Tesser, G.I. Comparison of four bifunctional reagents for coupling peptides to proteins and the effect of the three moieties on the immunogenicity of the conjugates. *J. Immunol. Meth.* 1989, **120**, 133-143
- 146 Shuler, K.R., Dunham, R.G. and Kanda, P. A simplified method for determination of peptide-protein molar ratios using amino acid analysis. *J. Immunol. Meth.* 1992, **156**, 137-149
- 147 McLean, G.W., Owsianka, A.M., Subak-Sharpe, J.H. and Marsden, H.S. Generation of anti-peptide and anti-protein sera. Effect of peptide presentation on immunogenicity. *Journal of Immunological Methods* 1991, **137**, 149-157
- 148 Sunthornnandh, P., Matangkasombut, P. and Ratanabanangkoon, K. Preparation, characterization and immunogenicity of various polymers and conjugates of elapid postsynaptic neurotoxins. *Mol. Immunol.* 1992, **29**, 501-510
- 149 Jones, G.L., Spencer, L., Lord, R. and Saul, A.J. Effect of context and adjuvant on the immunogenicity of recombinant proteins and peptide conjugates derived from the polymorphic malarial surface antigen MSA2. *Vaccine* 1996, **14**, 77-84
- 150 Morelli, G., del Valle, J., Lammel, C.J., Pohlner, J., Müller, K., Blake, M., *et al.* Immunogenicity and evolutionary variability of epitopes within IgA1 protease from serogroup A *Neisseria meningitidis*. *Mol. Microbiol.* 1994, **11**, 175-187
- 151 Achtman, M., Neibert, M., Crowe, B.A., Strittmatter, W., Kusecek, B., Weyse, E., *et al.* Purification and characterization of eight class 5 outer membrane protein variants from a clone of *Neisseria meningitidis* serogroup A. *J. Exp. Med.* 1988, **168**, 507-525
- 152 Laemmli, U.K. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970, **227**, 680-685
- 153 Anonymous. Introduction to cleavage techniques. Applied Biosystems, 1990
- 154 Rapp, W. RAPP Polymere, Eugenstr. 38/1, 7400 Tübingen, Germany (personal communication)
- 155 Rose, K., Zeng, W.G., Brown, L.E. and Jackson, D.C. A synthetic peptide-based polyoxime vaccine construct of high purity and activity. *Mol. Immunol.* 1995, **32**, 1031-1037

- 156 Pochon, S., Buchegger, F., Pelegrin, A., Mach, J.P., Offord, R.E., Ryser, J.E., et al. A novel derivative of the chelon desferrioxamine for site-specific conjugation to antibodies. *International Journal of Cancer* 1989, **43**, 1188-1194
- 157 Mutter, M., Tuchscherer, G.G., Miller, C., Altmann, K.H., Carey, Wyss, D.F., et al. Template assembled synthetic proteins with four helix-bundle topology total chemical synthesis and conformational studies. *Journal of the American Chemical Society* 1992, **114**, 1463-1470
- 158 Vilaseca, L.A., Rose, K., Werlen, R., Meunier, A., Offord, R.E., Nichols, C.L., et al. Protein conjugates of defined structure: Synthesis and use of a new carrier molecule. *Bioconjugate Chemistry* 1993, **4**, 515-520
- 159 Liu, F.T., Zinnecker, M., Hamaoka, T. and Katz, D.H. New procedures for preparation and isolation of conjugates of proteins and a synthetic copolymer of D-amino acids and immunochemical characterization of such conjugates. *Biochemistry* 1979, **18**, 690-693
- 160 Kitagawa, T. and Aikawa, T. Enzyme coupled immunoassay of insulin using a novel coupling reagent. *Journal of Biochemistry* 1976, **79**, 233-236
- 161 Chou, P.Y. and Fasman, G.D. Prediction of the secondary structure of proteins from their amino acid sequence. *Advances in Enzymology & Related Areas of Molecular Biology* 1978, **47**, 45-148
- 162 Garnier, J. Analysis of the accuracy and implications of simple methods for predicting the secondary structure of globular proteins. *J. Mol. Biol.* 1978, **120**, 97-120
- 163 Provencher, S.W. and Gloeckner, J. Estimation of globular protein secondary structure from circular dichroism. *Biochemistry* 1981, **20**, 33-37
- 164 Manavalan, P. and Johnson, W.C., Jr. Variable selection method improves the prediction of protein secondary structure from circular dichroism spectra. *Anal. Biochem.* 1987, **167**, 76-85
- 165 Meltzer, N.M., Tous, G.I., Gruber, S. and Stein, S. Gas-phase hydrolysis of proteins and peptides. *Analytical Biochemistry* 1987, **160**, 356-361
- 166 Yokote, Y., Arai, K.M. and Akahane, K. Recovery of tryptophan from 25-minute acid hydrolysates of protein. *Analytical Biochemistry* 1986, **152**, 245-249
- 167 Cochran, W.G. and Snedecor, G.W. Statistical Methods. Iowa State University Press, Ames, 1968
- 168 Blake, M.S. and Eastby, C. Studies on the gonococcal IgA1 protease II. Improved methods of enzyme purification and production of monoclonal antibodies to the enzyme. *J. Immunol. Meth.* 1991, **144**, 215-221
- 169 Olyhoek, T., Crowe, B.A. and Achtman, M. Clonal population structure of *Neisseria meningitidis* serogroup A isolated from epidemics and pandemics between 1915 and 1983. *Rev. Infect. Dis.* 1987, **9**, 665-692
- 170 Schaffner, W. and Weissmann, C. A rapid, sensitive and specific method for the determination of protein in dilute solutions. *Anal. Biochem.* 1973, **56**, 502-514

- 171 Thiesen, B., Greenwood, B., Brieske, N. and Achtman, M. Persistence of antibodies to meningococcal IgA1 protease versus decay of antibodies to group A polysaccharide and Opc protein. *Vaccine* 1997, **15**, 209-219
- 172 Virji, M., Makepeace, K., Ferguson, D.J.P., Achtman, M., Sarkari, J. and Moxon, E.R. Expression of the Opc protein correlates with invasion of epithelial and endothelial cells by *Neisseria meningitidis*. *Mol. Microbiol.* 1992, **6**, 2785-2795
- 173 Virji, M., Makepeace, K., Ferguson, D.J.P., Achtman, M. and Moxon, E.R. Meningococcal Opa and Opc proteins: their role in colonization and invasion of human epithelial and endothelial cells. *Mol. Microbiol.* 1993, **10**, 499-510
- 174 Rosenqvist, E., Høiby, E.A., Wedege, E., Kusecek, B. and Achtman, M. The 5C protein of *Neisseria meningitidis* is highly immunogenic in humans and induces bactericidal antibodies. *J. Infect. Dis.* 1993, **167**, 1065-1073
- 175 Merker, P., Tommassen, J., Virji, M., Sesardic, D. and Achtman, M. Two-dimensional structure of the Opc invasin from *Neisseria meningitidis*. *Mol. Microbiol.* 1997, **23**, 281-293
- 176 Seiler, A., Reinhardt, R., Sarkari, J., Caugant, D.A. and Achtman, M. Allelic polymorphism and site-specific recombination in the *opc* locus of *Neisseria meningitidis*. *Mol. Microbiol.* 1996, **19**, 841-856
- 177 Plaut, A.G. Microbial IgA proteases. *New England J. Med.* 1978, **298**, 1459-1463
- 178 Plaut, A.G. The IgA1 proteases of pathogenic bacteria. *Ann. Rev. Microbiol.* 1983, **37**, 603-622
- 179 Brooks, G.F., Lammel, C.J., Blake, M.S., Kusecek, B. and Achtman, M. Antibodies against IgA1 protease are stimulated both by clinical disease and by asymptomatic carriage of serogroup A *Neisseria meningitidis*. *J. Infect. Dis.* 1992, **166**, 1316-1321
- 180 del Valle, J. Epitopes for murine and human antibodies within the IgA1 protease from *Neisseria meningitidis*. 1995, 1-156
- 181 Morelli, G., Müller, K., Kusecek, B., et al. Epitopes to IgA1 protease of serogroup A meningococci. In: *Pathobiology and immunobiology of Neisseriaceae*. INSP, Cuernavaca, Mexico, 1994, pp. 543-548
- 182 Achtman, M. Max-Planck-Institut für molekulare Gentik, Ihnestr. 73, 14195 Berlin, Germany (personal communication)
- 183 Tam, J.P. Recent advances in multiple antigen peptides. *J. Immunol. Meth.* 1996, **196**, 17-32
- 184 Mutter, M. Nature's rules and chemist's tools: a way for creating novel proteins. *TIBS* 1988, **13**, 260-265
- 185 Tuchscherer, G., Servis, C., Corradin, G., Blum, U., Rivier, J. and Mutter, M. Total chemical synthesis, characterization, and immunological properties of an MHC class I model using the TASP concept for protein de novo design. *Protein Science* 1992, **1**, 1377-1386

- 186 Guiyoule, A., Rasoamanana, B., Buchrieser, C., Michel, P., Chanteau, S. and Carniel, E. Recent emergence of new variants of *Yersinia pestis* in Madagascar. *J. Clin. Microbiol.* 1997, **35**, 2826-2833
- 187 Stahl-Zeng, J. Perceptive Biosystems, Otto-von-Guericke-Ring 7, 65205 Wiesbaden, Germany (personal communication)
- 188 Waterhous, D.V. and Johnson, W.C., Jr. Importance of environment in determining secondary structure in proteins. *Biochemistry* 1994, **33**, 2121-2128
- 189 Busch, R., Reich, Z., Zaller, D.M., Sloan, V. and Mellins, E.D. Secondary structure composition and pH-dependent conformational changes of soluble recombinant HLA-DM. *Journal of Biological Chemistry* 1998, **273**, 27557-27564
- 190 Wang, L.J., Voloshin, O.N., Stasiak, A., Stasiak and Cameriniotero, R.D. Homologous DNA pairing domain peptides of RecA protein: Intrinsic propensity to form β -structures and filaments. *Journal of Molecular Biology* 1998, **277**, 1-11
- 191 Davies, S.A., Kelly, S.M., Price, N.C. and Bradshaw, J.P. Structural plasticity of the feline leukaemia virus fusion peptide: a circular dichroism study. *FEBS Letters* 1998, **425**, 415-418
- 192 Chanh, T.C., Kennedy, R.C. and Kanda, P. Synthetic peptides homologous to HIV transmembrane glycoprotein suppress normal human lymphocyte blastogenic response. *Cellular Immunology* 1988, **111**, 77-86
- 193 Carlsson, J., Drevin, H. and Axen, R. Protein thiolation and reversible protein-protein conjugation. N-Succinimidyl 3-(2-pyridyldithio)propionate, a new heterobifunctional reagent. *Biochemical Journal* 1978, **173**, 723-737
- 194 Lanford, R.E., White, R.G., Dunham, R.G. and Kanda, P. Effect of basic and nonbasic amino acid substitutions on transport induced by simian virus 40 T-antigen synthetic peptide nuclear transport signals. *Molecular & Cellular Biology* 1988, **8**, 2722-2729
- 195 Atassi, M.Z. and Saplin, B.J. Immunochemistry of sperm whale myoglobin. I. The specific interaction of some tryptic peptides and of peptides containing all the reactive regions of the antigen. *Biochemistry* 1968, **7**, 688-698
- 196 Briand, J.P., Muller, S. and Van Regenmortel, M.H.V. Synthetic peptides as antigens: Pitfalls of conjugation methods. *J. Immunol. Meth.* 1985, **78**, 59-69
- 197 Antoni, G. and Presentini, R. A least-squares computer method for the determination of the molecular ratio of conjugates between two different proteins from the results of the amino acid analysis. *Analytical Biochemistry* 1989, **179**, 158-161
- 198 Sedgwick, J.D. and Holt, P.G. A solid-phase immunoenzymatic technique for the enumeration of specific antibody-secreting cells. *J. Immunol. Meth.* 1983, **57**, 301-309
- 199 Czerniksky, C.C., Nilsson, L.A., Nygren, H., Ouchterlony, O. and Tarkowski, A. A solid-phase enzyme-linked immunospot (ELISPOT) assay for enumeration of specific antibody-secreting cells. *J. Immunol. Meth.* 1983, **65**, 109-121

- 200 Jerne, N.K., Henry, C., Nordin, A.A., Fuji, H., Koros, A.C. and Lefkovits, I. Plaque forming cells: methodology and theory. *Transplantation Reviews* 1974, **18**, 130-191
- 201 Gessner, A., Moskophidis, D. and Lehmann Grube, F. Enumeration of single IFN-gamma-producing cells in mice during viral and bacterial infection. *J. Immunol.* 1989, **142**, 1293-1298
- 202 Moskophidis, D. and Lehmann-Grube, F. The immune response of the mouse to lymphocytic choriomeningitis virus. IV. Enumeration of antibody-producing cells in spleens during acute and persistent infection. *J. Immunol.* 1984, **133**, 3366-3370
- 203 Zigterman, G.J., Verheul, A.F., Ernste, E.B., Rombouts, R.F., De Reuver, M.J., Jansze, M., et al. Measurement of the humoral immune response against Streptococcus pneumoniae type 3 capsular polysaccharide and oligosaccharide containing antigens by ELISA and ELISPOT techniques. *J. Immunol. Meth.* 1988, **106**, 101-107
- 204 Hartman, A.B., Van De Verg, L.L., Collins, H., Jr., Tang, D.B., Bendiuk, N.O., Taylor, D.N., et al. Local immune response and protection in the guinea pig keratoconjunctivitis model following immunization with Shigella vaccines. *Infection & Immunity* 1994, **62**, 412-420
- 205 Haq, J.A. and Szewczuk, M.R. Differential effect of aging on B-cell immune responses to cholera toxin in the inductive and effector sites of the mucosal immune system. *Infect. Immun.* 1991, **59**, 3094-3100
- 206 Fujihashi, K., Mcghee, J.R., Beagley, K.W., Mcpherson, D.T., Mcpherson, S.A., Huang, C.M., et al. Cytokine-specific ELISPOT assay. Single cell analysis of IL-2, IL-4 and IL-6 producing cells. *J. Immunol. Meth.* 1993, **160**, 181-189
- 207 Jackson, R.J., Fujihashi, K., Xu Amano, J., Kiyono, H., Elson, C.O. and McGhee, J.R. Optimizing oral vaccine induction of systemic and mucosal B-cell and antibody responses to tetanus toxoid by use of cholera toxin as an adjuvant. *Infection & Immunity* 1993, **61**, 4272-4279
- 208 Sedgwick, J.D. and Czerninsky, C. Detection of cell-surface molecules, secreted products of single cells and cellular proliferation by enzyme immunoassay. *J. Immunol. Meth.* 1992, **150**, 159-175
- 209 Larsson, M., Jin, X., Ramratnam, B., Ogg, G.S., Engelmayer, J., Demoitie, M.A., et al. A recombinant vaccinia virus based ELISPOT assay detects high frequencies of Pol-specific CD8 T cells in HIV-1-positive individuals. *AIDS* 1999, **13**, 767-777
- 210 Czerninsky, C., Moldoveanu, Z., Mestecky, J., Nilsson, L.A. and Ouchterlony, O. A novel two colour ELISPOT assay. I. Simultaneous detection of distinct types of antibody-secreting cells. *Journal of Immunological Methods* 1988, **115**, 31-37
- 211 Lakew, M., Nordstrom, I., Czerninsky, C. and Quiding-Jarbrink, M. Combined immunomagnetic cell sorting and ELISPOT assay for the phenotypic characterization of specific antibody-forming cells. *Journal of Immunological Methods* 1997, **203**, 193-198

- 212 Shirai, A., Holmes, K. and Klinman, D. Detection and quantitation of cells secreting IL-6 under physiologic conditions in BALB/c mice. *Journal of Immunology* 1993, **150**, 793-799
- 213 Slifka, M.K., Matloubian, M. and Ahmed, R. Bone marrow is a major site of long-term antibody production after acute viral infection. *Journal of Virology* 1995, **69**, 1895-1902
- 214 Autenrieth, I. Max von Pettenkofer-Institut für Hygiene und Medizinische Mikrobiologie, LMU München, Germany (personal communication)
- 215 Vadolas J, Davies JK, Wright PJ, Strugnell RA. Intranasal immunisation with liposomes induces strong mucosal immune responses. *Vaccine*. 1995;in press.
- 216 Vijayakrishnan, L., Sarkar, S., Roy, R.P. and Rao, K.V. B cell responses to a peptide epitope: IV. Subtle sequence changes in flanking residues modulate immunogenicity. *Journal of Immunology* 1997, **159**, 1809-1819
- 217 Neurath, A.R., Kent, S.B. and Strick, N. Antibody response to two synthetic peptides corresponding to residues 45-68 and 69-79 of the major protein of hepatitis B surface antigen. *Virus Research* 1984, **1**, 321-331
- 218 Vordermeier, H.-M., Harris, D.P., Moreno, C., Singh, M. and Ivanyi, J. The nature of the immunogen determines the specificity of antibodies and T cells to selected peptides of the 38 kDa mycobacterial antigen. *Int. Immunol.* 1995, **7**, 559-566
- 219 Bhatnagar, P.K., Papas, E., Blum, H.E., Milich, D.R., Nitecki, D., Karels, M.J., et al. Immune response to synthetic peptide analogues of hepatitis B surface antigen specific for the a determinant. *Proceedings of the National Academy of Sciences of the United States of America* 1982, **79**, 4400-4404
- 220 Wychowski, C., van der Werf, S., Siffert, O., Crainic, R., Bruneau, P. and Girard, M. A poliovirus type 1 neutralization epitope is located within amino acid residues 93 to 104 of viral capsid polypeptide VP1. *EMBO Journal* 1983, **2**, 2019-2024
- 221 Kennedy, R.C., Dreesman, G.R., Sparrow, J.T., Culwell, A.R., Sanchez, Ionescu-Matiu, I., et al. Inhibition of a common human anti-hepatitis B surface antigen idiotype by a cyclic synthetic peptide. *Journal of Virology* 1983, **46**, 653-655
- 222 Ionescu-Matiu, I., Kennedy, R.C., Sparrow, J.T., Culwell, A.R., Sanchez, Y., Melnick, J.L., et al. Epitopes associated with a synthetic hepatitis B surface antigen peptide. *Journal of Immunology* 1983, **130**, 1947-1952
- 223 Neurath, A.R., Kent, S.B. and Strick, N. Specificity of antibodies elicited by a synthetic peptide having a sequence in common with a fragment of a virus protein, the hepatitis B surface antigen. *Proceedings of the National Academy of Sciences of the United States of America* 1982, **79**, 7871-7875
- 224 De Franco, M., Gille-Perramant, M.F., Mevel, J.C. and Couderc, J. T helper subset involvement in two high antibody responder lines of mice (Biozzi mice): HI (susceptible) and HII (resistant) to collagen-induced arthritis. *European Journal of Immunology* 1995, **25**, 132-136

- 225 Ahlers, J.D., Dunlop, N., Alling, D.W., Nara, P.L. and Berzofsky, J.A. Cytokine-in-adjuvant steering of the immune response phenotype to HIV-1 vaccine constructs: granulocyte-macrophage colony-stimulating factor and TNF-alpha synergize with IL-12 to enhance induction of cytotoxic T lymphocytes. *Journal of Immunology* 1997, **158**, 3947-3958
- 226 Nishimura, T., Santa, K., Yahata, T., Sato, N., Ohta, A., Ohmi, Y., et al. Involvement of IL-4-producing Vbeta8.2+ CD4+ CD62L- CD45RB- T cells in non-MHC gene-controlled predisposition toward skewing into T helper type-2 immunity in BALB/c mice. *Journal of Immunology* 1997, **158**, 5698-5706
- 227 Muller, C.P., Theisen, D., Damien, B., Bouche, F., von der Ahe, I. and Ammerlaan, W. Problems in designing polyepitope constructs containing T-cell epitopes (TCE) and linear and conformational B-cell epitopes (BCE). International symposium on ADVANCES IN PEPTIDE, PROTEIN & NUCLEIC ACID VACCINE STRATEGIES 1998, Oxford, UK. 1999, in press
- 228 Klaus, G.G. and Cross, A.M. The influence of epitope density on the immunological properties of hapten-protein conjugates. I. Characteristics of the immune response to hapten-coupled albumen with varying epitope density. *Cellular Immunology* 1974, **14**, 226-241
- 229 Desaymard, C. and Howard, J.G. Role of epitope density in the induction of immunity and tolerance with thymus-independent antigens. II. Studies with 2,4-dinitrophenyl conjugates in vivo. *European Journal of Immunology* 1975, **5**, 541-545
- 230 Muller, S., Plaue, S., Couppez, M. and Van Regenmortel, M.H. Comparison of different methods for localizing antigenic regions in histone H2A. *Molecular Immunology* 1986, **23**, 593-601
- 231 Dayhoff, M.O. The origin and evolution of protein superfamilies. *Federation Proceedings* 1976, **35**, 2132-2138
- 232 Nisbet, A.D., Saundry, R.H., Moir, A.J., Fothergill, L.A. and Fothergill, J.E. The complete amino-acid sequence of hen ovalbumin. *European Journal of Biochemistry* 1981, **115**, 335-345
- 233 Bizzini, B., Blass, J., Turpin, A. and Raynaud, M. Chemical characterization of tetanus toxin and toxoid. Amino acid composition, number of SH and S-S groups and N-terminal amino acid. *European Journal of Biochemistry* 1970, **17**, 100-105
- 234 Amkraut, A.A., Malley, A. and Begley, D. Immunogenicity of hemocyanins and their subunits. *Journal of Immunology* 1969, **103**, 1301-1310
- 235 Vassart, G., Bacolla, A., Brocas, H., Christophe, D., de Martynoff, G., Leriche, A., et al. Structure, expression and regulation of the thyroglobulin gene. *Molecular & Cellular Endocrinology* 1985, **40**, 89-97
- 236 Briand, J.P., Andre, C., Tuailon, N., Herve, L., Neimark, J. and Muller, S. Multiple autoepitope presentation for specific detection of antibodies in primary biliary cirrhosis. *Hepatology* 1992, **16**, 1395-1403
- 237 Sarnesto, A., Ranta, S., Seppälä, I.J.T. and Mäkelä, O. Relative immunogenicity in mice of different regions of the human IgG. *Scand. J. Immunol.* 1983, **17**, 507-511

- 238 Van Ommen, R., Vredendaal, A.E. and Savelkoul, H.F. Secondary IgE responses in vivo are predominantly generated via gamma 1 epsilon-double positive B cells. *Scandinavian Journal of Immunology* 1994, **40**, 491-501
- 239 Hennessey, J.P., Jr. and Johnson, W.C., Jr. Information content in the circular dichroism of proteins. *Biochemistry* 1981, **20**, 1085-1094
- 240 Zhong, L. and Johnson, W.C., Jr. Environment affects amino acid preference for secondary structure. *Proc. Natl. Acad. Sci. (U. S. A.)* 1992, **89**, 4462-4465
- 241 Kojima, S., Kuriki, Y., Sato, Y., Arisaka, F., Kumagai, I., Takahashi, S., et al. Synthesis of alpha-helix-forming peptides by gene engineering methods and their characterization by circular dichroism spectra measurements. *Biochimica et Biophysica Acta* 1996, **1294**, 129-137
- 242 Chang, C.T., Wu, C.-S.C. and Yang, J.T. Circular dichroic analysis of protein conformation: Inclusion of the β -turns. *Analytical Biochemistry* 1978, **91**, 13-31
- 243 Dyson, H.J., Merutka, G., Walther, J.P., Lerner, R.A. and Wright, P.E. Folding of peptide fragments comprising the complete sequence of proteins. Models for initiation of protein folding. I. Myohemerythrin. *J. Mol. Biol.* 1992, **226**, 795-817
- 244 Dyson, H.J., Sayre, J.R., Merutka, G., Shin, H.C., Lerner, R.A. and Wright, P.E. Folding of peptide fragments comprising the complete sequence of proteins. Models for initiation of protein folding. II. Plastocyanin. *J. Mol. Biol.* 1993, **226**, 819-835
- 245 Johnson, W.C., Jr. Protein secondary structure and circular dichroism: a practical guide. *Proteins* 1990, **7**, 205-214
- 246 Compton, L.A. and Johnson, W.C., Jr. Analysis of protein circular dichroism spectra for secondary structure using a simple matrix multiplication. *Analytical Biochemistry* 1986, **155**, 155-167
- 247 Ward, L.D., Zhang, J.G., Checkley, G., Preston, B. and Simpson, R.J. Effect of pH and denaturants on the folding and stability of murine interleukin-6. *Prot. Sci.* 1993, **2**, 1291-1300
- 248 Yang, J.T., Wu, C.S. and Martinez, H.M. Calculation of protein conformation from circular dichroism. *Methods in Enzymology* 1986, **130**, 208-269
- 249 Bhattacharjya, S., Awasthi, S.K., Adiga, P.R. and Balaram, P. Folded conformations of antigenic peptides from riboflavin carrier protein in aqueous hexafluoroacetone. *Protein Science* 1998, **7**, 123-131
- 250 Yang, J.J., Buck, M., Pitkeathly, M., Kotik, M., Haynie, D.T., Dobson, C.M., et al. Conformational properties of four peptides spanning the sequence of hen lysozyme. *J. Mol. Biol.* 1995, **252**, 483-491
- 251 Hamada, D., Kuroda, Y., Tanaka, T. and Goto, Y. High helical propensity of the peptide fragments derived from beta-lactoglobulin, a predominantly beta-sheet protein. *J. Mol. Biol.* 1995, **254**, 737-746
- 252 Garnier, J., Osguthorpe, D.J. and Robson, B. Analysis of the accuracy and implications of simple methods for predicting the secondary structure of globular proteins. *Journal of Molecular Biology* 1978, **120**, 97-120

- 253 Yike, I., Ye, J., Zhang, Y., Manavalan, P., Gerken, T.A. and Dearborn, D.G. A recombinant peptide model of the first nucleotide-binding fold of the cystic fibrosis transmembrane conductance regulator: Comparison of wild-type and DELTA-F508 mutant forms. *Prot. Sci.* 1996, **5**, 89-97
- 254 Deroo, S., Elkasmi, K.C., Fournier, P., Theisen, D., Brons, R.C., Herrmann, M., *et al.* Enhanced antigenicity of a four-contact-residue epitope of the measles virus hemagglutinin protein by phage display libraries:evidence of a helical structure in the putative active site. *Molecular Immunology* 1998, **35**, 435-443
- 255 Krstenansky, J.L., Owen, T.J., Hagaman, K.A. and McLean, L.R. Short model peptides having a high alpha-helical tendency: design and solution properties. *FEBS Letters* 1989, **242**, 409-413
- 256 Dyson, H.J., Rance, M., Houghten, R.A., Wright, P.E. and Lerner, R.A. Folding of immunogenic peptide fragments of proteins in water solution. II. The nascent helix. *J. Mol. Biol.* 1993, **201**, 201-217
- 257 Zvi, A., Hiller, R. and Anglister, J. Solution conformation of a peptide corresponding to the principal neutralizing determinant of HIV-1IIIB: a two-dimensional NMR study. *Biochemistry* 1992, **31**, 6972-6979
- 258 Gras-Masse, H., Jolivet, M., Drobecq, H., Aubert, J.P., Beachey, E.H., Audibert, F., *et al.* Influence of helical organization on immunogenicity and antigenicity of synthetic peptides. *Mol. Immunol.* 1988, **25**, 673-678
- 259 Dyson, H.J., Rance, M., Houghten, R.A., Lerner, R.A. and Wright, P.E. Folding of immunogenic peptide fragments of proteins in water solution. I. Sequence requirements for the formation of a reverse turn. *J. Mol. Biol.* 1988, **201**, 161-200
- 260 Precheur, B., Bossus, M., Gras Masse, H., Quiniou, E., Tartar, A. and Craescu, C.T. NMR and circular dichroic studies of the solution structure of conformationally constrained antigenic peptides. *Eur. J. Biochem.* 1994, **220**, 415-425
- 261 Lamthanh, H., Leonetti, M., Nabedryk, E. and Menez, A. CD and FTIR studies of an immunogenic disulphide cyclized octadapeptide, a fragment of a snake curaremimetic toxin. *Biochim. Biophys. Acta* 1993, **1203**, 191-198
- 262 McInnes, C., Sonnichsen, F.D., Kay, C.M., Hodges, R.S. and Sykes, B.D. NMR solution structure and flexibility of a peptide antigen representing the receptor binding domain of *Pseudomonas aeruginosa*. *Biochemistry* 1993, **32**, 13432-13440
- 263 Chandrasekhar, K., Profy, A.T. and Dyson, H.J. Solution conformational preferences of immunogenic peptides derived from the principal neutralizing determinant of the HIV-1 envelope glycoprotein gp120. *Biochemistry* 1991, **30**, 9187-9194
- 264 Dyson, H.J., Satterthwait, A.C., Lerner, R.A. and Wright, P.E. Conformational preferences of synthetic peptides derived from the immunodominant site of the circumsporozoite protein of *Plasmodium falciparum* by ¹H NMR. *Biochemistry* 1990, **29**, 7828-7837
- 265 Londono, J.A., Gras-Masse, H., Dubeaux, C., Tartar, A. and Druilhe, P. Secondary structure and immunogenicity of hybrid synthetic peptides derived

- from two *Plasmodium falciparum* pre-erythrocytic antigens. *J. Immunol.* 1990, **145**, 1557-1563
- 266 O'Hern, P.A. Immunogenicity of peptides having pre-determined alpha-helical and alpha-alpha fold topologies. *Mol. Immunol.* 1991, **28**, 1047-1053
- 267 Laczko, I., Hollosi, M., Urge, L., Ugen, K.E., Weiner, D.B., Mantsch, H.H., *et al.* Synthesis and conformational studies of N-glycosylated analogues of the HIV-1 principal neutralizing determinant. *Biochemistry* 1992, **31**, 4282-4288
- 268 Dyson, H.J., Cross, K.J., Houghten, R.A., Wilson, I.A., Wright, P.E. and Lerner, R.A. The immunodominant site of a synthetic immunogen has a conformational preference in water for a type-II reverse turn. *Nature* 1985, **318**, 480-483
- 269 Dyson, H.J., Lerner, R.A. and Wright, P.E. The physical basis for induction of protein-reactive antipeptide antibodies. *Annual Review of Biophysics & Biophysical Chemistry* 1988, **17**, 305-324
- 270 Gibbs, A.C., Kondejewski, L.H., Gronwald, W., Nip, A.M., Hodges, R.S., Sykes, B.D., *et al.* Unusual β-sheet periodicity in small cyclic peptides. *Nature Structural Biology* 1998, **5**, 284-288
- 271 Nomizu, M., Utani, A., Beck, K., Otaka, A., Roller, P.P. and Yamada, Y. Mechanism of laminin chain assembly into a triple-stranded coiled-coil structure. *Biochemistry* 1996, **35**, 2885-2893
- 272 Monera, O.D., Zhou, N.E., Kay, C.M. and Hodges, R.S. Comparison of antiparallel and parallel two-stranded alpha-helical coiled-coils. *J. Biol. Chem.* 1993, **268**, 19218-19227
- 273 Lu, Y.L. and Derrick, J. Department of Biomolecular Sciences, UMIST, PO Box 88, Manchester M60 1QD, UK (personal communication)
- 274 Kyte, J. Absorption and emission of light. In: *Structure in protein chemistry*. Garland, 1995, pp. 407-413
- 275 Malorny, B. Max-Planck-Institut für molekulare Gentik, Ihnestr. 73, 14195 Berlin, Germany (personal communication)
- 276 Amit, A.G., Mariuzza, R.A., Phillips, S.E.V. and Poljak, R.J. Three-dimensional structure of an antigen-antibody complex at 2.8 Å resolution. *Science* 1986, **233**, 747-753
- 277 Tulip, W.R., Varghese, J.N., Laver, W.G., Webster, R.G. and Colman, P.M. Refined crystal structure of the influenza virus N9 neuroaminidase-NC41 Fab complex. *J. Mol. Biol.* 1992, 122-148
- 278 Sheriff, S., Silverton, E.W., Padlan, E.A., Cohen, G.H., Smith-Gill, S.J., Finzel, B.C., *et al.* Three-dimensional structure of an antibody-antigen complex. *Proc. Natl. Acad. Sci. (U. S. A.)* 1987, **84**, 8075-8079
- 279 Van Regenmortel, M.H.V. Molecular Dissection of Protein Antigens. Van Regenmortel, M. H. V. (Ed.). CRC Press, Boca Raton, 1992
- 280 Kyte, J. and Doolittle, R.F. A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* 1982, **157**, 105-132

- 281 Fleckenstein, B., Kalbacher, H., Muller, C.P., Stoll, D., Halder, T., Jung, G., *et al.* New ligands binding to the human leukocyte antigen class II molecule DRB1*0101 based on the activity pattern of an undecapeptide library. *European Journal of Biochemistry* 1996, **240**, 71-77
- 282 Hemmer, B., Fleckenstein, B.T., Vergelli, M., Jung, G., McFarland, Martin, R., *et al.* Identification of high potency microbial and self ligands for a human autoreactive class II-restricted T cell clone. *Journal of Experimental Medicine* 1997, **185**, 1651-1659
- 283 Zilber, B., Scherf, T., Levitt, M. and Anglister, J. NMR-derived model for a peptide-antibody complex. *Biochemistry* 1990, **29**, 10032-10041
- 284 Partidos, C.D., Delmas, A. and Steward, M.W. Structural requirements for synthetic immunogens to induce measles virus specific CTL responses. *Molecular Immunology* 1996, **33**, 1223-1229
- 285 Nowak, M.A. and McMichael, A.J. How HIV defeats the immune system. *Scientific American* 1995, **273**, 58-65
- 286 Melnick, J.L. Beginnings of the development of synthetic peptides as viral vaccines. *Annales de L'Institut Pasteur Virology* 1986, **137e**, 499-503
- 287 Langeveld, J.P., Kamstrup, S., Uttenthal, A., Strandbygaard, B., Vela, C., Dalsgaard, K., *et al.* Full protection in mink against mink enteritis virus with new generation canine parvovirus vaccines based on synthetic peptide or recombinant protein. *Vaccine* 1995, **13**, 1033-1037
- 288 Campbell, H. and Carter, H. Rational use of *Haemophilus influenzae* type b vaccine. *Drugs* 1993, **46**, 378-383
- 289 Kurikka, S., Kayhty, H., Saarinen, L., Ronnberg, P.R., Eskola, J. and Makela, P.H. Immunologic priming by one dose of *Haemophilus influenzae* type b conjugate vaccine in infancy. *J. Infect. Dis.* 1995, **172**, 1268-1272
- 290 Frisan, T., Levitsky, V., Polack, A. and Masucci, M.G. Phenotype-dependent differences in proteasome subunit composition and cleavage specificity in B cell lines. *Journal of Immunology* 1998, **160**, 3281-3289
- 291 Mosmann, T.R. and Sad, S. The expanding universe of T-cell subsets: Th1, Th2 and more. *Immunol. Today* 1996, **17**, 138-146
- 292 Gundlach, B.R., Wiesmuller, K.H., Junt, T., Kienle, S., Jung, G. and Walden, P. Specificity and degeneracy of minor histocompatibility antigen-specific MHC-restricted CTL. *Journal of Immunology* 1996, **156**, 3645-3651
- 293 Udaka, K., Wiesmuller, K.H., Kienle, S., Jung, G. and Walden, P. Decrypting the structure of major histocompatibility complex class I-restricted cytotoxic T lymphocyte epitopes with complex peptide libraries. *Journal of Experimental Medicine* 1995, **181**, 2097-2108
- 294 Van der Zee, R., Van Eden, W., Meloen, R.H., Noordzij, A. and Van Embden, J.D.A. Efficient mapping and characterization of a T-cell epitope by the simultaneous synthesis of multiple peptides. *Eur. J. Immunol.* 1989, **19**, 43-47

- 295 Deres, K., Schild, H., Wiesmuller, K.H., Jung, G. and Rammensee, H.G. In vivo priming of virus-specific cytotoxic T lymphocytes with synthetic lipopeptide vaccine. *Nature* 1989, **342**, 561-564
- 296 Borges, E., Wiesmuller, K.H., Jung, G. and Walden, P. Efficacy of synthetic vaccines in the induction of cytotoxic T lymphocytes. Comparison of the costimulating support provided by helper T cells and lipoamino acid. *Journal of Immunological Methods* 1994, **173**, 253-263
- 297 Brodie, S.J., Lewinsohn, D.A., Patterson, B.K., Jiyamapa, D., Krieger, J., Corey, L., *et al.* In vivo migration and function of transferred HIV-1-specific cytotoxic T cells. *Nature Medicine* 1999, **5**, 34-41
- 298 Elliott, S.L., Pye, S., Le, T., Mateo, L., Cox, J., Macdonald, L., *et al.* Peptide based cytotoxic T-cell vaccines; delivery of multiple epitopes, help, memory and problems. *Vaccine* 1999, **17**, 2009-2019
- 299 Jung, G. Eberhard-Karls-Universität, Institut für organische Chemie, Auf der Morgenstelle 18, 72076 Tübingen, Germany (personal communication)