

## 9 References

- **Antar, L. N., Afroz, R., Dichtenberg, J. B., Carroll, R. C., and Bassell, G. J.** (2004). Metabotropic glutamate receptor activation regulates fragile x mental retardation protein and FMR1 mRNA localization differentially in dendrites and at synapses. *J Neurosci* **24**, 2648-2655.
- **Avela, K., Lipsanen-Nyman, M., Idanheimo, N., Seemanova, E., Rosengren, S., Makela, T. P., Perheentupa, J., Chapelle, A. D., and Lehesjoki, A. E.** (2000). Gene encoding a new RING-B-box-Coiled-coil protein is mutated in mulibrey nanism. *Nat Genet* **25**, 298-301.
- **Avila, J., Dominguez, J., and Diaz-Nido, J.** (1994). Regulation of microtubule dynamics by microtubule-associated protein expression and phosphorylation during neuronal development. *Int J Dev Biol* **38**, 13-25.
- **Badano, J. L., Teslovich, T. M., and Katsanis, N.** (2005). The centrosome in human genetic disease. *Nat Rev Genet* **6**, 194-205.
- **Bagni, C., and Greenough, W. T.** (2005). From mRNP trafficking to spine dysmorphogenesis: the roots of fragile X syndrome. *Nat Rev Neurosci* **6**, 376-387.
- **Balch, C., and Dedman, J. R.** (1997). Annexins II and V inhibit cell migration. *Exp Cell Res* **237**, 259-263.
- **Barembaum, M., and Bronner-Fraser, M.** (2005). Early steps in neural crest specification. *Semin Cell Dev Biol* **16**, 642-646.
- **Barrallo-Gimeno, A., and Nieto, M. A.** (2005). The Snail genes as inducers of cell movement and survival: implications in development and cancer. *Development* **132**, 3151-3161.
- **Bashirullah, A., Cooperstock, R. L., and Lipshitz, H. D.** (1998). RNA localization in development. *Annu Rev Biochem* **67**, 335-394.
- **Berti, C., Fontanella, B., Ferrentino, R., and Meroni, G.** (2004). Mig12, a novel Opitz syndrome gene product partner, is expressed in the embryonic ventral midline and co-operates with Mid1 to bundle and stabilize microtubules. *BMC Cell Biol* **5**, 9.
- **Borden, K. L.** (1998). RING fingers and B-boxes: zinc-binding protein-protein interaction domains. *Biochem Cell Biol* **76**, 351-358.
- **Borden, K. L.** (2000). RING domains: master builders of molecular scaffolds? *J Mol Biol* **295**, 1103-1112.
- **Borden, K. L., Boddy, M. N., Lally, J., O'Reilly, N. J., Martin, S., Howe, K., Solomon, E., and Freemont, P. S.** (1995a). The solution structure of the RING finger domain from the acute promyelocytic leukaemia proto-oncoprotein PML. *Embo J* **14**, 1532-1541.
- **Borden, K. L., Lally, J. M., Martin, S. R., O'Reilly, N. J., Etkin, L. D., and Freemont, P. S.** (1995b). Novel topology of a zinc-binding domain from a protein involved in regulating early *Xenopus* development. *Embo J* **14**, 5947-5956.
- **Brendel, C., Rehbein, M., Kreienkamp, H. J., Buck, F., Richter, D., and Kindler, S.** (2004). Characterization of Staufen 1 ribonucleoprotein complexes. *Biochem J* **384**, 239-246.
- **Brown, V., Jin, P., Ceman, S., Darnell, J. C., O'Donnell, W. T., et al.** (2001). Microarray identification of FMRP-associated brain mRNAs and altered mRNA translational profiles in fragile X syndrome. *Cell* **107**, 477-487.
- **Brown, V., Small, K., Lakkis, L., Feng, Y., Gunter, C., Wilkinson, K. D., and Warren, S. T.** (1998). Purified recombinant Fmrp exhibits selective RNA binding as an intrinsic property of the fragile X mental retardation protein. *J Biol Chem* **273**, 15521-15527.
- **Buchner, G., Montini, E., Andolfi, G., Quaderi, N., Cainarca, S., Messali, S., Bassi, M. T., Ballabio, A., Meroni, G., and Franco, B.** (1999). MID2, a homologue of the Opitz

syndrome gene MID1: similarities in subcellular localization and differences in expression during development. *Hum Mol Genet* **8**, 1397-1407.

- **Bukau, B., and Horwich, A. L.** (1998). The Hsp70 and Hsp60 chaperone machines. *Cell* **92**, 351-366.
- **Burd, C. G., Matunis, E. L., and Dreyfuss, G.** (1991). The multiple RNA-binding domains of the mRNA poly(A)-binding protein have different RNA-binding activities. *Mol Cell Biol* **11**, 3419-3424.
- **Burkhard, P., Stetefeld, J., and Strelkov, S. V.** (2001). Coiled coils: a highly versatile protein folding motif. *Trends Cell Biol* **11**, 82-88.
- **Burrows, F., Zhang, H., and Kamal, A.** (2004). Hsp90 activation and cell cycle regulation. *Cell Cycle* **3**, 1530-1536.
- **Cainarca, S., Messali, S., Ballabio, A., and Meroni, G.** (1999). Functional characterization of the Opitz syndrome gene product (midin): evidence for homodimerization and association with microtubules throughout the cell cycle. *Hum Mol Genet* **8**, 1387-1396.
- **Cao, T., Borden, K. L., Freemont, P. S., and Etkin, L. D.** (1997). Involvement of the rfp tripartite motif in protein-protein interactions and subcellular distribution. *J Cell Sci* **110 ( Pt 14)**, 1563-1571.
- **Ceman, S., Brown, V., and Warren, S. T.** (1999). Isolation of an FMRP-associated messenger ribonucleoprotein particle and identification of nucleolin and the fragile X-related proteins as components of the complex. *Mol Cell Biol* **19**, 7925-7932.
- **Chattopadhyay, C., Hawke, D., Kobayashi, R., and Maity, S. N.** (2004). Human p32, interacts with B subunit of the CCAAT-binding factor, CBF/NF-Y, and inhibits CBF-mediated transcription activation in vitro. *Nucleic Acids Res* **32**, 3632-3641.
- **Chen, J., Peterson, R. T., and Schreiber, S. L.** (1998). Alpha 4 associates with protein phosphatases 2A, 4, and 6. *Biochem Biophys Res Commun* **247**, 827-832.
- **Chicurel, M. E., Singer, R. H., Meyer, C. J., and Ingber, D. E.** (1998). Integrin binding and mechanical tension induce movement of mRNA and ribosomes to focal adhesions. *Nature* **392**, 730-733.
- **Cohen, M. M., Jr.** (2002). Malformations of the craniofacial region: evolutionary, embryonic, genetic, and clinical perspectives. *Am J Med Genet* **115**, 245-268.
- **Compagni, A., Logan, M., Klein, R., and Adams, R. H.** (2003). Control of skeletal patterning by ephrinB1-EphB interactions. *Dev Cell* **5**, 217-230.
- **Condeelis, J.** (1995). Elongation factor 1 alpha, translation and the cytoskeleton. *Trends Biochem Sci* **20**, 169-170.
- **Consortium, F.** (1997). A candidate gene for familial Mediterranean fever. *Nat Genet* **17**, 25-31.
- **Cooper, G. M.** (2000). *The Cell, a Molecular Approach*.
- **Cordero, J. F., and Holmes, L. B.** (1978). Phenotypic overlap of the BBB and G syndromes. *Am J Med Genet* **2**, 145-152.
- **Cox, E. A., Bennin, D., Doan, A. T., O'Toole, T., and Huttenlocher, A.** (2003). RACK1 regulates integrin-mediated adhesion, protrusion, and chemotactic cell migration via its Src-binding site. *Mol Biol Cell* **14**, 658-669.
- **Cox, T. C.** (2004). Taking it to the max: the genetic and developmental mechanisms coordinating midfacial morphogenesis and dysmorphology. *Clin Genet* **65**, 163-176.
- **Cox, T. C., Allen, L. R., Cox, L. L., Hopwood, B., Goodwin, B., Haan, E., and Suthers, G. K.** (2000). New mutations in MID1 provide support for loss of function as the cause of X-linked Opitz syndrome. *Hum Mol Genet* **9**, 2553-2562.

- **Craig, E. A., Weissman, J. S., and Horwich, A. L.** (1994). Heat shock proteins and molecular chaperones: mediators of protein conformation and turnover in the cell. *Cell* **78**, 365-372.
- **Czar, M. J., Welsh, M. J., and Pratt, W. B.** (1996). Immunofluorescence localization of the 90-kDa heat-shock protein to cytoskeleton. *Eur J Cell Biol* **70**, 322-330.
- **Darnell, J. C., Jensen, K. B., Jin, P., Brown, V., Warren, S. T., and Darnell, R. B.** (2001). Fragile X mental retardation protein targets G quartet mRNAs important for neuronal function. *Cell* **107**, 489-499.
- **Davy, A., Aubin, J., and Soriano, P.** (2004). Ephrin-B1 forward and reverse signaling are required during mouse development. *Genes Dev* **18**, 572-583.
- **Davy, A., and Soriano, P.** (2005). Ephrin signaling in vivo: look both ways. *Dev Dyn* **232**, 1-10.
- **de Carcer, G.** (2004). Heat shock protein 90 regulates the metaphase-anaphase transition in a polo-like kinase-dependent manner. *Cancer Res* **64**, 5106-5112.
- **de Carcer, G., do Carmo Avides, M., Lallena, M. J., Glover, D. M., and Gonzalez, C.** (2001). Requirement of Hsp90 for centrosomal function reflects its regulation of Polo kinase stability. *Embo J* **20**, 2878-2884.
- **De Falco, F., Cainarca, S., Andolfi, G., Ferrentino, R., Berti, C., et al.** (2003). X-linked Opitz syndrome: novel mutations in the MID1 gene and redefinition of the clinical spectrum. *Am J Med Genet A* **120**, 222-228.
- **de Hoog, C. L., Foster, L. J., and Mann, M.** (2004). RNA and RNA binding proteins participate in early stages of cell spreading through spreading initiation centers. *Cell* **117**, 649-662.
- **Dennis, P. B., Fumagalli, S., and Thomas, G.** (1999). Target of rapamycin (TOR): balancing the opposing forces of protein synthesis and degradation. *Curr Opin Genet Dev* **9**, 49-54.
- **Di Como, C. J., and Arndt, K. T.** (1996). Nutrients, via the Tor proteins, stimulate the association of Tap42 with type 2A phosphatases. *Genes Dev* **10**, 1904-1916.
- **Disatnik, M. H., Boutet, S. C., Lee, C. H., Mochly-Rosen, D., and Rando, T. A.** (2002). Sequential activation of individual PKC isozymes in integrin-mediated muscle cell spreading: a role for MARCKS in an integrin signaling pathway. *J Cell Sci* **115**, 2151-2163.
- **Doxsey, S.** (2001). Re-evaluating centrosome function. *Nat Rev Mol Cell Biol* **2**, 688-698.
- **Dreyfuss, G., Kim, V. N., and Kataoka, N.** (2002). Messenger-RNA-binding proteins and the messages they carry. *Nat Rev Mol Cell Biol* **3**, 195-205.
- **Driever, W., and Nusslein-Volhard, C.** (1988). A gradient of bicoid protein in Drosophila embryos. *Cell* **54**, 83-93.
- **Duband, J. L., Monier, F., Delannet, M., and Newgreen, D.** (1995). Epithelium-mesenchyme transition during neural crest development. *Acta Anat (Basel)* **154**, 63-78.
- **Duvel, K., and Broach, J. R.** (2004). The role of phosphatases in TOR signaling in yeast. *Curr Top Microbiol Immunol* **279**, 19-38.
- **El-Hodiri, H. M., Shou, W., and Etkin, L. D.** (1997). xnf7 functions in dorsal-ventral patterning of the Xenopus embryo. *Dev Biol* **190**, 1-17.
- **Everett, A. D., and Brautigan, D. L.** (2002). Developmental expression of alpha4 protein phosphatase regulatory subunit in tissues affected by Opitz syndrome. *Dev Dyn* **224**, 461-464.
- **Fankhauser, C., Izaurralde, E., Adachi, Y., Wingfield, P., and Laemmli, U. K.** (1991). Specific complex of human immunodeficiency virus type 1 rev and nucleolar B23 proteins: dissociation by the Rev response element. *Mol Cell Biol* **11**, 2567-2575.

- **Feng, X., Tonnesen, M. G., Peerschke, E. I., and Ghebrehiwet, B.** (2002). Cooperation of C1q receptors and integrins in C1q-mediated endothelial cell adhesion and spreading. *J Immunol* **168**, 2441-2448.
- **Feng, Y., Absher, D., Eberhart, D. E., Brown, V., Malter, H. E., and Warren, S. T.** (1997). FMRP associates with polyribosomes as an mRNP, and the I304N mutation of severe fragile X syndrome abolishes this association. *Mol Cell* **1**, 109-118.
- **Fields, S., and Sternglanz, R.** (1994). The two hybrid system: an assay for protein-protein interaction. *Trends Genet* **10**, 286-292.
- **Filipenko, N. R., MacLeod, T. J., Yoon, C. S., and Waisman, D. M.** (2004). Annexin A2 is a novel RNA-binding protein. *J Biol Chem* **279**, 8723-8731.
- **Fingar, D. C., and Blenis, J.** (2004). Target of rapamycin (TOR): an integrator of nutrient and growth factor signals and coordinator of cell growth and cell cycle progression. *Oncogene* **23**, 3151-3171.
- **Ford, C. L., Randal-Whitis, L., and Ellis, S. R.** (1999). Yeast proteins related to the p40/laminin receptor precursor are required for 20S ribosomal RNA processing and the maturation of 40S ribosomal subunits. *Cancer Res* **59**, 704-710.
- **Gammill, L. S., and Bronner-Fraser, M.** (2003). Neural crest specification: migrating into genomics. *Nat Rev Neurosci* **4**, 795-805.
- **Gaudenz, K., Roessler, E., Quaderi, N., Franco, B., Feldman, G., et al.** (1998). Opitz G/BBB syndrome in Xp22: mutations in the MID1 gene cluster in the carboxy-terminal domain. *Am J Hum Genet* **63**, 703-710.
- **Gauthier, L. R., and Robbins, S. M.** (2003). Ephrin signaling: One raft to rule them all? One raft to sort them? One raft to spread their call and in signaling bind them? *Life Sci* **74**, 207-216.
- **Ghebrehiwet, B., Feng, X., Kumar, R., and Peerschke, E. I.** (2003). Complement component C1q induces endothelial cell adhesion and spreading through a docking/signaling partnership of C1q receptors and integrins. *Int Immunopharmacol* **3**, 299-310.
- **Gilbert, S. F.** (2003). *Developmental Biology*: (Sinauer associates)).
- **Gingras, A. C., Raught, B., and Sonenberg, N.** (2001). Regulation of translation initiation by FRAP/mTOR. *Genes Dev* **15**, 807-826.
- **Gong, C. X., Wegiel, J., Lidsky, T., Zuck, L., Avila, J., Wisniewski, H. M., Grundke-Iqbal, I., and Iqbal, K.** (2000). Regulation of phosphorylation of neuronal microtubule-associated proteins MAP1b and MAP2 by protein phosphatase-2A and -2B in rat brain. *Brain Res* **853**, 299-309.
- **Granata, A., Savery, D., Hazan, J., Cheung, B. M., Lumsden, A., and Quaderi, N. A.** (2005). Evidence of functional redundancy between MID proteins: implications for the presentation of Opitz syndrome. *Dev Biol* **277**, 417-424.
- **Grignani, F., Fagioli, M., Alcalay, M., Longo, L., Pandolfi, P. P., Donti, E., Biondi, A., Lo Coco, F., and Pelicci, P. G.** (1994). Acute promyelocytic leukemia: from genetics to treatment. *Blood* **83**, 10-25.
- **Grisendi, S., Bernardi, R., Rossi, M., Cheng, K., Khandker, L., Manova, K., and Pandolfi, P. P.** (2005). Role of nucleophosmin in embryonic development and tumorigenesis. *Nature* **437**, 147-153.
- **Gupta, S., and Knowlton, A. A.** (2005). HSP60, Bax, apoptosis and the heart. *J Cell Mol Med* **9**, 51-58.
- **Hall, B. K.** (2000). The neural crest as a fourth germ layer and vertebrates as quadroblastic not triploblastic. *Evol Dev* **2**, 3-5.

- **Hansen, M. D., Ehrlich, J. S., and Nelson, W. J.** (2002). Molecular mechanism for orienting membrane and actin dynamics to nascent cell-cell contacts in epithelial cells. *J Biol Chem* **277**, 45371-45376.
- **Hara, K., Maruki, Y., Long, X., Yoshino, K., Oshiro, N., Hidayat, S., Tokunaga, C., Avruch, J., and Yonezawa, K.** (2002). Raptor, a binding partner of target of rapamycin (TOR), mediates TOR action. *Cell* **110**, 177-189.
- **Hasegawa, N., Iwashita, T., Asai, N., Murakami, H., Iwata, Y., Isomura, T., Goto, H., Hayakawa, T., and Takahashi, M.** (1996). A RING finger motif regulates transforming activity of the rfp/ret fusion gene. *Biochem Biophys Res Commun* **225**, 627-631.
- **Hay, E. D.** (2005). The mesenchymal cell, its role in the embryo, and the remarkable signaling mechanisms that create it. *Dev Dyn* **233**, 706-720.
- **Hay, N., and Sonenberg, N.** (2004). Upstream and downstream of mTOR. *Genes Dev* **18**, 1926-1945.
- **Heitman, J., Movva, N. R., and Hall, M. N.** (1991). Targets for cell cycle arrest by the immunosuppressant rapamycin in yeast. *Science* **253**, 905-909.
- **Helbling, P. M., Tran, C. T., and Brandli, A. W.** (1998). Requirement for EphA receptor signaling in the segregation of *Xenopus* third and fourth arch neural crest cells. *Mech Dev* **78**, 63-79.
- **Helms, J. A., Cordero, D., and Tapadia, M. D.** (2005). New insights into craniofacial morphogenesis. *Development* **132**, 851-861.
- **Henrique, D., and Schweisguth, F.** (2003). Cell polarity: the ups and downs of the Par6/aPKC complex. *Curr Opin Genet Dev* **13**, 341-350.
- **Henry, J., Mather, I. H., McDermott, M. F., and Pontarotti, P.** (1998). B30.2-like domain proteins: update and new insights into a rapidly expanding family of proteins. *Mol Biol Evol* **15**, 1696-1705.
- **Hermanto, U., Zong, C. S., Li, W., and Wang, L. H.** (2002). RACK1, an insulin-like growth factor I (IGF-I) receptor-interacting protein, modulates IGF-I-dependent integrin signaling and promotes cell spreading and contact with extracellular matrix. *Mol Cell Biol* **22**, 2345-2365.
- **Hodges, M., Tissot, C., Howe, K., Grimwade, D., and Freemont, P. S.** (1998). Structure, organization, and dynamics of promyelocytic leukemia protein nuclear bodies. *Am J Hum Genet* **63**, 297-304.
- **Holder, N., and Klein, R.** (1999). Eph receptors and ephrins: effectors of morphogenesis. *Development* **126**, 2033-2044.
- **Huang, X., and Saint-Jeannet, J. P.** (2004). Induction of the neural crest and the opportunities of life on the edge. *Dev Biol* **275**, 1-11.
- **Huynh-Do, U., Stein, E., Lane, A. A., Liu, H., Cerretti, D. P., and Daniel, T. O.** (1999). Surface densities of ephrin-B1 determine EphB1-coupled activation of cell attachment through  $\alpha$ 5 $\beta$ 3 and  $\alpha$ 5 $\beta$ 1 integrins. *Embo J* **18**, 2165-2173.
- **Jacinto, E., and Hall, M. N.** (2003). Tor signalling in bugs, brain and brawn. *Nat Rev Mol Cell Biol* **4**, 117-126.
- **Jacinto, E., Loewith, R., Schmidt, A., Lin, S., Ruegg, M. A., Hall, A., and Hall, M. N.** (2004). Mammalian TOR complex 2 controls the actin cytoskeleton and is rapamycin insensitive. *Nat Cell Biol* **6**, 1122-1128.
- **Jana, N. R., Tanaka, M., Wang, G., and Nukina, N.** (2000). Polyglutamine length-dependent interaction of Hsp40 and Hsp70 family chaperones with truncated N-terminal

huntingtin: their role in suppression of aggregation and cellular toxicity. *Hum Mol Genet* **9**, 2009-2018.

- **Jang, C. Y., Lee, J. Y., and Kim, J.** (2004). Rps3, a DNA repair endonuclease and ribosomal protein, is involved in apoptosis. *FEBS Lett* **560**, 81-85.
- **Jensen, K., Shiels, C., and Freemont, P. S.** (2001). PML protein isoforms and the RBCC/TRIM motif. *Oncogene* **20**, 7223-7233.
- **Jerome, L. A., and Papaioannou, V. E.** (2001). DiGeorge syndrome phenotype in mice mutant for the T-box gene, Tbx1. *Nat Genet* **27**, 286-291.
- **Jiang, Y., and Broach, J. R.** (1999). Tor proteins and protein phosphatase 2A reciprocally regulate Tap42 in controlling cell growth in yeast. *Embo J* **18**, 2782-2792.
- **Jin, P., Alisch, R. S., and Warren, S. T.** (2004a). RNA and microRNAs in fragile X mental retardation. *Nat Cell Biol* **6**, 1048-1053.
- **Jin, P., Zarnescu, D. C., Ceman, S., Nakamoto, M., Mowrey, J., Jongens, T. A., Nelson, D. L., Moses, K., and Warren, S. T.** (2004b). Biochemical and genetic interaction between the fragile X mental retardation protein and the microRNA pathway. *Nat Neurosci* **7**, 113-117.
- **Jones, N. C., and Trainor, P. A.** (2005). Role of morphogens in neural crest cell determination. *J Neurobiol* **64**, 388-404.
- **Kanai, Y., Dohmae, N., and Hirokawa, N.** (2004). Kinesin transports RNA: isolation and characterization of an RNA-transporting granule. *Neuron* **43**, 513-525.
- **Kang, P., and Svoboda, K. K.** (2005). Epithelial-Mesenchymal Transformation during Craniofacial Development. *J Dent Res* **84**, 678-690.
- **Kang, Y., and Massague, J.** (2004). Epithelial-mesenchymal transitions: twist in development and metastasis. *Cell* **118**, 277-279.
- **Kaverina, I., Krylyshkina, O., and Small, J. V.** (1999). Microtubule targeting of substrate contacts promotes their relaxation and dissociation. *J Cell Biol* **146**, 1033-1044.
- **Kaverina, I., Rottner, K., and Small, J. V.** (1998). Targeting, capture, and stabilization of microtubules at early focal adhesions. *J Cell Biol* **142**, 181-190.
- **Kaverina, I. N., Minin, A. A., Gyoeva, F. K., and Vasiliev, J. M.** (1997). Kinesin-associated transport is involved in the regulation of cell adhesion. *Cell Biol Int* **21**, 229-236.
- **Kavi, H. H., Fernandez, H. R., Xie, W., and Birchler, J. A.** (2005). RNA silencing in Drosophila. *FEBS Lett* **579**, 5940-5949.
- **Kaytor, M. D., and Orr, H. T.** (2001). RNA targets of the fragile X protein. *Cell* **107**, 555-557.
- **Kazmin, D. A., Chinenov, Y., Larson, E., and Starkey, J. R.** (2003). Comparative modeling of the N-terminal domain of the 67kDa laminin-binding protein: implications for putative ribosomal function. *Biochem Biophys Res Commun* **300**, 161-166.
- **Keyes, W. M., and Sanders, E. J.** (2002). Regulation of apoptosis in the endocardial cushions of the developing chick heart. *Am J Physiol Cell Physiol* **282**, C1348-1360.
- **Khuri, S., Bakker, F. T., and Dunwell, J. M.** (2001). Phylogeny, function, and evolution of the cupins, a structurally conserved, functionally diverse superfamily of proteins. *Mol Biol Evol* **18**, 593-605.
- **Kiebler, M. A., Hemraj, I., Verkade, P., Kohrmann, M., Fortes, P., Marion, R. M., Ortin, J., and Dotti, C. G.** (1999). The mammalian stau protein localizes to the somatodendritic domain of cultured hippocampal neurons: implications for its involvement in mRNA transport. *J Neurosci* **19**, 288-297.

- **Kim, D. H., Sarbassov, D. D., Ali, S. M., King, J. E., Latek, R. R., Erdjument-Bromage, H., Tempst, P., and Sabatini, D. M.** (2002). mTOR interacts with raptor to form a nutrient-sensitive complex that signals to the cell growth machinery. *Cell* **110**, 163-175.
- **Kim, D. H., Sarbassov, D. D., Ali, S. M., Latek, R. R., Guntur, K. V., Erdjument-Bromage, H., Tempst, P., and Sabatini, D. M.** (2003). GbetaL, a positive regulator of the rapamycin-sensitive pathway required for the nutrient-sensitive interaction between raptor and mTOR. *Mol Cell* **11**, 895-904.
- **Kim, T. S., Jang, C. Y., Kim, H. D., Lee, J. Y., Ahn, B. Y., and Kim, J.** (2006). Interaction of Hsp90 with Ribosomal Proteins Protects from Ubiquitination and Proteasome-dependent Degradation. *Mol Biol Cell* **17**, 824-833.
- **Kimble, M., Khodjakov, A. L., and Kuriyama, R.** (1992). Identification of ubiquitous high-molecular-mass, heat-stable microtubule-associated proteins (MAPs) that are related to the Drosophila 205-kDa MAP but are not related to the mammalian MAP-4. *Proc Natl Acad Sci U S A* **89**, 7693-7697.
- **Kirby, M. L., Gale, T. F., and Stewart, D. E.** (1983). Neural crest cells contribute to normal aorticopulmonary septation. *Science* **220**, 1059-1061.
- **Kloc, M., and Etkin, L. D.** (2005). RNA localization mechanisms in oocytes. *J Cell Sci* **118**, 269-282.
- **Klugbauer, S., and Rabes, H. M.** (1999). The transcription coactivator HTIF1 and a related protein are fused to the RET receptor tyrosine kinase in childhood papillary thyroid carcinomas. *Oncogene* **18**, 4388-4393.
- **Knecht, A. K., and Bronner-Fraser, M.** (2002). Induction of the neural crest: a multigene process. *Nat Rev Genet* **3**, 453-461.
- **Kole, T. P., Tseng, Y., Jiang, I., Katz, J. L., and Wirtz, D.** (2005). Intracellular mechanics of migrating fibroblasts. *Mol Biol Cell* **16**, 328-338.
- **Kontges, G., and Lumsden, A.** (1996). Rhombencephalic neural crest segmentation is preserved throughout craniofacial ontogeny. *Development* **122**, 3229-3242.
- **Krainer, A. R., Mayeda, A., Kozak, D., and Binns, G.** (1991). Functional expression of cloned human splicing factor SF2: homology to RNA-binding proteins, U1 70K, and Drosophila splicing regulators. *Cell* **66**, 383-394.
- **Krumlauf, R.** (1993). Hox genes and pattern formation in the branchial region of the vertebrate head. *Trends Genet* **9**, 106-112.
- **Krylyshkina, O., Anderson, K. I., Kaverina, I., Upmann, I., Manstein, D. J., Small, J. V., and Toomre, D. K.** (2003). Nanometer targeting of microtubules to focal adhesions. *J Cell Biol* **161**, 853-859.
- **Krylyshkina, O., Kaverina, I., Kranewitter, W., Steffen, W., Alonso, M. C., Cross, R. A., and Small, J. V.** (2002). Modulation of substrate adhesion dynamics via microtubule targeting requires kinesin-1. *J Cell Biol* **156**, 349-359.
- **Kullander, K., and Klein, R.** (2002). Mechanisms and functions of Eph and ephrin signalling. *Nat Rev Mol Cell Biol* **3**, 475-486.
- **Landry, J. R., and Mager, D. L.** (2002). Widely spaced alternative promoters, conserved between human and rodent, control expression of the Opitz syndrome gene MID1. *Genomics* **80**, 499-508.
- **Lange, B. M., Bachi, A., Wilm, M., and Gonzalez, C.** (2000). Hsp90 is a core centrosomal component and is required at different stages of the centrosome cycle in Drosophila and vertebrates. *Embo J* **19**, 1252-1262.
- **Lasko, P.** (1999). RNA sorting in Drosophila oocytes and embryos. *Faseb J* **13**, 421-433.

- **Le Douarin, B., Zechel, C., Garnier, J. M., Lutz, Y., Tora, L., Pierrat, P., Heery, D., Gronemeyer, H., Chambon, P., and Losson, R.** (1995). The N-terminal part of TIF1, a putative mediator of the ligand-dependent activation function (AF-2) of nuclear receptors, is fused to B-raf in the oncogenic protein T18. *Embo J* **14**, 2020-2033.
- **Le Douarin, N. M., Creuzet, S., Couly, G., and Dupin, E.** (2004). Neural crest cell plasticity and its limits. *Development* **131**, 4637-4650.
- **Lehmann, R., and Nusslein-Volhard, C.** (1986). Abdominal segmentation, pole cell formation, and embryonic polarity require the localized activity of oskar, a maternal gene in *Drosophila*. *Cell* **47**, 141-152.
- **Li, P., Yang, X., Wasser, M., Cai, Y., and Chia, W.** (1997). Inscuteable and Staufien mediate asymmetric localization and segregation of prospero RNA during *Drosophila* neuroblast cell divisions. *Cell* **90**, 437-447.
- **Liang, P., and MacRae, T. H.** (1997). Molecular chaperones and the cytoskeleton. *J Cell Sci* **110 ( Pt 13)**, 1431-1440.
- **Lim, J., and Lu, K. P.** (2005). Pinning down phosphorylated tau and tauopathies. *Biochim Biophys Acta* **1739**, 311-322.
- **Liu, J., Prickett, T. D., Elliott, E., Meroni, G., and Brautigan, D. L.** (2001). Phosphorylation and microtubule association of the Opitz syndrome protein mid-1 is regulated by protein phosphatase 2A via binding to the regulatory subunit alpha 4. *Proc Natl Acad Sci U S A* **98**, 6650-6655.
- **Locascio, A., Manzanares, M., Blanco, M. J., and Nieto, M. A.** (2002). Modularity and reshuffling of Snail and Slug expression during vertebrate evolution. *Proc Natl Acad Sci U S A* **99**, 16841-16846.
- **Lopez de Silanes, I., Zhan, M., Lal, A., Yang, X., and Gorospe, M.** (2004). Identification of a target RNA motif for RNA-binding protein HuR. *Proc Natl Acad Sci U S A* **101**, 2987-2992.
- **Loreni, F., Iadevaia, V., Tino, E., Caldarola, S., and Amaldi, F.** (2005). RACK1 mRNA translation is regulated via a rapamycin-sensitive pathway and coordinated with ribosomal protein synthesis. *FEBS Lett* **579**, 5517-5520.
- **Lumsden, A., Sprawson, N., and Graham, A.** (1991). Segmental origin and migration of neural crest cells in the hindbrain region of the chick embryo. *Development* **113**, 1281-1291.
- **Lutsch, G., Stahl, J., Kargel, H. J., Noll, F., and Bielka, H.** (1990). Immunoelectron microscopic studies on the location of ribosomal proteins on the surface of the 40S ribosomal subunit from rat liver. *Eur J Cell Biol* **51**, 140-150.
- **Mamidipudi, V., Chang, B. Y., Harte, R. A., Lee, K. C., and Cartwright, C. A.** (2004a). RACK1 inhibits the serum- and anchorage-independent growth of v-Src transformed cells. *FEBS Lett* **567**, 321-326.
- **Mamidipudi, V., Zhang, J., Lee, K. C., and Cartwright, C. A.** (2004b). RACK1 regulates G1/S progression by suppressing Src kinase activity. *Mol Cell Biol* **24**, 6788-6798.
- **Mancilla, A., and Mayor, R.** (1996). Neural crest formation in *Xenopus laevis*: mechanisms of Xslug induction. *Dev Biol* **177**, 580-589.
- **McCahill, A., Warwicker, J., Bolger, G. B., Houslay, M. D., and Yarwood, S. J.** (2002). The RACK1 scaffold protein: a dynamic cog in cell response mechanisms. *Mol Pharmacol* **62**, 1261-1273.
- **Meares, G. P., Zmijewska, A. A., and Jope, R. S.** (2004). Heat shock protein-90 dampens and directs signaling stimulated by insulin-like growth factor-1 and insulin. *FEBS Lett* **574**, 181-186.



- **Mendes, S. W., Henkemeyer, M., and Liebl, D. J.** (2006). Multiple Eph receptors and B-class ephrins regulate midline crossing of corpus callosum fibers in the developing mouse forebrain. *J Neurosci* **26**, 882-892.
- **Meroni, G., and Diez-Roux, G.** (2005). TRIM/RBCC, a novel class of 'single protein RING finger' E3 ubiquitin ligases. *Bioessays* **27**, 1147-1157.
- **Merscher, S., Funke, B., Epstein, J. A., Heyer, J., Puech, A., et al.** (2001). TBX1 is responsible for cardiovascular defects in velo-cardio-facial/DiGeorge syndrome. *Cell* **104**, 619-629.
- **Miller, L. D., Lee, K. C., Mochly-Rosen, D., and Cartwright, C. A.** (2004). RACK1 regulates Src-mediated Sam68 and p190RhoGAP signaling. *Oncogene* **23**, 5682-5686.
- **Mitsiadis, T. A., Cheraud, Y., Sharpe, P., and Fontaine-Perus, J.** (2003). Development of teeth in chick embryos after mouse neural crest transplantations. *Proc Natl Acad Sci U S A* **100**, 6541-6545.
- **Mitsiadis, T. A., Salmivirta, M., Muramatsu, T., Muramatsu, H., Rauvala, H., Lehtonen, E., Jalkanen, M., and Thesleff, I.** (1995). Expression of the heparin-binding cytokines, midkine (MK) and HB-GAM (pleiotrophin) is associated with epithelial-mesenchymal interactions during fetal development and organogenesis. *Development* **121**, 37-51.
- **Mitsui, K., Nakayama, H., Akagi, T., Nekooki, M., Ohtawa, K., Takio, K., Hashikawa, T., and Nukina, N.** (2002). Purification of polyglutamine aggregates and identification of elongation factor-1alpha and heat shock protein 84 as aggregate-interacting proteins. *J Neurosci* **22**, 9267-9277.
- **Mnayer, L., Khuri, S., Al-Ali Merheby, H., Meroni, G., and Elsas, L. J.** (2005). A structure-function study of MID1 mutations associated with a mild Opitz phenotype. *Mol Genet Metab*.
- **Mohr, E., and Richter, D.** (2001). Messenger RNA on the move: implications for cell polarity. *Int J Biochem Cell Biol* **33**, 669-679.
- **Moore R.C, and R.J., C.** (2000). Association between elongation factor-1alpha and microtubules in vivo is domain dependent and conditional. *Cell Motil Cytoskeleton* **45**, 279-292.
- **Moore RC, Durso NA, and RJ, C.** (1998). Elongation factor-1alpha stabilizes microtubules in a calcium/calmodulin-dependent manner. *Cell Motil Cytoskeleton* **41**, 168-180.
- **Moore, R. C., and Cyr, R. J.** (2000). Association between elongation factor-1alpha and microtubules in vivo is domain dependent and conditional. *Cell Motil Cytoskeleton* **45**, 279-292.
- **Moore, R. C., Durso, N. A., and Cyr, R. J.** (1998). Elongation factor-1alpha stabilizes microtubules in a calcium/calmodulin-dependent manner. *Cell Motil Cytoskeleton* **41**, 168-180.
- **Moraes, F., Novoa, A., Jerome-Majewska, L. A., Papaioannou, V. E., and Mallo, M.** (2005). Tbx1 is required for proper neural crest migration and to stabilize spatial patterns during middle and inner ear development. *Mech Dev* **122**, 199-212.
- **Morales, A. V., Barbas, J. A., and Nieto, M. A.** (2005). How to become neural crest: from segregation to delamination. *Semin Cell Dev Biol* **16**, 655-662.
- **Mowat, D. R., Wilson, M. J., and Goossens, M.** (2003). Mowat-Wilson syndrome. *J Med Genet* **40**, 305-310.
- **Murai, K. K., and Pasquale, E. B.** (2004). Eph receptors, ephrins, and synaptic function. *Neuroscientist* **10**, 304-314.
- **Murata, K., Wu, J., and Brautigam, D. L.** (1997). B cell receptor-associated protein alpha4 displays rapamycin-sensitive binding directly to the catalytic subunit of protein phosphatase 2A. *Proc Natl Acad Sci U S A* **94**, 10624-10629.

- **Nakai, A., and Ishikawa, T.** (2001). Cell cycle transition under stress conditions controlled by vertebrate heat shock factors. *Embo J* **20**, 2885-2895.
- **Nanahoshi, M., Tsujishita, Y., Tokunaga, C., Inui, S., Sakaguchi, N., Hara, K., and Yonezawa, K.** (1999). Alpha4 protein as a common regulator of type 2A-related serine/threonine protein phosphatases. *FEBS Lett* **446**, 108-112.
- **Negrutskii, B. S., and El'skaya, A. V.** (1998). Eukaryotic translation elongation factor 1 alpha: structure, expression, functions, and possible role in aminoacyl-tRNA channeling. *Prog Nucleic Acid Res Mol Biol* **60**, 47-78.
- **Nieto, M. A., Sargent, M. G., Wilkinson, D. G., and Cooke, J.** (1994). Control of cell behavior during vertebrate development by Slug, a zinc finger gene. *Science* **264**, 835-839.
- **Nilsson, J., Sengupta, J., Frank, J., and Nissen, P.** (2004). Regulation of eukaryotic translation by the RACK1 protein: a platform for signalling molecules on the ribosome. *EMBO Rep* **5**, 1137-1141.
- **Nisole, S., Stoye, J. P., and Saib, A.** (2005). TRIM family proteins: retroviral restriction and antiviral defence. *Nat Rev Microbiol* **3**, 799-808.
- **Nobes, C. D., and Hall, A.** (1995). Rho, rac, and cdc42 GTPases regulate the assembly of multimolecular focal complexes associated with actin stress fibers, lamellipodia, and filopodia. *Cell* **81**, 53-62.
- **Noden, D. M., and Trainor, P. A.** (2005). Relations and interactions between cranial mesoderm and neural crest populations. *J Anat* **207**, 575-601.
- **Ogawa, K., Wada, H., Okada, N., Harada, I., Nakajima, T., Pasquale, E. B., and Tsuyama, S.** (2006). EphB2 and ephrin-B1 expressed in the adult kidney regulate the cytoarchitecture of medullary tubule cells through Rho family GTPases. *J Cell Sci* **119**, 559-570.
- **Ohta, K., Toriyama, M., Miyazaki, M., Murofushi, H., Hosoda, S., Endo, S., and Sakai, H.** (1990). The mitotic apparatus-associated 51-kDa protein from sea urchin eggs is a GTP-binding protein and is immunologically related to yeast polypeptide elongation factor 1 alpha. *J Biol Chem* **265**, 3240-3247.
- **Okuda, M.** (2002). The role of nucleophosmin in centrosome duplication. *Oncogene* **21**, 6170-6174.
- **Okuwaki, M., Tsujimoto, M., and Nagata, K.** (2002). The RNA binding activity of a ribosome biogenesis factor, nucleophosmin/B23, is modulated by phosphorylation with a cell cycle-dependent kinase and by association with its subtype. *Mol Biol Cell* **13**, 2016-2030.
- **Opitz, J. M., Frias J.L., Guttenberger, J.E, Pellet J.R..** (1969a). The G syndrome of multiple congenital anomalies. *Birth Defects Orig Artic Ser (V)* **2**, 95-102.
- **Opitz, J. M., Summitt, R.L., Smith, D.W.** (1969b). The BBB syndrome. Familial telecanthus with associated congenital anomalies. *Birth Defects Orig Artic Ser (V)* **2**, 86-94.
- **Palazzo, A. F., and Gundersen, G. G.** (2002). Microtubule-actin cross-talk at focal adhesions. *Sci STKE* **2002**, PE31.
- **Palmer, A., and Klein, R.** (2003). Multiple roles of ephrins in morphogenesis, neuronal networking, and brain function. *Genes Dev* **17**, 1429-1450.
- **Pasquale, E. B.** (2005). Eph receptor signalling casts a wide net on cell behaviour. *Nat Rev Mol Cell Biol* **6**, 462-475.
- **Pende, M., Um, S. H., Mieulet, V., Sticker, M., Goss, V. L., Mestan, J., Mueller, M., Fumagalli, S., Kozma, S. C., and Thomas, G.** (2004). S6K1(-/-)/S6K2(-/-) mice exhibit perinatal lethality and rapamycin-sensitive 5'-terminal oligopyrimidine mRNA translation and reveal a mitogen-activated protein kinase-dependent S6 kinase pathway. *Mol Cell Biol* **24**, 3112-3124.

- **Perkins, D. N., Pappin, D. J., Creasy, D. M., and Cottrell, J. S.** (1999). Probability-based protein identification by searching sequence databases using mass spectrometry data. *Electrophoresis* **20**, 3551-3567.
- **Peterson, R. T., Desai, B. N., Hardwick, J. S., and Schreiber, S. L.** (1999). Protein phosphatase 2A interacts with the 70-kDa S6 kinase and is activated by inhibition of FKBP12-rapamycin-associated protein. *Proc Natl Acad Sci U S A* **96**, 4438-4442.
- **Pinson, L., Auge, J., Audollent, S., Mattei, G., Etchevers, H., et al.** (2004). Embryonic expression of the human MID1 gene and its mutations in Opitz syndrome. *J Med Genet* **41**, 381-386.
- **Pla, P., Moore, R., Morali, O. G., Grille, S., Martinuzzi, S., Delmas, V., and Larue, L.** (2001). Cadherins in neural crest cell development and transformation. *J Cell Physiol* **189**, 121-132.
- **Poliakov, A., Cotrina, M., and Wilkinson, D. G.** (2004). Diverse roles of eph receptors and ephrins in the regulation of cell migration and tissue assembly. *Dev Cell* **7**, 465-480.
- **Prescott, N. J., Winter, R. M., and Malcolm, S.** (2001). Nonsyndromic cleft lip and palate: complex genetics and environmental effects. *Ann Hum Genet* **65**, 505-515.
- **Qi, M., Ikematsu, S., Maeda, N., Ichihara-Tanaka, K., Sakuma, S., Noda, M., Muramatsu, T., and Kadomatsu, K.** (2001). Haptotactic migration induced by midkine. Involvement of protein-tyrosine phosphatase zeta, Mitogen-activated protein kinase, and phosphatidylinositol 3-kinase. *J Biol Chem* **276**, 15868-15875.
- **Quaderi, N. A., Schweiger, S., Gaudenz, K., Franco, B., Rugarli, E. I., et al.** (1997). Opitz G/BBB syndrome, a defect of midline development, is due to mutations in a new RING finger gene on Xp22. *Nat Genet* **17**, 285-291.
- **Ramos, A., Hollingworth, D., and Pastore, A.** (2003). G-quartet-dependent recognition between the FMRP RGG box and RNA. *Rna* **9**, 1198-1207.
- **Raught, B., Gingras, A. C., and Sonenberg, N.** (2001). The target of rapamycin (TOR) proteins. *Proc Natl Acad Sci U S A* **98**, 7037-7044.
- **Reymond, A., Meroni, G., Fantozzi, A., Merla, G., Cairo, S., et al.** (2001). The tripartite motif family identifies cell compartments. *Embo J* **20**, 2140-2151.
- **Richman, J. M., Fu, K. K., Cox, L. L., Sibbons, J. P., and Cox, T. C.** (2002). Isolation and characterisation of the chick orthologue of the Opitz syndrome gene, Mid1, supports a conserved role in vertebrate development. *Int J Dev Biol* **46**, 441-448.
- **Richter, K., and Buchner, J.** (2001). Hsp90: chaperoning signal transduction. *J Cell Physiol* **188**, 281-290.
- **Robin, N. H., Feldman, G. J., Aronson, A. L., Mitchell, H. F., Weksberg, R., et al.** (1995). Opitz syndrome is genetically heterogeneous, with one locus on Xp22, and a second locus on 22q11.2. *Nat Genet* **11**, 459-461.
- **Robin, N. H., Opitz, J. M., and Muenke, M.** (1996). Opitz G/BBB syndrome: clinical comparisons of families linked to Xp22 and 22q, and a review of the literature. *Am J Med Genet* **62**, 305-317.
- **Robinson, V., Smith, A., Flenniken, A. M., and Wilkinson, D. G.** (1997). Roles of Eph receptors and ephrins in neural crest pathfinding. *Cell Tissue Res* **290**, 265-274.
- **Roessler, E., and Muenke, M.** (2001). Midline and laterality defects: left and right meet in the middle. *Bioessays* **23**, 888-900.
- **Roessler, E., and Muenke, M.** (2003). How a Hedgehog might see holoprosencephaly. *Hum Mol Genet* **12 Spec No 1**, R15-25.

- **Runyan, R. B., and Markwald, R. R.** (1983). Invasion of mesenchyme into three-dimensional collagen gels: a regional and temporal analysis of interaction in embryonic heart tissue. *Dev Biol* **95**, 108-114.
- **Santagati, F., and Rijli, F. M.** (2003). Cranial neural crest and the building of the vertebrate head. *Nat Rev Neurosci* **4**, 806-818.
- **Santiago, A., and Erickson, C. A.** (2002). Ephrin-B ligands play a dual role in the control of neural crest cell migration. *Development* **129**, 3621-3632.
- **Sarbassov, D. D., Ali, S. M., Kim, D. H., Guertin, D. A., Latek, R. R., Erdjument-Bromage, H., Tempst, P., and Sabatini, D. M.** (2004). Rictor, a novel binding partner of mTOR, defines a rapamycin-insensitive and raptor-independent pathway that regulates the cytoskeleton. *Curr Biol* **14**, 1296-1302.
- **Saurin, A. J., Borden, K. L., Boddy, M. N., and Freemont, P. S.** (1996). Does this have a familiar RING? *Trends Biochem Sci* **21**, 208-214.
- **Savagner, P.** (2001). Leaving the neighborhood: molecular mechanisms involved during epithelial-mesenchymal transition. *Bioessays* **23**, 912-923.
- **Schutte, B. C., and Murray, J. C.** (1999). The many faces and factors of orofacial clefts. *Hum Mol Genet* **8**, 1853-1859.
- **Schweiger, S., Foerster, J., Lehmann, T., Suckow, V., Muller, Y. A., et al.** (1999). The Opitz syndrome gene product, MID1, associates with microtubules. *Proc Natl Acad Sci U S A* **96**, 2794-2799.
- **Schweiger, S., and Schneider, R.** (2003). The MID1/PP2A complex: a key to the pathogenesis of Opitz BBB/G syndrome. *Bioessays* **25**, 356-366.
- **Sela-Donenfeld, D., and Kalcheim, C.** (1999). Regulation of the onset of neural crest migration by coordinated activity of BMP4 and Noggin in the dorsal neural tube. *Development* **126**, 4749-4762.
- **Selleck, M. A., and Bronner-Fraser, M.** (1995). Origins of the avian neural crest: the role of neural plate-epidermal interactions. *Development* **121**, 525-538.
- **Sengupta, B., Banerjee, A., and Sengupta, P. K.** (2005). Interactions of the plant flavonoid fisetin with macromolecular targets: insights from fluorescence spectroscopic studies. *J Photochem Photobiol B* **80**, 79-86.
- **Shiina, N., Gotoh, Y., Kubomura, N., Iwamatsu, A., and Nishida, E.** (1994). Microtubule severing by elongation factor 1 alpha. *Science* **266**, 282-285.
- **Shinmura, K., Tarapore, P., Tokuyama, Y., George, K. R., and Fukasawa, K.** (2005). Characterization of centrosomal association of nucleophosmin/B23 linked to Crm1 activity. *FEBS Lett* **579**, 6621-6634.
- **Short, K. M., and Cox, T. C.** (2006). Sub-classification of the rbcc/trim superfamily reveals a novel motif necessary for microtubule binding. *J Biol Chem*.
- **Short, K. M., Hopwood, B., Yi, Z., and Cox, T. C.** (2002). MID1 and MID2 homo- and heterodimerise to tether the rapamycin-sensitive PP2A regulatory subunit, alpha 4, to microtubules: implications for the clinical variability of X-linked Opitz GBBB syndrome and other developmental disorders. *BMC Cell Biol* **3**, 1.
- **Sittler, A., Walter, S., Wedemeyer, N., Hasenbank, R., Scherzinger, E., Eickhoff, H., Bates, G. P., Lehrach, H., and Wanker, E. E.** (1998). SH3GL3 associates with the Huntingtin exon 1 protein and promotes the formation of polyglu-containing protein aggregates. *Mol Cell* **2**, 427-436.
- **Sklan, E. H., Podoly, E., and Soreq, H.** (2006). RACK1 has the nerve to act: Structure meets function in the nervous system. *Prog Neurobiol*.

- Small, J. V., and Kaverina, I. (2005). Polarized cell motility: microtubules show the way, In Cell migration, D. Wedlich, ed. (WILEY-VCH verlag), pp. 15-31.
- **Smith, A., Robinson, V., Patel, K., and Wilkinson, D. G.** (1997). The EphA4 and EphB1 receptor tyrosine kinases and ephrin-B2 ligand regulate targeted migration of branchial neural crest cells. *Curr Biol* **7**, 561-570.
- **So, J., Suckow, V., Kijas, Z., Kalscheuer, V., Moser, B., et al.** (2005). Mild phenotypes in a series of patients with Opitz GBBB syndrome with MID1 mutations. *Am J Med Genet A* **132**, 1-7.
- **St Johnston, D.** (2005). Moving messages: the intracellular localization of mRNAs. *Nat Rev Mol Cell Biol* **6**, 363-375.
- **Stahl, J.** (2001). Ribosomal Proteins in Eukaryotes. *encyclopedia of life sciences*, 1-11.
- **Sternsdorf, T., Grotzinger, T., Jensen, K., and Will, H.** (1997). Nuclear dots: actors on many stages. *Immunobiology* **198**, 307-331.
- **Steventon, B., Carmona-Fontaine, C., and Mayor, R.** (2005). Genetic network during neural crest induction: from cell specification to cell survival. *Semin Cell Dev Biol* **16**, 647-654.
- **Steward, O., and Schuman, E. M.** (2003). Compartmentalized synthesis and degradation of proteins in neurons. *Neuron* **40**, 347-359.
- **Steward, O., and Worley, P.** (2001). Localization of mRNAs at synaptic sites on dendrites. *Results Probl Cell Differ* **34**, 1-26.
- **Storz, P., Hausser, A., Link, G., Dedio, J., Ghebrehiwet, B., Pfizenmaier, K., and Johannes, F. J.** (2000). Protein kinase C [micro] is regulated by the multifunctional chaperon protein p32. *J Biol Chem* **275**, 24601-24607.
- **Sung, Y. J., Conti, J., Currie, J. R., Brown, W. T., and Denman, R. B.** (2000). RNAs that interact with the fragile X syndrome RNA binding protein FMRP. *Biochem Biophys Res Commun* **275**, 973-980.
- **Sung, Y. J., Dolzhanskaya, N., Nolin, S. L., Brown, T., Currie, J. R., and Denman, R. B.** (2003). The fragile X mental retardation protein FMRP binds elongation factor 1A mRNA and negatively regulates its translation in vivo. *J Biol Chem* **278**, 15669-15678.
- **Tamanini, F., Meijer, N., Verheij, C., Willems, P. J., Galjaard, H., Oostra, B. A., and Hoogeveen, A. T.** (1996). FMRP is associated to the ribosomes via RNA. *Hum Mol Genet* **5**, 809-813.
- **Tang, H., Hornstein, E., Stolovich, M., Levy, G., Livingstone, M., Templeton, D., Avruch, J., and Meyuhas, O.** (2001a). Amino acid-induced translation of TOP mRNAs is fully dependent on phosphatidylinositol 3-kinase-mediated signaling, is partially inhibited by rapamycin, and is independent of S6K1 and rpS6 phosphorylation. *Mol Cell Biol* **21**, 8671-8683.
- **Tang, S. J., Meulemans, D., Vazquez, L., Colaco, N., and Schuman, E.** (2001b). A role for a rat homolog of stau6 in the transport of RNA to neuronal dendrites. *Neuron* **32**, 463-475.
- **Tapadia, M. D., Cordero, D. R., and Helms, J. A.** (2005). It's all in your head: new insights into craniofacial development and deformation. *J Anat* **207**, 461-477.
- **Tarapore, P., Okuda, M., and Fukasawa, K.** (2002). A mammalian in vitro centriole duplication system: evidence for involvement of CDK2/cyclin E and nucleophosmin/B23 in centrosome duplication. *Cell Cycle* **1**, 75-81.
- **Tarapore, P., Shinmura, K., Suzuki, H., Tokuyama, Y., Kim, S. H., Mayeda, A., and Fukasawa, K.** (2006). Thr199 phosphorylation targets nucleophosmin to nuclear speckles and represses pre-mRNA processing. *FEBS Lett* **580**, 399-409.
- **Thiery, J. P.** (2003). Epithelial-mesenchymal transitions in development and pathologies. *Curr Opin Cell Biol* **15**, 740-746.

- **Thomas, M. G., Martinez Tosar, L. J., Loschi, M., Pasquini, J. M., Correale, J., Kindler, S., and Boccaccio, G. L.** (2005). Staufen recruitment into stress granules does not affect early mRNA transport in oligodendrocytes. *Mol Biol Cell* **16**, 405-420.
- **Thorogood, P.** (1989). Developmental and Evolutionary Aspects of the Neural Crest. *Trends Neuroscience* **12**, 38-39.
- **Tokuyama, Y., Horn, H. F., Kawamura, K., Tarapore, P., and Fukasawa, K.** (2001). Specific phosphorylation of nucleophosmin on Thr(199) by cyclin-dependent kinase 2-cyclin E and its role in centrosome duplication. *J Biol Chem* **276**, 21529-21537.
- **Torok, M., and Etkin, L. D.** (2001). Two B or not two B? Overview of the rapidly expanding B-box family of proteins. *Differentiation* **67**, 63-71.
- **Trainor, P. A., and Krumlauf, R.** (2001). Hox genes, neural crest cells and branchial arch patterning. *Curr Opin Cell Biol* **13**, 698-705.
- **Trockenbacher, A., Suckow, V., Foerster, J., Winter, J., Krauss, S., Ropers, H. H., Schneider, R., and Schweiger, S.** (2001). MID1, mutated in Opitz syndrome, encodes an ubiquitin ligase that targets phosphatase 2A for degradation. *Nat Genet* **29**, 287-294.
- **Trojanowski, J. Q., and Lee, V. M.** (1995). Phosphorylation of paired helical filament tau in Alzheimer's disease neurofibrillary lesions: focusing on phosphatases. *Faseb J* **9**, 1570-1576.
- **Twigg, S. R., Kan, R., Babbs, C., Bochukova, E. G., Robertson, S. P., Wall, S. A., Morriss-Kay, G. M., and Wilkie, A. O.** (2004). Mutations of ephrin-B1 (EFNB1), a marker of tissue boundary formation, cause craniofrontonasal syndrome. *Proc Natl Acad Sci U S A* **101**, 8652-8657.
- **Vallee, R. B.** (1982). A taxol-dependent procedure for the isolation of microtubules and microtubule-associated proteins (MAPs). *J Cell Biol* **92**, 435-442.
- **Vasiliev, J. M., Gelfand, I. M., Domnina, L. V., Ivanova, O. Y., Komm, S. G., and Olshevskaia, L. V.** (1970). Effect of colcemid on the locomotory behaviour of fibroblasts. *J Embryol Exp Morphol* **24**, 625-640.
- **Vedeler, A., and Hollas, H.** (2000). Annexin II is associated with mRNAs which may constitute a distinct subpopulation. *Biochem J* **348 Pt 3**, 565-572.
- **Villace, P., Marion, R. M., and Ortin, J.** (2004). The composition of Staufen-containing RNA granules from human cells indicates their role in the regulated transport and translation of messenger RNAs. *Nucleic Acids Res* **32**, 2411-2420.
- **Vitelli, F., Morishima, M., Taddei, I., Lindsay, E. A., and Baldini, A.** (2002). Tbx1 mutation causes multiple cardiovascular defects and disrupts neural crest and cranial nerve migratory pathways. *Hum Mol Genet* **11**, 915-922.
- **Voos, W., and Rottgers, K.** (2002). Molecular chaperones as essential mediators of mitochondrial biogenesis. *Biochim Biophys Acta* **1592**, 51-62.
- **Waldo, K., Miyagawa-Tomita, S., Kumiski, D., and Kirby, M. L.** (1998). Cardiac neural crest cells provide new insight into septation of the cardiac outflow tract: aortic sac to ventricular septal closure. *Dev Biol* **196**, 129-144.
- **Wang, D., Baumann, A., Szebeni, A., and Olson, M. O.** (1994). The nucleic acid binding activity of nucleolar protein B23.1 resides in its carboxyl-terminal end. *J Biol Chem* **269**, 30994-30998.
- **Wieland, I., Jakubiczka, S., Muschke, P., Cohen, M., Thiele, H., Gerlach, K. L., Adams, R. H., and Wieacker, P.** (2004). Mutations of the ephrin-B1 gene cause craniofrontonasal syndrome. *Am J Hum Genet* **74**, 1209-1215.
- **Wilhelm, J. E., and Smibert, C. A.** (2005). Mechanisms of translational regulation in Drosophila. *Biol Cell* **97**, 235-252.

- 
- **Wilkie, A. O., and Morriss-Kay, G. M.** (2001). Genetics of craniofacial development and malformation. *Nat Rev Genet* **2**, 458-468.
  - **Winter, J.** (2003) Molekulare Charakterisierung des *MIDI*-Gens, thesis, Freien Universität Berlin, Berlin.
  - **Winter, J., Lehmann, T., Krauss, S., Trockenbacher, A., Kijas, Z., et al.** (2004). Regulation of the MID1 protein function is fine-tuned by a complex pattern of alternative splicing. *Hum Genet* **114**, 541-552.
  - **Wittmann-Liebold, B., and Graack, H.-R.** (2001). Ribosomal Proteins: Structure and Evolution. *encyclopedia of life sciences*, 1-10.
  - **Zalfa, F., Adinolfi, S., Napoli, I., Kuhn-Holsken, E., Urlaub, H., Achsel, T., Pastore, A., and Bagni, C.** (2005). Fragile X mental retardation protein (FMRP) binds specifically to the brain cytoplasmic RNAs BC1/BC200 via a novel RNA-binding motif. *J Biol Chem* **280**, 33403-33410.
  - **Zalfa, F., and Bagni, C.** (2005). Another view of the role of FMRP in translational regulation. *Cell Mol Life Sci* **62**, 251-252.
  - **Zalfa, F., Giorgi, M., Primerano, B., Moro, A., Di Penta, A., Reis, S., Oostra, B., and Bagni, C.** (2003). The fragile X syndrome protein FMRP associates with BC1 RNA and regulates the translation of specific mRNAs at synapses. *Cell* **112**, 317-327.
  - **Zhang, H., Shi, X., Paddon, H., Hampong, M., Dai, W., and Pelech, S.** (2004). B23/nucleophosmin serine 4 phosphorylation mediates mitotic functions of polo-like kinase 1. *J Biol Chem* **279**, 35726-35734.
  - **Zhang, W., Zong, C. S., Hermanto, U., Lopez-Bergami, P., Ronai, Z., and Wang, L. H.** (2006). RACK1 recruits STAT3 specifically to insulin and insulin-like growth factor 1 receptors for activation, which is important for regulating anchorage-independent growth. *Mol Cell Biol* **26**, 413-424.
  - **Zhao, W. Q., Chen, G. H., Chen, H., Pascale, A., Ravindranath, L., Quon, M. J., and Alkon, D. L.** (2003). Secretion of Annexin II via activation of insulin receptor and insulin-like growth factor receptor. *J Biol Chem* **278**, 4205-4215.