

8. REFERENCES

- Abdelaziz, A. I. et al. (2004): Functional characterization of the human atrial essential myosin light chain (hALC-1) in a transgenic rat model. *J. Mol. Med.* 82, 265-274
- Abelmann, W. H. and Lorell, B. H. (1989): The challenge of cardiomyopathy. *J. Am. Coll. Cardiol.* 13, 1219-1239
- Ackerman, M. J. et al. (2002): Prevalence and Age-Dependence of Malignant Mutations in the Beta-Myosin Heavy Chain and Troponin T Genes in Hypertrophic Cardiomyopathy. A Comprehensive Outpatient Perspective. *J. Am. Coll. Cardiol.* 39, 2042-2048
- Afroze, T. et al. (2003): Calcineurin-independent regulation of plasma membrane Ca^{2+} -ATPase-4 in the vascular smooth muscle cell cycle. *Am. J. Physiol. Cell. Physiol.* 285, C88-C95
- Aharonovitz, O. et al. (1998): Stimulation of 42/44 kDa mitogen-activated protein kinases by arginine vasopressin in rat cardiomyocytes. *Biochim. Biophys. Acta* 1401, 105-111
- Aikawa, R. et al. (2002): Integrins play a critical role in mechanical stress-induced p38 MAPK activation. *Hypertension* 39, 233-238
- Albert, P. R. and Tashjian, A. H. Jr. (1984): Relationship of Thyrotropin-releasing Hormone-induced Spike and Plateau Phases in Cytosolic Free Ca^{2+} Concentrations to Hormone Secretion. *J. Biol. Chem.* 259, 15350-15363
- Albert, P. R. and Tashjian, A. H. Jr. (1986): Ionomycin acts as an ionophore to release TRH-regulated Ca^{2+} stores from GH4C1 cells. *Am. J. Physiol.* 251, C887-91
- Anger, M. et al. (1998): Cellular distribution of Ca^{2+} pumps and Ca^{2+} release channels in rat cardiac hypertrophy induced by aortic stenosis. *Circulation* 98, 2477-2486
- Aramburu, J. et al. (1998): Selective inhibition of NFAT activation by a peptide spanning the calcineurin targeting site of NFAT. *Mol. Cell* 1, 627-637
- Aramburu, J. et al. (2004): Calcineurin: a central controller of signalling in eukaryotes. *EMBO reports* 5, 343-348
- Ashrafian, H. et al. (2003): Hypertrophic cardiomyopathy: a paradigm for myocardial energy depletion. *Trends Genet.* 19, 263-268
- Auckland, L. M. et al. (1986): Cardiac myosin light and heavy chain isotypes in Tetralogy of Fallot. *Cardiovasc. Res.* 20, 828-863

- Ausubel, F. M. et al.: Current Protocols in Molecular Biology, John Wiley and Sons, Inc., 2003, Volume 1 – 4
- Barger, P. M. and Kelly, D. P. (2000): PPAR signalling in the control of cardiac energy metabolism. *Trends Cardiovasc. Med.* 10, 238-245
- Barton, P. and Buckingham, M. E. (1985): The myosin alkali light chain proteins and their genes. *Biochem. J.* 231, 249-261
- Berg, J. S. et al. (2001): A millenial myosin census. *Mol. Biol. Cell* 12, 780-794
- Berk, B. C. et al. (1987): Early agonist-mediated ionic events in cultured vascular smooth muscle cells. *J. Biol. Chem.* 262, 5065-5072
- Berne, R. M. and Levy, M. N.: Physiology, Third Edition, St. Louis, 1993
- Berridge M. J. (1993): Inositol trisphosphate and calcium signalling. *Nature* 361, 315-325
- Bird, E. et al. (2001): Significant Reduction in Cardiac Fibrosis and Hypertrophy in Spontaneously Hypertensive Rats (SHR) Treated with a V_{1a} Receptor Antagonist. *Circulation* 17, 892
- Blaeser, F. et al. (2000): Ca²⁺-dependent Gene Expression Mediated by MEF2 Transcription Factors. *J. Biol. Chem.* 275, 197-209
- Boknik, P. et al. (2001): Enhanced protein phosphorylation in hypertensive hypertrophy. *Cardiovasc. Res.* 51, 717-728
- Bowling, N. et al. (1999): Increased Protein Kinase C Activity and Expression of Ca²⁺-Sensitive Isoforms in the Failing Human Heart. *Circulation* 99, 384-391
- Bowman, J. C. et al. (1997): Expression of protein kinase C β in the heart causes hypertrophy in adult mice and sudden death in neonates. *J. Clin. Invest.* 100, 2189-95
- Braun, A. P. and Schulman, H. (1995): The multifunctional calcium/calmodulin-dependent protein kinase: From form to function. *Annu. Rev. Physiol.* 57, 417-445
- Braz, J. C. et al. (2002): PKC α regulates the hypertrophic growth of cardio-myocytes through extracellular signal regulated kinase ½ (ERK1/2). *J. Cell Biol.* 156, 905-19
- Brenner, B. (1988): Effect of Ca²⁺ on cross-bridge turnover kinetics in skinned single rabbit psoas fibers: implications for regulation of muscle contraction. *Proc. Natl. Acad. Sci. USA* 85, 3265-3269
- Brostrom, M. A. et al. (2000): Vasopressin-induced hypertrophy in H9c2 heart-derived myocytes. *Int. J. Biochem. Cell Biol.* 32, 993-1006

- Budworth, J. and Gescher, A. (1995): Differential inhibition of cytosolic and membrane-derived protein kinase C activity by staurosporine and other kinase inhibitors. FEBS Lett. 362, 139-142
- Bueno, O. F. et al. (2000): The MEK1-ERK1/2 signaling pathway promotes compensated cardiac hypertrophy in transgenic mice. EMBO J. 19, 6341-50
- Bueno, O. F. et al. (2002): Calcineurin and hypertrophic heart disease: novel insights and remaining questions. Cardiovasc. Res. 53, 806-821
- Buskin, J. N. and Hauschka, S. D. (1989): Identification of a myocyte nuclear factor that binds to the muscle-specific enhancer of the mouse muscle creatine kinase gene. Mol. Cell. Biol. 9, 2627-2640
- Caforio, A. L. et al. (1992): Identification for the α and β myosin heavy chain isoforms as major autoantigens in dilated cardiomyopathy. Circulation. 85, 1734-1742
- Caforio, A. L. and Goodwin, J. F. (1993): Aspetti attuali della patogenesi e del trattamento della cardiomiopatia dilatativa. Primary Cardiol. 8, 77-83
- Cannon B. W.: The Wisdom of the Body, 1932
- Casale, P. N. et al. (1986): Value of echocardiographic measurement of left ventricular mass in predicting cardiovascular morbid events in hypertensive men. Ann. Intern. Med. 105, 173-178
- Catala, F. et al. (1995): A Skeletal Muscle-Specific Enhancer Regulated by Factors Binding to E and CarG Boxes Is Present in the Promoter of the Mouse Myosin Light-Chain 1A Gene. Mol. Cell. Biol. 15, 4585-4596
- Chandrashekhar, Y. et al. (2003): The role of arginine vasopressin and its receptors in the normal and failing rat heart. J. Mol. Cell. Cardiol. 35, 495-504
- Chatila, T. et al. (1996): A unique phosphorylation-dependent mechanism for the activation of calcium/calmodulin-dependent protein kinase type IV/Gr. J. Biol. Chem. 271, 21542-21548
- Chen, L. et al. (1998): Structure of the DNA-binding domains from NFAT, Fos and Jun bound specifically to DNA. Nature 392, 42-48
- Chen, W. C. and Chen, C. C. (1999): Signal Transduction of Arginine Vasopressin-Induced Arachidonic Acid Release in H9c2 Cardiac Myoblasts: Role of Ca^{2+} and the Protein Kinase C-Dependent Activation of p42 Mitogen-Activated Protein Kinase. Endocrinology 140, 1639-1648
- Chin, D. and Means, A. R. (2000): Calmodulin: a prototypical calcium sensor. Trends Cell Biol. 10, 322-328

- Chien, K. R. (2000): Meeting Koch's postulates for calcium signalling in cardiac hypertrophy. *J. Clin. Invest.* 105, 1339-1342
- Choukroun, G. et al. (1998): Role of the stress-activated protein kinases in endothelin-induced cardiomyocyte hypertrophy. *J. Clin. Invest.* 102, 1311-20
- Cingolani, H. E. and Camilion de Hurtado, M. C. (2002): Na⁺-H⁺ exchanger inhibition: a new antihypertrophic tool. *Circ. Res.* 90, 751-53
- Clerk, A. et al. (1994): Differential Activation of Protein Kinase C Isoforms by Endothelin-1 and Phenylephrine and Subsequent Stimulation of p42 and p44 Mitogen-activated Protein Kinases in Ventricular Myocytes Cultured from Neonatal Rat Hearts. *J. Biol. Chem.* 269, 32848-32857
- Clerk, A. et al. (1998): Stimulation of the p38 mitogen-activated protein kinase pathway in neonatal rat ventricular myocytes by the G-protein-coupled receptor agonists, endothelin-1 and phenylephrine: a role in cardiac myocyte hypertrophy ? *J. Cell Biol.* 142, 523-535
- Coghlan, V. M. et al. (1995): Association of protein kinase A and protein phosphatase 2B with a common anchoring protein. *Science* 267, 108-111
- Cohen, A. et al. (1988): Promoter analysis of myosin alkali light chain genes expressed in mouse striated muscle. *Nucleic Acids Res.* 16, 10037-10052
- Colomer, J. M. et al. (2003): Pressure overload selectively up-regulates Ca²⁺/calmodulin-dependent protein kinase II in vivo. *Mol. Endocrinol.* 17, 183-192
- Cruzalegui, F. H. and Means, A. R. (1993): Biochemical characterization of the multifunctional Ca²⁺/calmodulin-dependent protein kinase type IV expressed in insect cells. *J. Biol. Chem.* 268, 26171-26178
- D'Angelo, D. D. et al. (1997): Transgenic G α q overexpression induces cardiac contractile failure in mice. *Proc. Natl. Acad. Sci. USA* 94, 8121-26
- Dantzig, J. A. et al. (1992): Reversal of the cross-bidge force-generating transition by photogeneration of phosphate in rabbit psoas muscle fibers. *J. Physiol.* 451, 247-278
- Daubas, P. et al. (1985): A comparison between mammalian and avian fast skeletal muscle alkali myosin light chain genes: regulatory implications. *Nucleic Acids Res.* 13, 4623-43
- Davies, M. J. and McKenna, W. J. (1995): Hypertrophic cardiomyopathy: pathology and pathogenesis. *Histopathology* 26, 493-500

- Davis, R. J. (1993): The Mitogen-activated Protein Kinase Signal Transduction Pathway. *J. Biol. Chem.* 268, 14553-14556
- Depre, C. et al. (1998): Unloaded heart in vivo replicates fetal gene expression of cardiac hypertrophy. *Nat. Med.* 4, 1269-75
- De Windt, L. J. et al. (2000a): Calcineurin promotes protein kinase C and c-Jun NH₂ terminal kinase activation in the heart. Cross-talk between cardiac hypertrophic signaling pathways. *J. Biol. Chem.* 275, 13571-13579
- De Windt, L. J. et al. (2000b): Calcineurin-mediated hypertrophy protects cardiomyocytes from apoptosis in vitro and in vivo: An apoptosis-independent model of dilated heart failure. *Circ. Res.* 86, 255-263
- Dorn II, G. W. et al. (2003): Phenotyping hypertrophy: eschew obfuscation. *Circ. Res.* 92, 1171- 1175
- Ebert, T. J. et al. (1986): Vasopressin reduces cardiac function and augments cardiopulmonary baroreflex resistance increases in man. *J. Clin. Invest.* 77, 1136-1142
- Edman, C. F. and Schulman, H. (1994): Identification and characterization of δB-CaM kinase and δC-CaM kinase from rat heart, two new multifunctional Ca²⁺/calmodulin-dependent protein kinase isoforms. *Biochim. Biophys. Acta* 1221, 89-101
- Eisenberg, E. and Hill, T. L. (1985): Muscle contraction and free energy transduction in biological systems. *Science* 227, 999-1006
- Elliott, P. and McKenna, W. J. (2004): Hypertrophic cardiomyopathy. *Lancet* 363, 1881-1891
- Exton, J. H. (1990): Signaling through phosphatidylcholine breakdown. *J. Biol. Chem.* 265, 1-4
- Fallot, A. (1888): Contribution a l'anatomie pathologique de la maladie bleue (cyanose cardiac). *Marseille Med.* 25, 418-420
- Fatkin, D. et al. (2000): An abnormal Ca²⁺ response in mutant sarcomere protein-mediated familial hypertrophic cardiomyopathy. *J. Clin. Invest.* 106, 1351-1359
- Fewell, J. G. et al. (1998): Functional Significance of Cardiac Myosin Essential Light Chain Isoform Switching in Transgenic Mice. *J. Clin. Invest.* 101, 2630-2639
- Fiedler, B. and Wollert, K. C. (2004): Interference of antihypertrophic molecules and signaling pathways with the Ca²⁺-calcineurin-NFAT cascade in cardiac myocytes. *Cardiovasc. Res.* 63, 450-457

- Fishman, J. B. et al. (1985): Internalization, recycling, and redistribution of vasopressin receptors in rat hepatocytes. *J. Biol. Chem.* 260, 12641-12546
- Fodor, W. L. et al. (1989): Human ventricular/slow twitch myosin alkali light chain gene characterization, sequence, and chromosomal location. *J. Biol. Chem.* 264, 2143-2149
- Frey, N. et. al. (2000): Decoding calcium signals involved in cardiac growth and function. *Nat. Med.* 6, 1221-27
- Frey, N. and Olson, E. N. (2003): Cardiac hypertrophy: the good, the bad, and the ugly. *Annu. Rev. Physiol.* 65, 45-79
- Fuentes, J. J. et al. (1995): A new human gene from the Down syndrome critical region encodes a proline-rich protein highly expressed in fetal brain and heart. *Hum. Mol. Genet.* 4, 1935-1944
- Fujisawa, S. and Iijima, T. (1999): On the inotropic actions of arginine vasopressin in ventricular muscle of the guinea pig heart. *Jpn. J. Pharmacol.* 81, 309-312
- Fukuzawa, J. et al. (1999): Arginine vasopressin increases the rate of protein synthesis in isolated perfused adult rat heart via the V₁ receptor. *Mol. Cell. Biochem.* 195, 93-98
- Gallot-Payet, N. et al. (1991): Involvement of protein kinase C in the coupling between the V₁-vasopressin receptor and phospholipase C in rat glomerulosa cells: effects on aldosterone secretion. *Endocrinology* 129, 623-634
- Ganguly P. K. et al. (1989): Altered sympathetic system and adrenoreceptors during the development of cardiac hypertrophy. *Am. Heart J.* 118, 520-525
- Ganguly P. K. and Sherwood G. R. (1991): Noradrenaline turnover and metabolism in myocardium following aortic constriction in rats. *Cardiovasc. Res.* 25, 579-585
- Geisterfer, A. A. F. and Owens, G. K. (1989): Arginine-vasopressin-induced hypertrophy of cultured rat aortic smooth muscle cells. *Hypertension* 14, 413-420
- Glennon, P. E. et al. (1996): Depletion of mitogen-activated protein kinase using an antisense oligodeoxynucleotide approach downregulates the phenylephrine-induced hypertrophic response in rat cardiac myocytes. *Circ. Res.* 78, 954-61
- Goldsmith, S. R. (1987): Vasopressin as vasopressor. *Am. J. Med.* 82, 1213-1219
- Goldsmith, S. R. (2002): Vasopressin antagonists in CHF: ready for clinical trials? *Cardiovasc. Res.* 54, 13-15
- Grunig, E. et al. (1998): Frequency and phenotypes of familial dilated cardiomyopathy. *J. Am. Coll. Cardiol.* 31, 186-194

- Gruver, C. L. et al. (1993): Targeted developmental overexpression of calmodulin induces proliferative and hypertrophic growth of cardiomyocytes in transgenic mice. *Endocrinology* 133, 376-388
- Hagemann, D. et al. (2001): Expression of Ca^{2+} /calmodulin-dependent protein kinase II δ -subunit isoforms in rats with hypertensive cardiac hypertrophy. *Mol. Cell. Biochem.* 220, 69-76
- Han, J. et al. (1997): Activation of the transcription factor MEF2C by the MAP kinase p38 in inflammation. *Nature* 386, 296-99
- Harada, K. et al. (1998): Pressure overload induces cardiac hypertrophy in angiotensin II type 1A receptor knockout mice. *Circulation* 97, 1952-1959
- Hayashi, A. et al. (1999): PKC ν , a new member of the protein kinase C family, composes a fourth subfamily with PKC μ . *Biochim. Biophys. Acta* 1450, 99-106
- Heist, E. K. and Schulman, H. (1998): The role of Ca^{2+} /calmodulin-dependent protein kinases within the nucleus. *Cell Calcium* 23, 103-114
- Heist, E. K. et al. (1998): Phosphorylation at the nuclear localization signal of Ca^{2+} /calmodulin-dependent protein kinase II blocks its nuclear targeting. *J. Biol. Chem.* 273, 19763-19771
- Herzig, T. C. et al. (1997): Angiotensin II type 1a receptor gene expression in the heart: AP-1 and GATA-4 participate in the response to pressure overload. *Proc. Natl. Acad. Sci. USA* 94, 7543-7548
- Hescheler, J. et al. (1991): Morphological, Biochemical, and Electrophysiological Characterization of a Clonal Cell (H9c2) Line From Rat Heart. *Circ. Res.* 69, 1476-1486
- Hirota, H. et al. (1999): Loss of a gp130 cardiac muscle cell survival pathway is a critical event in the onset of heart failure during biomechanical stress. *Cell* 97, 189-98
- Ho, N. et al. (1996): Activation Protein-1-Dependent Transcriptional Activation of Interleukin 2 Gene by Ca^{2+} /Calmodulin Kinase Type IV/Gr. *J. Exp. Med.* 184, 101-112
- Hoch, B. et al. (1998): Differentiation-Dependent Expression of Cardiac δ -CaMKII Isoforms. *J. Cell. Biochem.* 68, 259-268
- Hoch, B., et al. (1999): Identification and expression of δ -isoforms of the multifunctional Ca^{2+} /calmodulin-dependent protein kinase in failing and nonfailing human myocardium. *Circ. Res.* 84, 713-721

- Hoey, T. et al. (1995): Isolation of two new members of the NF-AT gene family and functional characterization of the NF-AT proteins. *Immunity* 2, 461-472
- Hogan, P. G. et al. (2003): Transcriptional regulation by calcium, calcineurin, and NFAT. *Genes Dev.* 17, 2205-2232
- Hollenberg, S. M. et al. (1995): Identification of a New Family of Tissue-Specific Basic Helix-Loop-Helix Proteins with a Two-Hybrid System. *Mol. Cell. Biol.* 15, 3813-3822
- Holmes, K. C. (1997): The swinging lever-arm hypothesis of muscle contraction. *Curr. Biol.* 7, R112-R118
- Hook, S. S. and Means, A. R. (2001): Ca²⁺/CaM-dependent kinases: from activation to function. *Annu. Rev. Pharmacol. Toxicol.* 41, 471-505
- Howard, J. and Spudich, J. A. (1996): Is the lever arm of myosin a molecular elastic element? *Proc. Natl. Acad. Sci. USA* 93, 4462-4464
- Hu, H. and Sachs, F. (1997): Stretch-activated ion channels in the heart. *J. Mol. Cell. Cardiol.* 29, 1511-1523
- Huang, L. et al. (2001): Increased contractility and altered Ca²⁺ transients of mouse heart myocytes conditionally expressing PKCβ. *Am. J. Physiol. Cell. Physiol.* 280, C1114-C20
- Hudmon, A. and Schulman, H. (2002): Neuronal Ca²⁺/calmodulin-dependent protein kinase II: the role of structure and autoregulation in cellular function. *Annu. Rev. Biochem.* 71, 473-510
- Hupf, H. et al. (1999): Evidence for a Vasopressin System in the Rat Heart. *Circ. Res.* 84, 365-370
- Hynes, R. O. (1992): Integrins: versatility, modulation and signaling in cell adhesion. *Cell* 69, 11-25
- Ichida, M. and Finkel, T. (2001): Ras regulates NFAT3 activity in cardiac myocytes. *J. Biol. Chem.* 276, 3524-3530
- Im, S. H. and Rao, A. (2004): Activation and Deactivation of Gene Expression by Ca²⁺/Calcineurin-NFAT-mediated Signaling. *Mol. Cells* 18, 1-9
- Innis M. S. et al. (1988): DNA Sequencing with *Thermus Acquaticus* DNA Polymerase and Direct Sequencing of Polymerase chain reaction-amplified DNA. *Proc. Natl. Acad. Sci. U.S.A.* 85, 9436-9440
- Irving, M. et al. (1995): Tilting of the light chain region of myosin during step length changes and active force generation in skeletal muscle. *Nature* 375, 688-691

- Isoda, K. et al. (2001): Myocardial hypertrophy in transgenic mice overexpressing human interleukin 1 α . *J. Card. Fail.* 7, 355-364
- Izumo, S. et al. (1988): Proto-oncogene induction and reprogramming of cardiac gene expression produced by pressure overload. *Proc. Natl. Acad. Sci. USA* 85, 339-343
- James, M. A. et al. (2000): Wall stress and hypertension. *J. Cardiovasc. Risk* 7, 187-190
- Jans, D. A. et al. (1990): Lateral mobility of the phospholipase C-activating vasopressin V₁-type receptor in A7r5 smooth muscle cells: a comparison with the adenylate cyclase-coupled V₂-receptor. *EMBO J.* 9, 2693-2699
- Jensen, K. F. et al. (1991): Nuclear and axonal localization of Ca²⁺/calmodulin-dependent protein kinase type Gr in rat cerebellar cortex. *Proc. Natl. Acad. Sci. USA* 88, 2850-2853
- Kanaseki, T. et al. (1991): Structural features of Ca²⁺/calmodulin-dependent protein kinase II revealed by electron microscopy. *J. Cell Biol.* 115, 1049-1060
- van Kerckhoven, R. et al. (2002): Chronic vasopressin V_{1A} but not V₂ receptor antagonism prevents heart failure in chronically-infarcted rats. *Eur. J. Pharmacol.* 449, 135-141
- Khayyal, M. A. et al. (1985): Effects of vasopressin on the coronary circulation: reserve and regulation during ischemia. *Am. J. Physiol.* 248, H516-522
- Kim, J. M. et al. (1999): Phosphatidylinositol 3-Kinase Regulates Differentiation of H9c2 Cardiomyoblasts Mainly through the Protein Kinase B/Akt-Independent Pathway. *Arch. Biochem. Biophys.* 367, 67-73
- Kimes, B. W. and Brandt, B. L. (1976): Properties of a Clonal muscle cell line from rat heart. *Exp. Cell Res.* 98, 367-381
- Kingsbury, T. J. and Cunningham, K. W. (2000): A conserved family of calcineurin regulators. *Genes Dev.* 14, 1595-1604
- Kirchhefer, U. et al. (1999): Activity of cAMP-dependent protein kinase and Ca²⁺/calmodulin-dependent protein kinase in failing and nonfailing human hearts. *Cardiovasc. Res.* 42, 254-261
- Klee, C. B. Interaction of calmodulin with Ca²⁺ and target proteins. In: *Calmodulin*. Cohen, P. and Klee, C. B. eds. The Netherlands: Elsevier, Amsterdam, 1988
- Klee, C. B. et al. (1988): Calcineurin. *Adv Enzymol Relat Areas Mol Biol* 61, 149-200
- Klee, C. B. et al. (1998): Regulation of the calmodulin-stimulated protein phosphatase, calcineurin. *J. Biol. Chem.* 273, 13367-13370

- Knowlton, K. U. et al. (1993): The α 1A-adrenergic receptor subtype mediates biochemical, molecular, and morphologic features of cultured myocardial cell hypertrophy. *J. Biol. Chem.* 268, 15374-15380
- Kolodziejczyk, S. M. et al. (1999): MEF2 is upregulated during cardiac hypertrophy and is required for normal post-natal growth of the myocardium. *Curr. Biol.* 9, 1203-1206
- Komuro, I. et al. (1988): Expression of cellular oncogenes in the myocardium during the developmental stage and pressure-overload hypertrophy of the rat heart. *Circ. Res.* 62, 1075-1079
- Komuro, I. et al. (1996): Mechanical stretch activates the stress-activated protein kinases in cardiac myocytes. *FASEB J.* 10, 631-36
- Kovacic, J. C. and Muller, D. et al. (2003): Hypertrophic cardiomyopathy: state-of-the-art review, with focus on the management of outflow obstruction. *Intern. Med. J.* 33, 521-529
- Kubota, T. et al. (1997): Cardiac-specific overexpression of tumor necrosis factor- α causes lethal myocarditis in transgenic mice. *J. Card. Fail.* 3, 117-124
- Kurabayashi, M. et al. (1988): Molecular cloning and characterization of human atrial and ventricular myosin alkali light chain cDNA clones. *J. Biol. Chem.* 263, 13930-13936
- Kusumoto, K. et al. (2001): Na^+/H^+ exchange inhibition reduces hypertrophy and heart failure after myocardial infarction in rats. *Am. J. Physiol. Heart Circ. Physiol.* 280, H738-H745
- Kuwahara, F. (2002): Transforming growth factor- β function blocking prevents myocardial fibrosis and diastolic dysfunction in pressure-overloaded rats. *Circulation* 106, 130-135
- Laemmli, U. K. (1970): Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680-685
- Lai, M. M. et al. (1998): Cain, a novel physiologic protein inhibitor of calcineurin. *J. Biol. Chem.* 273, 18325-18331
- Lapointe, M. C. et al. (1990): Extracellular calcium regulates expression of the gene for atrial natriuretic factor. *Hypertension*. 15, 20-28
- Lassar, A. B. et al. (1989): MyoD is a sequence-specific DNA binding protein requiring a region of myc homology to bind to the muscle creatine kinase enhancer. *Cell* 58, 823-831

- Lee, C. R. et al. (2003): Vasopressin: A new target for the treatment of heart failure. *Am. Heart J.* 146, 9-18
- Lee, J. C. and Edelman, A. M. (1994): A protein activator of Ca^{2+} -calmodulin-dependent protein kinase1a. *J. Biol. Chem.* 269, 2158-2164
- Leinwand, L. (2001): Calcineurin inhibition and cardiac hypertrophy: A matter of balance. *Proc. Natl. Acad. Sci. USA* 98, 2947-2949
- Leinwand, L. A. et al. (1983): Isolation and characterization of human myosin heavy chain genes. *Proc. Natl. Acad. Sci. USA* 80, 3716-3720
- Lembo, G. et al. (1996): Elevated blood pressure and enhanced myocardial contractility in mice with severe IGF-1 deficiency. *J. Clin. Invest.* 98, 2648-2655
- Lewis, J. M. and Schwartz, A. M. (1995): Mapping in vivo associations of cytoplasmic proteins with integrin $\beta 1$ cytoplasmic domain mutants. *Mol. Biol. Cell.* 6, 151-160
- Levy, D. et al. (1990): Prognostic implications of echocardiographically determined left ventricular mass in the Framingham Heart Study. *N. Engl. J. Med.* 322, 1561-1566
- Li, Q. et al. (1997): Overexpression of insulin-like growth factor-1 in mice protects from myocyte death after infarction, attenuating ventricular dilation, wall stress, and cardiac hypertrophy. *J. Clin. Invest.* 100, 1991-1999
- Lin, Q. et al. (1997): Control of Mouse Cardiac Morphogenesis and Myogenesis by Transcription Factor MEF2C. *Science* 276, 1404-1407
- Lips, D. J. et al. (2003): Molecular determinants of myocardial hypertrophy and failure: alternative pathways for beneficial and maladaptive hypertrophy. *Eur. Heart J.* 24, 883-896
- Lipshultz, S. E. et al. (2003): The incidence of pediatric cardiomyopathy in two regions of the United States. *N. Engl. J. Med.* 348, 1647-1655
- Liu, C. and Hermann, T. E. (1978): Characterization of Ionomycin as a Calcium Ionophore. *J. Biol. Chem.* 253, 5892-5894
- Liu, P. et al. (1999): Vasopressin-Evoked $[\text{Ca}^{2+}]_i$ Responses in Neonatal Rat Cardiomyocytes. *J. Cardiovasc. Pharmacol.* 34, 540-546
- Lombardi, V. et al. (1995): Elastic distortion of myosin heads and repriming of the working stroke in muscle. *Nature* 374, 553-555
- Louie, E. K. and Edwards III, L. C. (1994): Hypertrophic cardiomyopathy. *Prog. Cardiovasc. Dis.* 36, 275-308
- Lu, J. et al. (2000): Signal-dependent activation of the MEF2 transcription factor by dissociation from histone deacetylases. *Proc. Natl. Acad. Sci. USA* 97, 4070-4075

- Lutz, W. et al. (1991): Vasopressin receptor-mediated endocytosis: current review. *Am. J. Physiol.* 261, F1-F13
- Lyakh, L. et al. (1997): Expression of NFAT-Family Proteins in Normal Human T Cells. *Mol. Cell Biol.* 17, 2475-2484
- Macián, F. et al. (2001): Partners in transcription: NFAT and AP-1. *Oncogene* 20, 2476-2489
- Maier, L. S. and Bers, D. M. (2002): Calcium Cycling. Calcium, Calmodulin, and Calcium-Calmodulin Kinase II: Heartbeat to Heartbeat and Beyond. *J. Mol. Cell. Cardiol.* 34, 919-939
- Marks, A. R. (2003): Calcium and the heart: a question of life and death. *J. Clin. Invest.* 11, 597-600
- Maron, B. J. (1997): Hypertrophic cardiomyopathy. *Lancet* 350, 127-133
- Maron, B. J. (2002): Hypertrophic cardiomyopathy: a systematic review. *JAMA* 287, 1308-1320
- Maron, B. J. et al. (1980): Sudden death in young athletes. *Circulation* 62, 218-229
- Maron, B. J. et al. (1995): Cardiac disease in young trained athletes. Insights into methods for distinguishing athlete's heart from structural heart disease, with particular emphasis on hypertrophic cardiomyopathy. *Circulation* 91, 1569-1601
- Martino, T. A. et al.: Enterovirus myocarditis and dilated cardiomyopathy: a review of clinical and experimental studies. In: Rotbart, H. A., Editor. *Human Virus Infections*, Washington DC: Medical Society for Microbiology, 1995, 291-350
- Martiny-Baron, G. et al. (1993): Selective Inhibition of Protein Kinase C Isoenzymes by the Indolocarbozole Gö 6976. *J. Biol. Chem.* 268, 9194-9197
- Mayer, P. et al. (1995): Novel and uncommon isoforms of the calcium sensing enzyme calcium/calmodulin-dependent protein kinase II in heart tissue. *Basic Res. Cardiol.* 90, 372-379
- McDonough, P. M. and Glembotski, C. C. (1992): Induction of atrial natriuretic factor and myosin light chain-2 gene expression in cultured ventricular myocytes by electrical stimulation of contraction. *J. Biol. Chem.* 267, 11665-11668
- McKenna, W. J. et al. (1998): Genes and disease expression in hypertrophic cardiomyopathy. *Lancet* 352, 1162-1163
- McKinsey, T. A. et al. (2000): Activation of the myocyte enhancer factor-2 transcription factor by calcium/calmodulin-dependent protein kinase-stimulated binding of 14-3-3 to histone deacetylase 5. *Proc. Natl. Acad. Sci. USA* 97, 14400-14405

- McKinsey, T. A. et al. (2002): MEF2: a calcium-dependent regulator of cell division, differentiation and death. *Trends Biochem. Sci.* 27, 40-47
- Means, A. R. et al. (1991): A novel Ca^{2+} /calmodulin-dependent protein kinase and a male germ cell-specific calmodulin-binding protein are derived from the same gene. *Mol. Cell. Biol.* 11, 3960-3971
- Meerson, F. Z. and Javich, M. P. (1982): Isoenzyme pattern and activity of myocardial creatine phosphokinase under heart adaptation to prolonged overload. *Basic Res. Cardiol.* 77, 349-358
- Mejía-Alvarez, R. et al. (1994): Simultaneous expression of cardiac and skeletal muscle isoforms of the L-type Ca^{2+} channel in a rat heart muscle cell line. *J. Physiol.* 478, 315-329
- Mercadier, J. J. et al. (1989): Atrial natriuretic factor gene expression in rat ventricle during experimental hypertension. *Am. J. Physiol.* 257, H979-H987
- Milano, C. A. et al. (1994): Myocardial expression of a constitutively active $\alpha 1\text{B}$ -adrenergic receptor in transgenic mice induces cardiac hypertrophy. *Proc. Natl. Acad. Sci. USA* 91, 10109-10113
- Mochly-Rosen, D. et al (1991): Identification of intracellular receptor proteins for activated protein kinase C. *Proc. Natl. Acad. Sci. USA* 88, 3997-4000
- Mogensen, J. et al. (2003): Idiopathic restrictive cardiomyopathy is part of the clinical expression of cardiac Troponin I mutations. *J. Clin. Invest.* 111, 209-216
- Molkentin, J. D. (2000): Calcineurin and beyond: cardiac hypertrophic signaling. *Circ. Res.* 87, 731-738
- Molkentin, J. D. and Dorn II, G. W. (2001): Cytoplasmic signaling pathways that regulate cardiac hypertrophy. *Annu. Rev. Physiol.* 63, 391-426
- Molkentin J. D. and Olson E. N. (1997): GATA4: a novel transcriptional regulator of cardiac hypertrophy ? *Circulation* 96, 3833-3835
- Molkentin, J. D. et al. (1998): A calcineurin-dependent transcriptional pathway for cardiac hypertrophy. *Cell* 93, 215-228
- Moncrief, N. D. et al. (1990): Evolution of EF-hand calcium-modulated proteins: Relationships based on amino acid sequences. *J. Mol. Evol.* 30, 522-562
- Morano, I. (1999): Tuning the human heart molecular motors by myosin light chains. *J. Mol. Med.* 77, 544-555
- Morano, I. and Haase, H. (1997): Different actin affinities of human cardiac essential myosin light chain isoforms. *FEBS Lett.* 408, 71-74

- Morano, I. et al. (1997): Changes in essential myosin light chain isoform expression provide a molecular basis of isometric force regulation in the failing human heart. *J Mol. Cell. Cardiol.* 29, 1177-1187
- Morano, M. et al. (1996): Regulation of Human Heart Contractility by Essential Myosin Light Chain Isoforms. *J. Clin. Invest.* 98, 467-473
- Mulvagh, S. L. et al. (1987): A hemodynamic load in vivo induces cardiac expression of the cellular oncogene, c-myc. *Biochem. Biophys. Res. Commun.* 147, 627-636
- Nagai, R. et al. (1989): Regulation of myocardial Ca^{2+} -ATPase and phospholamban mRNA expression in response to pressure overload and thyroid hormone. *Proc. Natl. Acad. Sci. USA* 86, 2966-2970
- Nagueh, S. F. et al. (1999): Changes in left ventricular diastolic function 6 months after nonsurgical septal reduction therapy for hypertrophic obstructive cardiomyopathy. *Circulation* 99, 344-347
- Nairn, A. C. et al. (1985): Identification of calmodulin-dependent protein kinase III and its major Mr. 100,000 substrate in mammalian tissues. *Proc. Natl. Acad. Sci. USA* 82, 7939-7943
- Nairn, A. C. and Greengard, P. (1987): Purification and characterization of Ca^{2+} /calmodulin-dependent protein kinase I from bovine brain. *J. Biol. Chem.* 262, 7273-7281
- Nakamura Y. et al. (2000): Hypertrophic growth of cultured neonatal rat heart cells mediated by vasopressin V_{1A} receptor. *Eur. J. Pharmacol.* 391, 39-48
- Naya, F. J. et al. (1999): Transcriptional activity of MEF2 during mouse embryogenesis monitored with a MEF2-dependent transgene. *Development* 126, 2045-2052
- Neumann, D. A. (1994): Autoimmunity and dilated cardiomyopathy. *Mayo Clin. Proc.* 69, 193-195
- Nielsen, S. et al. (1999): Physiology and pathophysiology of renal aquaporins. *J. Am. Soc. Nephrol.* 10, 647-663
- Nishimura, R. A. et al. (2003): Hypertrophic Cardiomyopathy/A Patient Perspective. *Circulation* 108, e133-e135
- Nugent, A. W. et al. (2003): The Epidemiology of Childhood Cardiomyopathy in Australia. *N. Engl. J. Med.* 348, 1639-1646
- Okamura, H. et al. (2000): Concerted Dephosphorylation of the Transcription Factor NFAT1 Induces a Conformational Switch that Regulates Transcriptional Activity. *Mol. Cell* 6, 539-550

- Olson, E. N. and Klein, W. H. (1994): bHLH factors in muscle development: dead lines and commitments, what to leave in and what to leave out. *Genes Dev.* 8, 1-8
- Olson, E. N. and Molkentin, D. J. (1999): Prevention of cardiac hypertrophy by calcineurin inhibition. *Circ. Res.* 84, 623-632
- Olson, E. N. and Williams, R. S. (2000): Calcineurin signaling and muscle remodeling. *Cell* 101, 689-692
- Pass, J. M. et al. (2001): PKC ϵ activation induces dichotomous cardiac phenotypes and modulates PKC ϵ -RACK interactions and RACK expression. *Am. J. Physiol. Heart Circ. Physiol.* 280, H946-H955
- Passier, R. et al. (2000): CaM kinase signaling induces cardiac hypertrophy and activates the MEF2 transcription factor in vivo. *J. Clin. Invest.* 105, 1395-1406
- Pellieux, C. et al. (2001): Dilated cardiomyopathy and impaired cardiac hypertrophic response to angiotensin II in mice lacking FGF-2. *J. Clin. Invest.* 108, 1843-1851
- Poche R.: Strukturelle Veränderungen des druck- und volumenüberlasteten Herzens. In: Herzkrankheiten. Pathophysiologie, Diagnostik, Therapie. H. Roskamm, H. Reindell, 4. Auflage, Springer Verlag, Berlin, 1996, 414-424
- Pollard, T. D. and Earnshaw, W. C.: Cell Biology, Saunders (Imprint of Elsevier), Philadelphia, Pennsylvania, 2004
- Post, G. R. et al. (1996): Dissociation of p44 and p42 mitogen-activated protein kinase activation from receptor-induced hypertrophy in neonatal rat ventricular myocytes. *J. Biol. Chem.* 271, 8452-57
- Price, K. M. et al. (1980): Human atrial and ventricular myosin light-chains subunits in the adult and during development. *Biochem. J.* 191, 571-580
- Ramirez, M. T. et al. (1997): The nuclear δ B isoform of Ca^{2+} /calmodulin-dependent protein kinase II regulates atrial natriuretic factor gene expression in ventricular myocytes. *J. Biol. Chem.* 272, 31203-31208
- Ranger, A. M. et al. (1998): The transcription factor NF-ATc is essential for cardiac valve formation. *Nature* 392, 186-190
- Rao, A. et al. (1997): Transcription factors of the NFAT Family: Regulation and Function. *Annu. Rev. Immunol.* 15, 707-747
- Rayment, I. et al. (1995): Structural interpretation of the mutations in the β -cardiac myosin that have been implicated in familial hypertrophic cardiomyopathy. *Proc. Natl. Acad. Sci. USA* 92, 3864-3868

- Redwood, C. S. et al. (1999): Properties of mutant contractile proteins that cause hypertrophic cardiomyopathy. *Cardiovasc. Res.* 44, 20-36
- Revis, N. W. et al. (1977): Lactate dehydrogenase isoenzymes in the human hypertrophic heart. *Cardiovasc. Res.* 11, 172-176
- Ritter, O. et al. (1999a): Expression of atrial myosin light chains but not α -myosin heavy chains is correlated in vivo with increased ventricular function in patients with hypertrophic obstructive cardiomyopathy. *J. Mol. Med.* 77, 677-685.
- Ritter, O. et al. (1999b): Remodeling of the Hypertrophied Human Myocardium by Cardiac bHLH Transcription Factors. *J. Cell. Biochem.* 74, 551-561
- Ritter, O. et al. (2002): Calcineurin in Human Heart Hypertrophy. *Circulation* 105, 2265-2269
- Roman, B. B. et al. (2001): PKC- β is not necessary for cardiac hypertrophy. *Am. J. Physiol. Heart Circ. Physiol.* 280, H2264-H2270
- Rooney, J. W. et al. (1995): Coordinate and cooperative roles for NF-AT and AP-1 in the regulation of the murine IL-4 gene. *Immunity* 2, 473-483
- Ross, R. S. (2004): Molecular and mechanical synergy: cross-talk between integrins and growth factor receptors. *Cardiovasc. Res.* 63, 381-390
- Rothermel, B. et al. (2000): A protein encoded within the Down syndrome critical region is enriched in striated muscles and inhibits calcineurin signaling. *J. Biol. Chem.* 275, 8719-8725
- Rothermel, B. et al. (2001): Myocyte-enriched calcineurin-interacting protein, MCIP1, inhibits cardiac hypertrophy in vivo. *Proc. Natl. Acad. Sci. USA* 98, 3328-3333
- Rotter, M. et al. (1991): The human embryonic myosin alkali light chain gene: use of alternative promoters and 3' non-coding regions. *Nucleic Acids Res.* 19, 1497-1504
- Ruknudin, A. et al. (1993): Stretch-activated ion channels in tissue-cultured chick heart. *Am. J. Physiol.* 264, H960-H972
- Ruwhof, C. and Van der Laarse, A. (2000): Mechanical stress-induced cardiac hypertrophy: mechanisms and signal transduction pathways. *Cardiovasc. Res.* 47, 23-37
- Sadoshima, J. and Izumo, S. (1995): Rapamycin selectively inhibits angiotensin II-induced increase in protein synthesis in cardiac myocytes in vitro. Potential role of 70-kD S6 kinase in angiotensin II-induced cardiac hypertrophy. *Circ. Res.* 77, 1040-1052

- Sadoshima, J. and Izumo, S. (1997): The cellular and molecular response of cardiac myocytes to mechanical stress. *Annu. Rev. Physiol.* 59, 551-571
- Sadoshima, J. et al. (1993): Autocrine release of angiotensin II mediates stretch-induced hypertrophy of cardiac myocytes in vitro. *Cell* 75, 977-984
- Saito, T. et al. (2003): Roles of calcineurin and calcium/calmodulin-dependent protein kinase II in pressure overload-induced cardiac hypertrophy. *J. Mol. Cell. Cardiol.* 35, 1153-1160
- Sakagami, H. and Kondo, H. (1993): Cloning and sequencing of a gene encoding the beta polypeptide of Ca^{2+} /calmodulin-dependent protein kinase IV and its expression confined to the mature cerebellar granule cells. *Brain Res. Mol. Brain* 19, 215-218
- Sambrook, J. et al.: *Molecular Cloning, A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Plainview, New York, 1989, Volume 1 – 3
- Sampey, D. B. et al. (1999): Vasopressin V_2 receptor enhances gain of baroreflex in conscious spontaneously hypertensive rats. *Am. J. Physiol.* 276, R872-879
- Sandler, H. and Dodge, H. T. (1963): Left ventricular tension and stress in man. *Circ. Res.* 13, 91-104
- Sangiorgi, M. (2003): Clinical and epidemiological aspects of cardiomyopathies: a critical review of current knowledge. *Eur. J. Intern. Med.* 14, 5-17
- Schaub, M. C. et al. (1984): Myosin isoenzymes in human hypertrophic heart. Shift of atrial myosin heavy chains and ventricular light chains. *Eur. Heart J.* 5, 85-93
- Schmidt R. F. and Thews G.: *Physiologie des Menschen*, 26th edition, Springer Verlag, Berlin, 1995
- Schultz, J. E. et al. (1999): Fibroblast growth factor-2 mediates pressure-induced hypertrophic response. *J. Clin. Invest.* 104, 709-719
- Schultz Jel, J. et al. (2002): TGF- β 1 mediates the hypertrophic cardiomyocyte growth induced by angiotensin II. *J. Clin. Invest.* 109, 787-796
- Schunkert, H. et al. (1990): Increased rat cardiac angiotensin converting enzyme activity and mRNA expression in pressure overload left ventricular hypertrophy. Effects on coronary resistance, contractility, and relaxation. *J. Clin. Invest.* 86, 1913-1920
- Schwartz, K. et al. (1986): α -Skeletal muscle actin mRNAs accumulate in hypertrophied adult rat hearts. *Circ. Res.* 59, 551-555
- Seharaseyon, J. et al. (1990): Human Embryonic/Atrial Myosin Alkali Light Chain Gene: Characterization, Sequence, and Chromosomal Location. *Genomics* 7, 289-293

- Sei, C. A. et al. (1991): α -Adrenergic stimulation of atrial natriuretic factor expression in cardiac myocytes requires calcium influx, protein kinase C and calmodulin-regulated pathways. *J. Biol. Chem.* 266, 15910-15916
- Sellers, J. R.: Myosin. 2nd edn, Protein Profile. Oxford University Press, Oxford, 1999
- Sellmayer, A. et al. (1991): Endogenous non-cyclooxygenase metabolites of arachidonic acid modulate growth and mRNA levels of immediate-early response genes in rat mesangial cells. *J. Biol. Chem.* 266, 3800-3807
- Shapiro, L. M. and McKenna, W. J. (1983): Distribution of left ventricular hypertrophy in hypertrophic cardiomyopathy: a two-dimensional echocardiographic study. *J. Am. Coll. Cardiol.* 2, 437-444
- Shaw, K.T.-Y. et al. (1995): Immunosuppressive drugs prevent a rapid dephosphorylation of transcription factor NFAT1 in stimulated immune cells. *Proc. Natl. Acad. Sci. USA* 92, 11205-11209
- Sheng, M. et al. (1991): CREB: a Ca^{2+} -regulated transcription factor phosphorylated by calmodulin-dependent kinases. *Science* 252, 1427-1430
- Simmerman, H. K. B. et al. (1986): Sequence analysis of phospholamban: Identification of phosphorylation sites and two major structural domains. *J. Biol. Chem.* 261, 13333-13341
- Siri, F. M. (1988): Sympathetic changes during development of cardiac hypertrophy in aortic-constricted rats. *Am. J. Physio. Heart Circ. Physiol.* 255, H452-H457
- Soderling, T. R. (1999): The Ca^{2+} -calmodulin-dependent protein kinase cascade. *Trends Biochem. Sci.* 24, 232-236
- Soderling, T. R. et al. (2001): Cellular Signaling through Multifunctional Ca^{2+} /calmodulin-dependent Protein kinase II. *J. Biol. Chem.* 276, 3719-3722
- Sonnenberg, H. (1986): Mechanisms of release and renal tubular action of atrial natriuretic factor. *Fed. Proc.* 45, 2106-2110
- Spirito, P. et al. (1997): The management of hypertrophic cardiomyopathy. *N. Engl. J. Med.* 336, 775-785
- Srinivasan, M. et al. (1994): Alternative splicing introduces a nuclear localization signal that targets multifunctional CaM kinase to the nucleus. *J. Cell Biol.* 126, 839-852
- Srivastava, D. et al. (1995): A subclass of bHLH proteins required for cardiac morphogenesis. *Science* 270, 1995-1999
- Stepkowski, D. (1995): The role of skeletal muscle myosin light chain N-terminal fragments. *FEBS Lett.* 374, 6-11

- Sütsch, G. et al. (1992): Hemodynamic performance and myosin light chain-1 expression in the hypertrophied left ventricle in aortic valve disease before and after valve replacement. *Circ. Res.* 70, 1035-1043
- Sugden, P. H. and Clerk, A. (1998): «Stress-responsive» mitogen-activated protein kinases (c-Jun N-terminal kinases and p38 mitogen-activated protein kinases) in the myocardium. *Circ. Res.* 83, 345-352
- Sun, L. et al. (2000): Activation of HSF and selective increase in heat-shock proteins by acute dexamethasone treatment. *Am. J. Physiol.* 278, H1091-H1097
- Sun, P. et al. (1994): Differential activation of CREB by Ca^{2+} /calmodulin-dependent protein kinases type II and type IV involves phosphorylation of a site that negatively regulates activity. *Genes Dev.* 8, 2527-2539
- Sun, P. et al. (1996): Regulation of Activating Transcription Factor-1 and the cAMP Response Element-binding protein by Ca^{2+} /calmodulin-dependent Protein Kinases Type I, II and IV. *J. Biol. Chem.* 271, 3066-3073
- Sweeney, H. L. (1995): Function of the N-terminus of the myosin essential light chain of vertebrate striated muscle. *Biophys. J.* 68, 112-119
- Swynghedauw, B. (1999): Molecular Mechanisms of Myocardial Remodeling. *Physiol. Rev.* 79, 215-262
- Tahara, A. et al. (1998): Effect of YM087, a potent nonpeptide vasopressin antagonist, on vasopressin-induced protein synthesis in neonatal rat cardiomyocyte. *Cardiovasc. Res.* 38, 198-205
- Takeishi, Y. et al. (2000): Transgenic overexpression of constitutively active protein kinase C ϵ causes concentric cardiac hypertrophy. *Circ. Res.* 86, 1218-23
- Takeishi, Y. et al. (2001): Src and multiple MAP kinase activation in cardiac hypertrophy and congestive heart failure under chronic pressure-overload: comparison with acute mechanical stretch. *J. Mol. Cell. Cardiol.* 33, 1637-48
- Takewaki, S. et al. (1995): Activation of Na^+-H^+ antiporter (NHE-1) gene expression during growth, hypertrophy and proliferation of the rabbit cardiovascular system. *J. Mol. Cell. Cardiol.* 27, 729-42
- Tarone, G. and Lembo, G. (2003): Molecular interplay between mechanical and humoral signalling in cardiac hypertrophy. *Trends in Mol. Med.* 9, 376-382
- Thibonnier, M. (1992): Signal transduction of V₁-vascular vasopressin receptors. *Regul. Pept.* 38, 1-11

- Thibonnier, M. (2004): Genetics of Vasopressin Receptors. *Curr. Hypertens. Rep.* 6, 21-26
- Thibonnier, M. et al. (1987): The human platelet vasopressin receptor and its intracellular messengers: a key role of divalent cations. *J. Cardiovasc. Pharmacol.* 10, 24-29
- Thibonnier, M. et al. (1991a): Multiple signaling pathways of V₁-vascular AVP receptors of A7r5 cells. *Endocrinology* 129, 2845-2856
- Thibonnier, M. et al. (1991b): V₁-vascular AVP receptors of A7r5 cells activate the arachidonic acid cyclooxygenase and epoxygenase pathways. *J. Am. Soc. Nephrol.* 2, 432
- Thibonnier, M. et al.: Structure of the human V_{1a} vasopressin receptor gene. In : Neurohypophysis: Recent Progress of Vasopressin and Oxytocin Research, edited by Saito, T., Kurokawa, T., Yoshida, S. Elsevier Science, Amsterdam, 1995, 553-571
- Thibonnier, M. et al. (1996): Structure, Sequence, Expression, and Chromosomal Localization of the Human V_{1a} Vasopressin Receptor Gene. *Genomics* 31, 327-334
- Thibonnier, M. et al.: Signal transduction pathways of the human V₁-vascular, V₂-renal, V₃-pituitary AVP, and oxytocin receptors. In: Progress in Brain Research, edited by Urban, I. J. A., Burbach, J. P. H. and De Wied, D. The Netherlands: Elsevier Science, Amsterdam, 1998, 143-158
- Thomas, T. et al. (1998): The bHLH Factors, dHAND and eHAND, specify pulmonary and systemic cardiac ventricles independently of left-right sidedness. *Dev. Biol.* 196, 228-236
- Tian, R. et al. (1999): Long-term expression of protein kinase C in adult mouse hearts improves postischemic recovery. *Proc. Natl. Acad. Sci. USA* 96, 13536-41
- Tobimatsu, T. and Fujisawa, H. (1989): Tissue specific expression of four types of rat calmodulin-dependent protein kinase II transcripts. *J. Biol. Chem.* 264, 17907-17912
- Tokumitsu, H. et al. (1995): Characterization of a Ca²⁺/calmodulin-dependent protein kinase cascade. Molecular cloning and expression of calcium/calmodulin-dependent protein kinase. *J. Biol. Chem.* 270, 19320-19324
- Tombes, R. M. et al. (1995): G₁ Cell Cycle Arrest and Apoptosis Are Induced in NIH 3T3 Cells by KN-93, an Inhibitor of CaMK-II (the Multifunctional Ca²⁺/CaM Kinase). *Cell Growth Differ.* 6, 1063-1070

- Toullec, D. et al. (1991): The Bisindolylmaleimide GF 109203X Is a Potent and Selective Inhibitor of Protein Kinase C. *J. Biol. Chem.* 266, 15771-15781
- Towbin, J. A. and Bowles, N. E. (2002): The failing heart. *Nature* 415, 227-233
- Tsien, R. W. and Wheeler, D. B.: Voltage gated calcium channels. Calcium as a Cellular Regulator. Carafoli, E. and Klee, C. eds. Oxford Univ. Press, New York, 1999
- Uemura, A. et al. (1998): Demonstration of a Ca^{2+} /calmodulin-dependent protein kinase cascade in the hog heart. *Biochem. Biophys. Res. Commun.* 249, 355-360
- Ueyama, T. et al. (1999): Endothelin-1 activates p38 mitogen-activated protein kinase via endothelin-A receptor in rat myocardial cells. *Mol. Cell. Biochem.* 199, 119-24
- Van der Laarse, A. R. C. et al. (1998): Molecular aspects of cardiac hypertrophy and heart failure. *Cardiologie*
- Walker, B. R. et al. (1988): Direct cardiac effects of vasopressin: role of V_1 - and V_2 -vasopressinergic receptors. *Am. J. Physiol.* 255, H261-H265
- Wange, R. L. et al. (1991): Photoaffinity labeling of two rat liver plasma membrane proteins with [^{32}P] gamma-azidoanilido GTP in response to vasopressin. *J. Biol. Chem.* 266, 11409-11412
- Weeds, A. G. and Pope, B. (1971): Substructure of the myosin molecule. *Nature* 234, 85-88
- Wegner, M. et al. (1992): Calcium regulated phosphorylation within the leucine zipper of C/EBP β . *Science* 256, 370-373
- Welsh, C. et al. (1990): Vasopressin stimulates phospholipase D activity against phosphatidylcholine in vascular smooth muscle cells. *Lipids* 25, 675-684
- Wendell S. A. et al. (2000): Renin-angiotensin system and sympathetic nervous system in cardiac pressure-overload hypertrophy. *Am. J. Physiol. Heart Circ. Physiol.* 279, H2797-H2806
- Wynne, J. and Braunwald, E.: The cardiomyopathies and myocarditides. In: Braunwald, E., Zipes, DP., Libby, P., eds. *Heart Disease: A Textbook of cardiovascular Medicine*. 6th edn, W.B. Saunders, Philadelphia, Pennsylvania, 2001, 1751-1806
- Xu, A. et al. (1993): Phosphorylation and activation of the Ca^{2+} -pumping ATPase of cardiac sarcoplasmic reticulum by Ca^{2+} /calmodulin-dependent protein kinase. *J. Biol. Chem.* 268, 8394-8397
- Xu, Y.-J. and Gopalakrishnan, V. (1991): Vasopressin increases cytosolic free (Ca^{2+}) in the neonatal rat cardiomyocyte. Evidence for V_1 subtype receptors. *Circ. Res.* 69, 239-245

- Xu, Y-J. et al. (1999): Vasopressin accelerates protein synthesis in neonatal rat cardiomyocytes. *Mol. Cell. Biochem.* 195, 183-190
- Yaffe, D. (1968): Retention of differentiation potentialities during prolonged cultivation of myogenic cells. *Proc. Natl. Acad. Sci. USA* 61, 477-483
- Yamazaki, T. et al. (1998): Role of ion channels and exchangers in mechanical stretch-induced cardiomyocyte hypertrophy. *Circ. Res.* 82, 430-37
- Yang, J. et al. (2000): Independent signals control expression of the calcineurin inhibitory proteins MCIP1 and MCIP2 in striated muscles. *Circ. Res.* 87, E61-68
- Yang, T. T. et al. (2002): Phosphorylation of NFATc4 by p38 mitogen-activated protein kinases. *Mol. Cell. Biol.* 22, 3892-3904
- Yano, M. et al. (1998): Differential activation of cardiac c-jun amino-terminal kinase and extracellular signal-regulated kinase in angiotensin II-mediated hypertension. *Circ. Res.* 83, 752-60
- Yokokura, H. et al. (1997): Isolation and comparison of rat cDNA encoding Ca^{2+} /calmodulin-dependent protein kinase I isoforms. *Biochim. Biophys. Acta* 1338, 8-12
- Zacharzowsky, U. B. et al. (2002): Analysis of the Energetic State of Heart Cells After Adenovirus-Mediated Expression of hALC-1. *J. Cell. Biochem.* 86, 422-431
- Zhang, C. L. et al. (2002): Class II Histone Deacetylases Act as Signal-Responsive Repressors of Cardiac Hypertrophy. *Cell* 110, 479-488
- Zhang, T. et al. (2003): The δ_C -isoform of CaMKII is activated in cardiac hypertrophy and induces dilated cardiomyopathy and heart failure. *Circ. Res.* 92, 912-919
- Zhang, T. and Brown, J. H. (2004): Role of Ca^{2+} /calmodulin-dependent protein kinase II in cardiac hypertrophy and heart failure. *Cardiovasc. Res.* 63, 476-486
- Zhu, W. et al. (2000): Ca^{2+} /calmodulin-dependent kinase II and calcineurin play critical roles in endothelin-1-induced cardiomyocyte hypertrophy. *J. Biol. Chem.* 275, 15239-15245
- Zimmermann, K. et al. (1990): Heterogenic mRNAs with an Identical Protein-coding Region of the Human Embryonic Myosin Alkali Light Chain in Skeletal Muscle Cells. *J. Mol. Biol.* 211, 505-513