

Literaturverzeichnis

Adam CL, Kyle CE, Young P (1996). Seasonal patterns of growth, voluntary food intake and plasma concentrations of prolactin, insulin-like growth factor-1, LH and gonadal steroids in male and female pre-pubertal red deer (*Cervus elaphus*) reared in either natural photoperiod or constant daylength. Animal Science 62: 605 - 613.

Adam CL, Kyle CE, Young P, Atkinson T (1995). Effect of Nutritional Growth Restriction on Timing of Reproductive Development and Plasma-Concentrations of Insulin-Like Growth-Factor-I and Growth-Hormone in Male Red Deer (*Cervus-Elaphus*) Reared in Constant Photoperiod. Animal Science 61: 155 - 160.

Adham IM, Burkhardt E, Benahmed M, Engel W (1993). Cloning of a cDNA for a novel insulin-like peptide of the testicular Leydig cells. J.Biol.Chem. 268: 26668 - 26672.

Bardin CW, Yan Cheng C, Mustow NA, Gunsalus GL (1994). The Sertoli Cell. In: The Physiology of Reproduction (Hrsg.: Knobil E, Neill JD), 1291-1334. Raven Press, New York.

Bartke A, Sinha Hikim AP, Russell LD (1996). Leydig cell structure and function in seasonal breeders. In: The Leydig Cell (Hrsg.: Payne AH, Hardy MP, Russell LD), 431-450. Cache River Press, Vienna, Illinois.

Benahmed M, Morera AM, Chauvin MC, de Peretti E (1987). Somatomedin C/insulin-like growth factor 1 as a possible intratesticular regulator of Leydig cell activity. Mol.Cell Endocrinol. 50: 69 - 77.

Benton L, Shan LX, Hardy MP (1995). Differentiation of adult Leydig cells. J.Steroid Biochem.Mol.Biol. 53: 61 - 68.

Berndston WE, Desjardins C (1974). The cycle of the seminiferous epithelium and spermatogenesis in the bovine testis. Am.J.Anat. 140: 167 - 179.

Blottner S, Roelants H (1998). Calculation of spermatogenic transformations based on dual-flow cytometric analysis of testicular tissue in seasonal breeders. Andrologia 30: 331 - 337.

Blottner S, Roelants H, Wagener A, Wenzel UD (1999). Testicular mitosis, meiosis and apoptosis in mink (*Mustela vison*) during breeding and non-breeding seasons. Anim Reprod.Sci. 57: 237 - 249.

Böhme G, Pier C (1986). Ordnungskriterien für den Zyklus des Epithels der Tubuli seminiferi, nach Untersuchungen beim Kater (*Felis catus*). Berl.Münch.Tierärztl.Wschr. 99: 232 - 236.

Brodowski AK, Jewgenow K, Pielowski Z, Blottner S (2001). Saisonale Veränderungen histologisch- morphometrischer Parameter im Testis von Feldhasen. Zeit.Jagdwiss. 47: 26 - 33.

Bronson FH, Heidemann PD (1994). Seasonal Regulation of Reproduction in Mammals. In: The Physiology of Reproduction (Hrsg.: Knobil E, Neill JD), 541-584. Raven Press, New York.

- Bursch W, Oberhammer F, Schulte-Hermann R (1992). Cell death by apoptosis and its protective role against disease. *Trends Pharmacol.Sci.* 13: 245 - 251.
- Cancilla B, Davies A, Ford-Perriss M, Risbridger GP (2000). Discrete cell- and stage-specific localisation of fibroblast growth factors and receptor expression during testis development. *J.Endocrinol.* 164: 149 - 159.
- Carpenter G, Wahl MI (1990). The epidermal growth factor family. In: *Peptide growth factors and their receptors* (Hrsg.: Spron MB, Roberts AB), Springer Verlag,
- Caussanel V, Tabone E, Hendrick JC, Dacheux F, Benahmed M (1997). Cellular distribution of transforming growth factor betas 1, 2, and 3 and their types I and II receptors during postnatal development and spermatogenesis in the boar testis. *Biol.Reprod.* 56: 357 - 367.
- Caussanel V, Tabone E, Mauduit C, Dacheux F, Benahmed M (1996). Cellular distribution of EGF, TGFalpha and their receptor during postnatal development and spermatogenesis of the boar testis. *Mol.Cell Endocrinol.* 123: 61 - 69.
- Clermont Y (1972). Kinetics of spermatogenesis in mammals: seminiferous epithelium cycle and spermatogonial renewal. *Physiol Rev.* 52: 198 - 236.
- Clermont Y, Leblond CP (1955). Spermiogenesis of man, monkey, ram and other mammals as shown by the "periodic acid- schiff" technique. *Am.J.Anat.* 96: 229 - 254.
- D'Occhio MJ, Aspden WJ (1999). Endocrine and reproductive responses of male and female cattle to agonists of gonadotrophin-releasing hormone. *J.Reprod.Fertil.Suppl* 54: 101 - 114.
- Daughaday WH, Rotwein P (1989). Insulin-like growth factors I and II. Peptide, messenger ribonucleic acid and gene structures, serum, and tissue concentrations. *Endocr.Rev.* 10: 68 - 91.
- de Kretser DM, Kerr JB (1994). The Cytology of the Testis. In: *The Physiology of Reproduction* (Hrsg.: Knobil E, Neill JD), 1177-1290. Raven Press, New York.
- de Kretser DM, Loveland KL, Meinhardt A, Simorangkir D, Wreford N (1998). Spermatogenesis. *Hum.Reprod.* 13 Suppl 1: 1 - 8.
- Desjardins C, Ewing LL (1993). Leydig Cells: Structure and Function. In: *Cell and Molecular Biology of the Testis* (Hrsg.: Desjardins C, Ewing LL), 137-165. Oxford University Press, Oxford.
- Di Guglielmo GM, Le Roy C, Goodfellow AF, Wrana JL (2003). Distinct endocytic pathways regulate TGF-beta receptor signalling and turnover. *Nat.Cell Biol.* 5: 410 - 421.
- Dissel-Emiliani FM, Boer-Brouwer M, De Rooij DG (1996). Effect of fibroblast growth factor-2 on Sertoli cells and gonocytes in coculture during the perinatal period. *Endocrinology* 137: 647 - 654.
- Ergun S, Kilic N, Fiedler W, Mukhopadhyay AK (1997). Vascular endothelial growth factor and its receptors in normal human testicular tissue. *Mol.Cell Endocrinol.* 131: 9 - 20.

Fawcett DW (1973). Observations on the organization of the interstitial tissue of the testis and on the occluding cell junctions in the seminiferous epithelium. *Adv.Biosci.* 10: 83 - 99.

Flores JM, Sanchez MA, Gonzalez M, Pizarro M (1998). Caprine testicular hypoplasia associated with sexual reversion decreases the expression of insulin-like growth factor II (IGF-II) mRNA in testes. *Anim Reprod.Sci.* 52: 279 - 288.

Franca LR, Godinho CL (2003). Testis Morphometry, Seminiferous Epithelium Cycle Length, and Daily Sperm Production in Domestic Cats (*Felis catus*). *Biol.Reprod.* 68: 1554 - 1561.

Franck Lissbrant I, Lissbrant E, Persson A, Damber JE, Bergh A (2003). Endothelial cell proliferation in male reproductive organs of adult rat is high and regulated by testicular factors. *Biol.Reprod.* 68: 1107 - 1111.

Gluckman PD, Ambler GR (1993). What is the function of circulating insulin-like growth factor-2 in postnatal life? *Mol.Cell Endocrinol.* 92: C1 - C3.

Godin I, Wylie CC (1991). TGF beta 1 inhibits proliferation and has a chemotropic effect on mouse primordial germ cells in culture. *Development* 113: 1451 - 1457.

Goeritz F, Quest M, Wagener A, Fassbender M, Broich A, Hildebrandt TB, Hofmann RR, Blottner S (2003). Seasonal timing of sperm production in roe deer: interrelationship among changes in ejaculate parameters, morphology and function of testis and accessory glands. *Theriogenology* 59: 1487 - 1502.

Gospodarowicz D, Ferrara N, Schweigerer L, Neufeld G (1987). Structural characterization and biological functions of fibroblast growth factor. *Endocr.Rev.* 8: 95 - 114.

Hagenas L, Ritzen EM, Plooen L, Hansson V, French FS, Nayfeh SN (1975). Sertoli cell origin of testicular androgen-binding protein (ABP). *Mol.Cell Endocrinol.* 2: 339 - 350.

Hardy MP, Mendis-Handagama SM, Zirkin BR, Ewing LL (1987). Photoperiodic variation of Leydig cell numbers in the testis of the golden hamster: a possible mechanism for their renewal during recrudescence. *J.Exp.Zool.* 244: 269 - 276.

Hartung J, Schoppmeyer K (1986). Zur zyklischen Entwicklung der Hoden beim Reh. Beiträge zur Jagd- und Wildforschung 14: 214 - 221.

Hartung J, Schoppmeyer K (1994). Beitrag zu Hodenveränderungen beim Rehwild. Beiträge zur Jagd- und Wildforschung 19: 117 - 120.

Harvey MB, Kaye PL (1992). IGF-2 stimulates growth and metabolism of early mouse embryos. *Mech.Dev.* 38: 169 - 173.

Hikim AP, Bartke AJ, Russell LD (1988). The seasonal breeding hamster as a model to study structure-function relationships in the testis. *Tissue Cell* 20: 63 - 78.

Hikim AP, Hikim IS, Amador AG, Bartke A, Woolf A, Russell LD (1991). Reinitiation of spermatogenesis by exogenous gonadotropins in a seasonal breeder, the woodchuck (*Marmota monax*), during gonadal inactivity. *Am.J.Anat.* 192: 194 - 213.

Hochereau-de Reviers MT, Lincoln GA (1978). Seasonal variation in the histology of the testis of the red deer, *Cervus elaphus*. *J.Reprod.Fertil.* 54: 209 - 213.

Hoffmann B, Barth D, Karg H (1978). Progesterone and estrogen levels in peripheral plasma of the pregnant and nonpregnant roe deer (*Capreolus capreolus*). *Biol.Reprod.* 19: 931 - 935.

Hombach-Klonisch S, Kauffold J, Rautenberg T, Steger K, Tetens F, Fischer B, Klonisch T (2000). Relaxin-like factor (RLF) mRNA expression in the fallow deer. *Mol.Cell Endocrinol.* 159: 147 - 158.

Hombach-Klonisch S, Tetens F, Kauffold J, Steger K, Fischer B, Klonisch T (1999). Molecular cloning and localization of caprine relaxin-like factor (RLF) mRNA within the goat testis. *Mol.Reprod.Dev.* 53: 135 - 141.

Ivell R, Balvers M, Domagalski R, Ungefroren H, Hunt N, Schulze W (1997). Relaxin-like factor: a highly specific and constitutive new marker for Leydig cells in the human testis. *Mol.Hum.Reprod.* 3: 459 - 466.

Ivell R, Bathgate RA (2002). Reproductive biology of the relaxin-like factor (RLF/INSL3). *Biol.Reprod.* 67: 699 - 705.

Jeyakumar M, Suresh R, Krishnamurthy HN, Moudgal NR (1995). Changes in testicular function following specific deprivation of LH in the adult male rabbit. *J.Endocrinol.* 147: 111 - 120.

Johnson L, Matt KS, Bartke A, Nguyen HB, Le HT (1987). Effect of photoperiod on the size of the Leydig cell population and the rate of recruitment of Leydig cells in adult Syrian hamsters. *Biol.Reprod.* 37: 727 - 738.

Kierszenbaum AL (1994). Mammalian spermatogenesis in vivo and in vitro: a partnership of spermatogenic and somatic cell lineages. *Endocr.Rev.* 15: 116 - 134.

Klonisch T, Ivell R, Balvers M, Kliesch S, Fischer B, Bergmann M, Steger K (1999). Expression of relaxin-like factor is down-regulated in human testicular Leydig cell neoplasia. *Mol.Hum.Reprod.* 5: 104 - 108.

Korpelainen EI, Karkkainen MJ, Tenhunen A, Lakso M, Rauvala H, Vierula M, Parvinen M, Alitalo K (1998). Overexpression of VEGF in testis and epididymis causes infertility in transgenic mice: evidence for nonendothelial targets for VEGF. *J.Cell Biol.* 143: 1705 - 1712.

Laemmli UK (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680 - 685.

Lahr G, Mayerhofer A, Seidl K, Bucher S, Grothe C, Knochel W, Gratzl M (1992). Basic fibroblast growth factor (bFGF) in rodent testis. Presence of bFGF mRNA and of a 30 kDa bFGF protein in pachytene spermatocytes. *FEBS Lett.* 302: 43 - 46.

Leblond CP, Clermont Y (1952a). Definition of the stages of the cycle of the seminiferous epithelium in the rat. *Ann.N.Y.Acad.Sci.* 55: 548 - 573.

Leblond CP, Clermont Y (1952b). Spermiogenesis of rat, mouse, hamster and guinea pig as revealed by the "periodic acid-fuchsin sulphurous acid" technique. *Am.J.Anat.* 90: 167 - 206.

- Lejeune H, Habert R, Saez JM (1998). Origin, proliferation and differentiation of Leydig cells. *J.Mol.Endocrinol.* 20: 1 - 25.
- LeRoith D, Roberts CT, Jr. (1993). Insulin-like growth factors. *Ann.N.Y.Acad.Sci.* 692: 1 - 9.
- Lin T, Blaisdell J, Haskell JF (1987). Transforming growth factor-beta inhibits Leydig cell steroidogenesis in primary culture. *Biochem.Biophys.Res.Commun.* 146: 387 - 394.
- Lincoln GA (1989). Seasonal aspects in testicular physiology in seasonal mammals. In: *The Testis* (Hrsg.: Burger H, de Kretser DM), 229-385. Raven Press, New York.
- Marchlewska-Koj A, Kruczak M (1988). Seasonal variation in testicular activity of roe deer (*Capreolus capreolus*). *Folia Biol.(Krakow)* 36: 119 - 125.
- Markwald RR, Davis RW, Kainer RA (1971). Histological and histochemical periodicity of cervine Leydig cells in relation to antler growth. *Gen.Comp Endocrinol.* 16: 268 - 280.
- Massague J (2000). How cells read TGF-beta signals. *Nat.Rev.Mol.Cell Biol.* 1: 169 - 178.
- Mayerhofer A, Russell LD, Grothe C, Rudolf M, Gratzl M (1991). Presence and localization of a 30-kDa basic fibroblast growth factor-like protein in rodent testes. *Endocrinology* 129: 921 - 924.
- Mendis-Handagama SM (1991). Mitosis in normal adult guinea pig Leydig cells. *J.Androl* 12: 240 - 243.
- Mendis-Handagama SM, Ariyaratne HB (2001). Differentiation of the adult Leydig cell population in the postnatal testis. *Biol.Reprod.* 65: 660 - 671.
- Mendis-Handagama SM, Ewing LL (1990). Sources of error in the estimation of Leydig cell numbers in control and atrophied mammalian testes. *J.Microsc.* 159 (Pt 1): 73 - 82.
- Monfort SL, Brown JL, Bush M, Wood TC, Wemmer C, Vargas A, Williamson LR, Montali RJ, Wildt DE (1993a). Circannual inter-relationships among reproductive hormones, gross morphometry, behaviour, ejaculate characteristics and testicular histology in Eld's deer stags (*Cervus eldi thamin*). *J.Reprod.Fertil.* 98: 471 - 480.
- Monfort SL, Brown JL, Wood TC, Wemmer C, Vargas A, Williamson LR, Wildt DE (1993b). Seasonal patterns of basal and GnRH-induced LH, FSH and testosterone secretion in Eld's deer stags (*Cervus eldi thamin*). *J.Reprod.Fertil.* 98: 481 - 488.
- Morera AM, Cochet C, Keramidas M, Chauvin MA, de Peretti E, Benahmed M (1988). Direct regulating effects of transforming growth factor beta on the Leydig cell steroidogenesis in primary culture. *J.Steroid Biochem.* 30: 443 - 447.
- Mullaney BP, Skinner MK (1991). Growth factors as mediators of testicular cell-cell interactions. *Baillieres Clin.Endocrinol.Metab* 5: 771 - 790.
- Nakayama Y, Yamamoto T, Abe SI (1999). IGF-I, IGF-II and insulin promote differentiation of spermatogonia to primary spermatocytes in organ culture of newt testes. *Int.J.Dev.Biol.* 43: 343 - 347.

Nakazumi H, Sasano H, Maehara I, Orikasa S (1996). Transforming growth factor-alpha, epidermal growth factor, and epidermal growth factor receptor in human testis obtained from biopsy and castration: immunohistochemical study. *Tohoku J.Exp.Med.* 178: 381 - 388.

Nalbandian A, Dettin L, Dym M, Ravindranath N (2003). Expression of vascular endothelial growth factor receptors during male germ cell differentiation in the mouse. *Biol.Reprod.* 69: 985 - 994.

Neaves WB (1973). Changes in testicular Leydig cells and in plasma testosterone levels among seasonally breeding rock hyrax. *Biol.Reprod.* 8: 451 - 466.

Nieschlag E, Behre HM (1996). *Andrologie*. (Hrsg.: Nieschlag E, Behre HM) Springer, Berlin.

Nilsen- Hamilton M (1990). Transforming growth factor beta and its actions on cellular growth and differentiation. In: *Growth factors and development* (Hrsg.: Nilsen- Hamilton M), 96-136. Academic Press, Inc., San Diego.

Nishimura T, Utsunomiya Y, Hoshikawa M, Ohuchi H, Itoh N (1999). Structure and expression of a novel human FGF, FGF-19, expressed in the fetal brain. *Biochim.Biophys.Acta* 1444: 148 - 151.

O'Donnell L, McLachlan RI, Wreford NG, Robertson DM (1994). Testosterone promotes the conversion of round spermatids between stages VII and VIII of the rat spermatogenic cycle. *Endocrinology* 135: 2608 - 2614.

Ortavant R, Courot M (1977). Spermatogenesis in domestic animals. In: *Reproduction in domestic animals* (Hrsg.: Cole HT, Cupps PT), 202-224. Academic press, New York.

Pan X, Cao GW, Ke CW, He X, Qi ZT (1999). Structure and Function of Vascular Endothelial Growth Factor and Its Related Proteins. *Sheng Wu Hua Xue Yu Sheng Wu Wu Li Xue Bao.(Shanghai)* 31: 357 - 361.

Pelletier RM (1986). Cyclic formation and decay of the blood-testis barrier in the mink (*Mustela vison*), a seasonal breeder. *Am.J.Anat.* 175: 91 - 117.

Pimentel E (1994a). *Handbook of Growth Factors Vol.1: General Basic Aspects*. CRC Press, Boca Raton.

Pimentel E (1994b). *Handbook of Growth Factors Vol.2: Peptide Growth factors*. CRC Press, Boca Raton.

Reyes E, Bubenik GA, Lobos A, Schams D, Bartos L (1997). Seasonal levels of cortisol, IGF-1 and triiodothyronine in adult male pudu (*Pudu puda*). *Folia Zoologica* 46: 109 - 116.

Roelants H, Schneider F, Goritz F, Streich J, Blottner S (2002). Seasonal Changes of Spermatogonial Proliferation in Roe Deer, Demonstrated by Flow Cytometric Analysis of c-kit Receptor, in Relation to Follicle-Stimulating Hormone, Luteinizing Hormone, and Testosterone. *Biol.Reprod.* 66: 305 - 312.

Rotwein P, Bichell DP, Kikuchi K (1993). Multifactorial regulation of IGF-I gene expression. *Mol.Reprod.Dev.* 35: 358 - 363.

Russell LD, Ettlin R, Sinha Hikim AP, Clegg E (1990a). Evaluation of the Testis. Cache River Press, Vienna, Illinois.

Russell LD, Ren HP, Sinha H, I, Schulze W, Sinha Hikim AP (1990b). A comparative study in twelve mammalian species of volume densities, volumes, and numerical densities of selected testis components, emphasizing those related to the Sertoli cell. Am.J.Anat. 188: 21 - 30.

Schlatt S, Meinhardt A, Nieschlag E (1997). Paracrine regulation of cellular interactions in the testis: factors in search of a function. Eur.J.Endocrinol. 137: 107 - 117.

Schulze C (1992). Interaktionen zwischen Leydigzellen - peritubulären Zellen - Sertolizellen und ihre Bedeutung für die Steuerung der Spermatogenese. Ann.Anat. 509 - 515.

Sempere AJ, Mauget R, Mauget C (1998). Reproductive physiology of roe deer. In: The European Roe Deer: The Biology of Success (Hrsg.: Andersen R, Duncan P, Linnel DC), 161-188. Scandinavian University Press,

Sharpe RM (1994). Regulation of Spermatogenesis. In: The Physiology of Reproduction (Hrsg.: Knobil E, Neill JD), 1363-1434. Raven Press, New York.

Sharpe RM, Maddocks S, Kerr JB (1990). Cell-cell interactions in the control of spermatogenesis as studied using Leydig cell destruction and testosterone replacement. Am.J.Anat. 188: 3 - 20.

Sharpe RM, Maddocks S, Millar M, Kerr JB, Saunders PT, McKinnell C (1992). Testosterone and spermatogenesis. Identification of stage-specific, androgen-regulated proteins secreted by adult rat seminiferous tubules. J.Androl 13: 172 - 184.

Short RV, Mann T (1966). The sexual cycle of a seasonally breeding mammal, the roebuck (*Capreolus capreolus*). J.Reprod.Fertil. 12: 337 - 351.

Sinha Hikim AP, Bartke A, Russell LD (1988). Morphometric studies on hamster testes in gonadally active and inactive states: light microscope findings. Biol.Reprod. 39: 1225 - 1237.

Skinner MK (1987). Cell-cell interactions in the testis. Ann.N.Y.Acad.Sci. 513: 158 - 171.

Skinner MK (1991). Cell-cell interactions in the testis. Endocr.Rev. 12: 45 - 77.

Skinner MK, Norton JN, Mullaney BP, Rosselli M, Whaley PD, Anthony CT (1991). Cell-cell interactions and the regulation of testis function. Ann.N.Y.Acad.Sci. 637: 354 - 363.

Skinner MK, Takacs K, Coffey RJ (1989). Transforming growth factor-alpha gene expression and action in the seminiferous tubule: peritubular cell-Sertoli cell interactions. Endocrinology 124: 845 - 854.

Sluka P, O'Donnell L, Stanton PG (2002). Stage-specific expression of genes associated with rat spermatogenesis: characterization by laser-capture microdissection and real-time polymerase chain reaction. Biol.Reprod. 67: 820 - 828.

Smith PK, Krohn RI, Hermanson GT, Mallia AK, Gartner FH, Provenzano MD, Fujimoto EK, Goeke NM, Olson BJ, Klenk DC (1985). Measurement of protein using bicinchoninic acid. Anal.Biochem. 150: 76 - 85.

- Soder O, Bang P, Wahab A, Parvinen M (1992). Insulin-like growth factors selectively stimulate spermatogonial, but not meiotic, deoxyribonucleic acid synthesis during rat spermatogenesis. *Endocrinology* 131: 2344 - 2350.
- Spiteri-Grech J, Nieschlag E (1993). Paracrine factors relevant to the regulation of spermatogenesis--a review. *J.Reprod.Fertil.* 98: 1 - 14.
- Steger K (2001). Spermatogenese und Spermatogenesestörungen. *Reproduktionsmedizin* 17: 137 - 148.
- Strbenc M, Bavdek SV (2001). Apoptosis and proliferation in the testes of the brown hare (*Lepus europaeus*) under the influence of photoperiod. *Pflugers Arch.* 442: R161 - R162.
- Sundqvist C, Ellis LC, Bartke A (1988). Reproductive endocrinology of the mink (*Mustela vison*). *Endocr.Rev.* 9: 247 - 266.
- Suresh R, Medhamurthy R, Moudgal NR (1995). Comparative studies on the effects of specific immunoneutralization of endogenous FSH or LH on testicular germ cell transformations in the adult bonnet monkey (*Macaca radiata*). *Am.J.Reprod.Immunol.* 34: 35 - 43.
- Suttie JM, Fennessy PF, Corson ID, Laas FJ, Crosbie SF, Butler JH, Gluckman PD (1989). Pulsatile growth hormone, insulin-like growth factors and antler development in red deer (*Cervus elaphus scoticus*) stags. *J.Endocrinol.* 121: 351 - 360.
- Suttie JM, White RG, Breier BH, Gluckman PD (1991). Photoperiod associated changes in insulin-like growth factor-I in reindeer. *Endocrinology* 129: 679 - 682.
- Tahka KM, Ruokonen A, Wallgren H, Teravainen T (1983). Temporal changes in testicular histology and steroidogenesis in juvenile bank voles (*Clethrionomys glareolus*, Schreber) subjected to different photoperiods. *Endocrinology* 112: 1420 - 1426.
- Teerds KJ, De Rooij DG, Rommerts FF, van dT, I, Wensing CJ (1989). Turnover time of Leydig cells and other interstitial cells in testes of adult rats. *Arch.Androl* 23: 105 - 111.
- ten Dijke P, Hansen P, Iwata KK, Pieler C, Foulkes JG (1988). Identification of another member of the transforming growth factor type beta gene family. *Proc.Natl.Acad.Sci.U.S.A* 85: 4715 - 4719.
- Tsutsumi O, Kurachi H, Oka T (1986). A physiological role of epidermal growth factor in male reproductive function. *Science* 233: 975 - 977.
- Vannelli BG, Barni T, Orlando C, Natali A, Serio M, Balboni GC (1988). Insulin-like growth factor-I (IGF-I) and IGF-I receptor in human testis: an immunohistochemical study. *Fertil.Steril.* 49: 666 - 669.
- Vasilatos-Younken R, Dunnington EA, Siegel PB, McMurtry JP (1997). Tissue-specific alterations in insulin-like growth factor-I concentrations in response to 3,3',5-triiodo-L-thyronine supplementation in the growth hormone receptor-deficient sex-linked dwarf chicken. *Gen.Comp Endocrinol.* 105: 31 - 39.

Verhoeven G (1992). Local control systems within the testis. Baillieres Clin.Endocrinol.Metab 6: 313 - 333.

Wagener A (2002). Saisonale Expression testikulärer Wachstumsfaktoren beim Reh. Dissertation, Mathematisch-Naturwissenschaftliche Fakultät der Universität Potsdam.

Wagener A, Blottner S, Goritz F, Fickel J (2000). Detection of growth factors in the testis of roe deer (*Capreolus capreolus*). Anim Reprod.Sci. 64: 65 - 75.

Webster JR, Corson ID, Littlejohn RP, Stuart SK, Suttie JM (1996). Effects of season and nutrition on growth hormone and insulin-like growth factor-I in male red deer. Endocrinology 137: 698 - 704.

Wrobel KH, Kessler M, Schimmel M (1993). Quantitative Evaluations of the Tubular Epithelium in the Testis of the Fallow Deer (*Dama-Dama*). Reproduction in Domestic Animals 28: 1 - 13.

Yan YC, Sun YP, Zhang ML (1998). Testis epidermal growth factor and spermatogenesis. Arch.Androl 40: 133 - 146.

Young KA, Nelson RJ (2000). Short photoperiods reduce vascular endothelial growth factor in the testes of *Peromyscus leucopus*. Am.J.Physiol Regul.Integr.Comp Physiol 279: R1132 - R1137.

Young KA, Zirkin BR, Nelson RJ (1999). Short photoperiods evoke testicular apoptosis in white-footed mice (*Peromyscus leucopus*). Endocrinology 140: 3133 - 3139.

Yu Z, Guo R, Ge Y, Ma J, Guan J, Li S, Sun X, Xue S, Han D (2003). Gene expression profiles in different stages of mouse spermatogenic cells during spermatogenesis. Biol.Reprod. 69: 37 - 47.

Zayed AE, Hifny A, Abou-Elmagd A, Wrobel KH (1995). Seasonal changes in the intertubular tissue of the camel testis (*Camelus dromedarius*). Anat.Anz. 177: 199 - 212.

Zhang ML, Yan YC, Sun YP, Koide SS (1997). Identification and expression of epidermal growth factor gene in mouse testis. Cell Res. 7: 51 - 59.

Zheng WX, Butwell TJ, Heckert L, Griswold MD, Bellve AR (1990). Pleiotropic actions of the seminiferous growth factor on two testicular cell lines: comparisons with acidic and basic fibroblast growth factors. Growth Factors 3: 73 - 82.

Zirkin BR (1993). Regulation of spermatogenesis in the adult mammal: gonadotropins and androgens. In: Cell and Molecular Biology of the Testis (Hrsg.: Desjardins C, Ewing LL), 166-188. Oxford University Press, New York.