

## 6.0 Abkürzungsverzeichnis

2-DE	Zweidimensionale Gelelektrophorese
5'-RACE	5'-Rapid amplification of cDNA ends
ABTS	2,2-Azino-bis (3-Ethylbenzthiazolin-6-sulfonat)
AIF	Apoptosis inducing factor
APIT	Aplysia punctata ink toxin
bp	Basenpaare
cDNA	Complementary DNA
CHX	Cycloheximid
DBM	Dinukleotidbindendes Motiv
DiTriSec	Di-Tri-Sticky end cloning
EAS	Essentielle Aminosäuren
ESI-MS	Electrospray ionisation mass spectrometry
FAD	Flavinadenindinukleotid
FCS	Fetal calf serum
FMN	Flavinmononukleotid
GOX	Glukoseoxidase
H <sub>2</sub> O <sub>2</sub>	Hydrogenperoxid
HPLC	High performance liquid chromatography
HRP	Meerrettichperoxidase
kB	Kilobasen
kD	Kilodalton
KRG	Krebs-Ringer
LAAO	L-Aminosäurenoxidase
LD	Letale Dosis
LOX	Lysinoxidase
MALDI	Matrix assisted laser desorption/ionization
MBTH	3-Methyl-2-Benzothiazolonhydrazon Hydrochlorid
MPIIB	Max-Planck-Institut für Infektionsbiologie
MPO	Myeloperoxidase
NAC	N-acetylcystein
NEAS	Nichtessentielle Aminosäuren
PBS	Phosphatbuffered saline

---

PCR	Polychainreaction
pI	Isoelektrischer Punkt
PI	Propidiumiodid
PLA <sub>2</sub>	Phospholipase A <sub>2</sub>
PRX	Peroxiredoxin
RLU	Relative light units
ROS	Reaktive Oxygenspezien
RPMI	Roswell Park Memorial Institute
RT°C	Raumtemperatur
SDS-PAGE	Sulfododecylsulfat-Polyacrylamidgelektrophorese
TNF	Tumor necrosis factor
TUNEL	TdT-mediated dUTP nick-end labeling