

7. Literatur

- Afzelius A (1910) Verhandlungen der Dermatologischen Gesellschaft zu Stockholm. Sitzung vom 28. Oktober 1909. *Archives of Dermatology and Syphology* 101: 403-406
- Afzelius A (1921) Erythema chronicum migrans. *Acta Dermato-Venerologica* (Stockholm) 2: 120-125
- Alekseev AN, Dubinina HV, Antykova LP, Dzhivanyan TI, Rijpkema SGT, Verbeek-de Kruif N, Cinco M (1998) Tick-borne borrelioses pathogen identification in *Ixodes* ticks (Acarina, Ixodidae) collected in St. Petersburg and Kaliningrad baltic regions of Russia. *Journal of Medical Entomology* 35: 136-142
- Anderson JF, Magnarelli LA, Stafford KC (1990) Birdfeeding ticks transstadially transmit *Borrelia burgdorferi* that infect Syrian hamsters. *Journal of Wildlife Disease* 26:1-10
- Baranton G, Postic D, Saint Girons I, Boerlin P, Piffaretti JC, Assous M, Grimont PAD (1992) Delineation of *Borrelia burgdorferi* sensu stricto, *Borrelia garinii* sp. nov., and group VS461 associated with Lyme borreliosis. *International Journal of Systematic Bacteriology* 42: 378-383
- Baril C, Richaud C, Baranton G, Saint-Girons I (1989) Linear chromosome of *Borrelia burgdorferi*. *Research in Microbiology* 140: 507-516
- Basta J, Plch J, Hulinska D, Daniel M (1999) Incidence of *Borrelia garinii* and *Borrelia afzelii* in *Ixodes ricinus* ticks in an urban environment, Prague, Czech Republic, between 1995 and 1998. *European Journal of Clinical and Microbiology Infectious Diseases* 18: 515-517
- Battaly GR and Fish D (1993) Relative importance of bird species as hosts for immature *Ixodes dammini* (Acari: Ixodidae) in a suburban residential landscape of southern New York State. *Journal of Medical Entomology* 30: 740-747
- Buchwald A (1883) Ein Fall von diffuser idiopathischer Haut-Atrophie. *Archives of Dermatology and Syphology* 10: 553-556
- Burgdorfer W, Barbour AG, Hayes SF, Benach JL, Grunwaldt E, Davis JP (1982) Lyme disease - a tick-borne spirochetosis? *Science* 216: 1317-1319
- Calhoun JB. The ecology and sociology of the Norway rat. Bethesda, MD: United States Department of Health, Education and Welfare, 1963: 288, Publication Health Service publication no. 1008
- Canica MM, Nato F, du Merle L, Mazie JC, Baranton G, Postic D (1993) Monoclonal antibodies for identification of *Borrelia afzelii* sp. nov. associated with late cutaneous manifestation of Lyme borreliosis. *Scandinavian Journal of Infectious Diseases* 25: 441-448

- Cinco M, Padovan D, Murgia R, Poldini L, Frusteri L, Van den Pol I, Verbeek-De Kruif N, Rijpkema SGT, Maroli M (1998) Rate of infection of *Ixodes ricinus* ticks with *Borrelia burgdorferi* sensu stricto, *Borrelia garinii*, *Borrelia afzelii* and group VS116 in an endemic focus of Lyme disease in Italy. *European Journal of Clinical Microbiology and Infectious Diseases* 17: 90-94
- De Boer R, Hovius KE, Nohlmans MKE, Gray JS (1993) The woodmouse (*Apodemus sylvaticus*) as a reservoir of tick-transmitted spirochetes (*Borrelia burgdorferi*) in the Netherlands. *Zentralblatt für Bakteriologie* 279: 404-416
- De Michelis S, Sewell HS, Collares-Pereira M, Santos-Reis M, Schouls LM, Benes V, Holmes EC, Kurtenbach K (2000) Genetic diversity of *Borrelia burgdorferi* sensu lato in ticks from mainland Portugal. *Journal of Clinical Microbiology* 38: 2128-2133
- De Silva AM, Zeidner NS, Zhang Y, Dolan MC, Piesman J, Fikrig E (1999) Influence of Outer surface protein A Antibody on *Borrelia burgdorferi* within feeding ticks. *Infection and Immunity* 67:30-35
- Donahue JG, Piesman J, Spielman A (1987) Reservoir competence of white-footed mice for Lyme disease spirochetes. *American Journal of Tropical Medicine and Hygiene* 36: 92-96
- Eiffert H, Ohlenbusch A, Christen HJ, Thomssen R, Spielman A, Matuschka F-R (1995) Nondifferentiation between Lyme disease spirochetes from vector ticks and human cerebrospinal fluid. *The Journal of Infectious Diseases* 171: 476-479
- Escudero R, Barral M, Perez A, Vitutia MM, Garcia-Perez AL, Jimenez S, Sellek RE, Anda P (2000) Molecular and pathogenic characterization of *Borrelia burgdorferi* sensu lato isolates from Spain. *Journal of Clinical Microbiology* 38: 4026-4033
- Fikrig E, Telford III SR, Barthold SW, Kantor FS, Spielman A, Flavell RA (1992) Elimination of *Borrelia burgdorferi* from vector ticks feeding on OspA-immunized mice. *Proceedings of the National Academy of Sciences* 89: 5418-5421
- Fukunaga M, Sohnaka M (1992) Tandem repeat of the 23S and 5S ribosomal RNA genes in *Borrelia burgdorferi*; the etiological agent of Lyme disease. *Biochemical and Biophysical Research Communications* 183: 952-957
- Gern L, Siegenthaler M, Hu CM, Leuba-Garcia S, Humair PF, Moret J (1994) *Borrelia burgdorferi* in rodents (*Apodemus flavicollis* and *A. sylvaticus*): Duration and enhancement of infectivity for *Ixodes ricinus* ticks. *European Journal of Epidemiology* 10: 75-80
- Gern L, Estrada-Pena A, Frandsen F, Gray JS, Jaenson TGT, Jongejan F, Kahl O, Korenberg E, Mehl R, Nuttall PA (1998) European reservoir hosts of *Borrelia burgdorferi* sensu lato. *Zentralblatt für Bakteriologie* 287: 196-204
- Gern L, Hu CM, Kocianova E, Vyrosteckova V, Rehacek J (1999) Genetic diversity of *Borrelia burgdorferi* sensu lato isolates obtained from *Ixodes ricinus* ticks collected in

Slovakia. *European Journal of Epidemiology* 15: 665-669

Gray JS, Schönberg A, Postic D, Belfaiza J, Saint-Girons I (1996) First isolation and characterisation of *Borrelia garinii*, agent of Lyme borreliosis, from Irish ticks. *Irish Journal of Medical Sciences* 165:24-26

Gray JS, Robertson JN, Key S (2000) Limited role of rodents as reservoirs of *Borrelia burgdorferi* sensu lato in Ireland. *European Journal of Epidemiology* 16: 101-103

Hansen K, Hindersson P, Strandberg Pedersen N (1988) Measurement of antibodies to the *Borrelia burgdorferi* flagellum improves serodiagnosis in Lyme disease. *Journal of Clinical Microbiology* 26: 338-346

Hovius KE, Stark LAM, Bleumink-Pluym NMC, van de Pol I, Verbeek-de Kruif N, Rijpkema SGT, Schouls LM, Houwers DJ (1999) Presence and distribution of *Borrelia burgdorferi* sensu lato species in internal organs and skin of naturally infected symptomatic and asymptomatic dogs, as detected by polymerase chain reaction. *Veterinary Quarterly* 21:54-58

Howe TR, Mayer LW, Barbour AG (1985) A single recombinant plasmid expressing two major outer surface proteins of the Lyme disease spirochete. *Science* 227:645-646

Howe TR, Laquier W, Barbour AG (1986) Organization of genes encoding two outer surface membrane proteins of the Lyme disease agent *Borrelia burgdorferi* within a single transcriptional unit. *Infection and Immunity* 54: 207-212

Hu CM, Humair PF, Wallich R, Gern L (1997) *Apodemus* sp. rodents, reservoir hosts for *Borrelia afzelii* in an endemic area in Switzerland. *Zentralblatt für Bakteriologie* 285: 558-564

Hu CM, Wilske B, Fingerle V, Lobet Y, Gern L (2001) Transmission of *Borrelia garinii* OspA Serotype 4 to BALB/c mice by *Ixodes ricinus* ticks collected in the field. *Journal of Clinical Microbiology* 39: 1169-1171

Huegli D, Hu CM, Humair PF, Wilske B, Gern L (2002) *Apodemus* species mice are reservoir hosts of *Borrelia garinii* OspA serotype 4 in Switzerland. *Journal of Clinical Microbiology* 40: 4735-4737

Humair PF, Turrian N, Aeschlimann A, and Gern L (1993) *Borrelia burgdorferi* in a focus of Lyme borreliosis: epizootiologic contribution of small mammals. *Folia Parasitologica* 40: 65-70

Humair PF, Peter O, Wallich R, Gern L (1995) Strain variation of Lyme disease spirochetes isolated from *Ixodes ricinus* ticks and rodents collected in two endemic areas in Switzerland. *Journal of Medical Entomology* 32: 433-438

Humair PF, Postic D, Wallich R, Gern L (1998a) An avian reservoir (*Turdus merula*) of the Lyme borreliosis spirochetes. *Zentralblatt für Bakteriologie* 287: 521-538

Humair PF, Gern L (1998b) Relationship between *Borrelia burgdorferi* sensu lato species, red squirrels (*Sciurus vulgaris*) and *Ixodes ricinus* in enzootic areas in Switzerland. *Acta Tropica* 69: 213-227

Humair PF, Rais O, Gern L (1999) Transmission of *Borrelia afzelii* from *Apodemus* mice and *Clethrionomys* voles to *Ixodes ricinus* ticks: differential transmission pattern and overwintering maintenance. *Parasitology* 118: 33-42

Ishiguro F, Takada N, Nakata K, Yano Y, Suzuki H, Masuzawa T, Yanagihara Y (1996) Reservoir competence of the vole, *Clethrionomys rufocanus bedfordiae*, for *Borrelia garinii* or *Borrelia afzelii*. *Microbiology and Immunology* 40: 67-69

Jaenson TGT und Tälleklint L (1992) Incompetence of roe deer as reservoirs of the Lyme borreliosis spirochete. *Journal of Medical Entomology* 29: 813-817

Jaenson TGT und Tälleklint L (1996) Lyme borreliosis spirochetes in *Ixodes ricinus* (Acari: Ixodidae) and the varying hare on isolated islands in the Baltic Sea. *Journal of Medical Entomology* 33: 339-343

Janetzki C, Kahl O, Knülle W: The Mongolian gerbil, *Meriones unguiculatus*, as a laboratory reservoir for *Borrelia burgdorferi*. Proc. 1st Int. Conf. On Tick-Borne Pathogens at the Host-Vector Interface: An Agenda for research (UG Munderloh and TJ Kurtti, eds.). St. Paul, Minnesota, USA, 15-18 September (1992) 143

Johnson RC, Schmid GP, Hyde FW, Steigerwalt AG, Brenner DJ (1984) *Borrelia burgdorferi* sp. nov.: etiologic agent of Lyme disease. *International Journal of Systematic Bacteriology* 34: 496-497

Junttila J, Peltomaa M, Soini H, Marjamäki M, Viljanen MK (1999) Prevalence of *Borrelia burgdorferi* in *Ixodes ricinus* ticks in urban recreational areas of Helsinki. *Journal of Clinical Microbiology* 37: 1361-1365

Kahl O: The Mongolian gerbil, *Meriones unguiculatus*, as a laboratory host of immature ixodid ticks. Proc. 1st Int. Conf. On Tick-Borne Pathogens at the Host-Vector Interface: An Agenda for research (UG Munderloh and TJ Kurtti, eds.). St. Paul, Minnesota, USA, 15-18 September (1992) 300

Kahl O, Janetzki-Mittmann C, Gray JS, Jonas R, Stein J, de Boer R (1998) Risk of infection with *Borrelia burgdorferi* sensu lato for a host in relation to the duration of nymphal *Ixodes ricinus* feeding and the method of tick removal. *Zentralblatt für Bakteriologie* 287: 41-52

Kimsey RB und Spielman A (1990) Motility of Lyme disease spirochetes in fluids as viscous as the extracellular matrix. *The Journal of Infectious Diseases* 162: 1205-1208

Kirstein F, Rijpkema SGT, Molkenboer M, Gray JS (1997a) The distribution and prevalence of *B. burgdorferi* genomospecies in *Ixodes ricinus* ticks in Ireland. *European Journal of*

Epidemiology 13: 67-72

Kirstein F, Rijpkema SGT, Molkenboer M, Gray JS (1997b) Local variations in the distribution and prevalence of *Borrelia burgdorferi* sensu lato genomospecies in *Ixodes ricinus* ticks. *Applied and Environmental Microbiology* 63: 1102-1106

Kurtenbach K, Dizij A, Arndt S, Seitz HM, Margos G, Moter SE, Kramer MD, Wallich R, Schaible UE, Simon MM (1994) Differential immune responses to *Borrelia burgdorferi* in european wild rodent species influence spirochete transmission to *Ixodes ricinus* L. (Acari Ixodidae). *Infection and Immunity* 62: 5344-5352

Kurtenbach K, Sewell H-S, Ogden NH, Randolph SE, Nuttall PA (1998a) Serum complement sensitivity as a key factor in Lyme disease ecology. *Infection and Immunity* 66: 1248-1251

Kurtenbach K, Peacey M, Rijpkema SGT, Hoodless AN, Nuttall PA, and Randolph SE (1998b) Differential transmission of the genospecies of *Borrelia burgdorferi* sensu lato by game birds and small rodents in England. *Applied and Environmental Microbiology* 64: 1169-1174

Kurtenbach K, Carey D, Hoodless AN, Nuttall PA, and Randolph SE (1998c) Competence of pheasants as reservoirs for Lyme disease spirochetes. *Journal of Medical Entomology* 35:77-81

Kurtenbach K, De Michelis S, Etti S, Schäfer SM, Sewell HS, Brade V, Kraiczy P (2002) Host association of *Borrelia burgdorferi* sensu lato – the key role of host complement. *Trends in Microbiology* 10: 74-79

Lane RS (1990) Susceptibility of the western fence lizard (*Sceloporus occidentalis*) to the Lyme borreliosis spirochete (*Borrelia burgdorferi*). *American Journal of Tropical Medicine and Hygiene* 42: 75-82

Lane RS, Quistad GB (1998) Borreliacidal factor in the blood of the western fence lizard (*Sceloporus occidentalis*). *Journal of Parasitology* 84: 29-34

Le Fleche A, Postic D, Girardet K, Peter O, Baranton G (1997) Characterization of *Borrelia lusitaniae* sp. nov. by 16S ribosomal DNA sequence analysis. *International Journal of Systematic Bacteriology* 47: 921-925

Levine FF, Wilson ML, Spielman A (1985) Mice as reservoir of the Lyme disease spirochete. *American Journal of Tropical Medicine and Hygiene* 34: 355-360

Liebisch G, Sohns B, Bautsch W (1998) Detection and typing of *Borrelia burgdorferi* sensu lato in *Ixodes ricinus* ticks attached to human skin by PCR. *Journal of Clinical Microbiology* 36: 3355-3358

Masuzawa T, Iwaki A, Sato Y, Miyamoto K, Korenberg EI, Yanagihara Y (1997) Genetic diversity of *Borrelia burgdorferi* sensu lato isolated in far eastern Russia. *Microbiology and*
62

Immunology 41: 595-600

Mather TN, Telford III SR, MacLachlan AB, Spielman A (1989) Incompetence of catbirds as reservoirs for the Lyme disease spirochete (*Borrelia burgdorferi*). *Journal of Parasitology* 75:66-69

Matuschka F-R und Spielman A (1986) The emergence of Lyme disease in a changing environment in North America and Central Europe. *Experimental & Applied Acarology* 2: 337-353

Matuschka F-R, Lange R, Spielman A, Richter D, Fischer P (1990) Subadult *Ixodes ricinus* (Acari: Ixodidae) on rodents in Berlin, West Germany. *Journal of Medical Entomology* 27: 385-390

Matuschka F-R, Fischer P, Musgrave K, Richter D, Spielman A (1991) Hosts on which nymphal *Ixodes ricinus* most abundantly feed. *American Journal of Tropical Medicine and Hygiene* 44: 100-107

Matuschka F-R, Fischer P, Heiler M, Blümcke S, Spielman A (1992a) Stage-associated risk of transmission of the Lyme disease spirochete by European *Ixodes* ticks. *Parasitology Research* 78: 695-698

Matuschka F-R, Fischer P, Heiler M, Richter D, Spielman A (1992b) Capacity of European animals as reservoir hosts for the Lyme disease spirochete. *The Journal of Infectious Diseases* 165: 479-483

Matuschka F-R and Spielman A (1992c) Loss of Lyme disease spirochetes from *Ixodes ricinus* ticks feeding on European blackbirds. *Experimental Parasitology* 74: 151-158

Matuschka F-R, Heiler M, Eiffert H, Fischer P, Lotter H, Spielman A (1993) Diversionary role of hoofed game in the transmission of Lyme disease spirochetes. *American Journal of Tropical Medicine and Hygiene* 48: 693-699

Matuschka F-R, Eiffert H, Ohlenbusch A, Spielman A (1994a) Amplifying role of edible dormice in Lyme disease transmission in central Europe. *The Journal of Infectious Diseases* 179: 122-127

Matuschka F-R, Ohlenbusch A, Eiffert H, Richter D, Spielman A (1996a) Characteristics of Lyme disease spirochetes in archived European ticks. *The Journal of Infectious Diseases* 174:424-426

Matuschka F-R, Endepols S, Richter D, Ohlenbusch A, Eiffert H, Spielman A (1996b) Risk of urban Lyme disease enhanced by the presence of rats. *The Journal of Infectious Diseases* 174: 1108-1111

Matuschka F-R, Endepols S, Richter D, Spielman A (1997) Competence of urban rats as reservoir hosts for Lyme disease spirochetes. *Journal of Medical Entomology* 34: 489-493

Matuschka F-R, Allgöwer R, Spielman A, Richter D (1999) Characteristics of garden dormice that contribute to their capacity as reservoirs for Lyme disease spirochetes. *Applied and Environmental Microbiology* 65: 707-711

Matuschka F-R, Schinkel TW, Klug B, Spielman A, Richter D (2000) Relative incompetence of European rabbits for Lyme disease spirochaetes. *Parasitology* 121:297-302

Matuschka F-R and Richter D (2002) Mosquitoes and soft ticks cannot transmit Lyme disease spirochetes. *Parasitology Research* 88:283-284

Monin R, Gern L, Aeschlimann A (1989) A study of the different modes of transmission of *Borrelia burgdorferi* by *Ixodes ricinus*. In: Stanek G (Hrsg.) Lyme borreliosis II, Zentralblatt für Bakteriologie (Suppl.18), Gustav Fischer Verlag, Stuttgart, New York, S.14-20

Nakao M, Miyamoto K, Fukunaga M (1994) Lyme disease spirochetes in Japan: Enzootic transmission cycles in birds, rodents, and *Ixodes persulcatus* ticks. *The Journal of Infectious Diseases* 170: 878-882

Nakao M, Miyamoto K (1995) Mixed infection of different *Borrelia* species among *Apodemus speciosus* mice in Hokkaido, Japan. *Journal of Clinical Microbiology* 33: 490-492

Nohlmans LMKE, De Boer R, Van den Bogaard AEJM, Van Boven CPA (1995) Genotypic and phenotypic analysis of *Borrelia burgdorferi* isolates from the Netherlands. *Journal of Clinical Microbiology* 33: 119-125

O'Connell S, Granström M, Gray JS, Stanek G (1998) Epidemiology of European Lyme borreliosis. *Zentralblatt für Bakteriologie* 287: 229-240

Ohlenbusch A (1996a) Beiträge zur Diagnostik und Pathogenese der Lyme-Borreliose und zur Transmission des Erregers *Borrelia burgdorferi*. Dissertation, Universität Göttingen, Cuvillier Verlag Göttingen

Ohlenbusch A, Matuschka F-R, Richter D, Christen HJ, Thomssen R, Spielman A, Eiffert H (1996b) Etiology of the acrodermatitis chronica atrophicans lesion in Lyme disease. *The Journal of Infectious Diseases* 174: 421-423

Olsén B, Jaenson TGT, Noppa L, Bunikis J, Bergström S (1993) A Lyme borreliosis cycle in seabirds and *Ixodes uriae* ticks. *Nature* 362: 340-342

Olsén B, Jaenson TGT, Bergström S (1995a) Prevalence of *Borrelia burgdorferi* sensu lato-infected ticks on migrating birds. *Applied and Environmental Microbiology* 61: 3082-3087

Pichon B, Godfroid E, Hoyois B, Bollen A, Rodhain F, Perez-Eid C (1995) Simultaneous infection of *Ixodes ricinus* nymphs by two *Borrelia burgdorferi* sensu lato species: possible implications for clinical manifestations. *Emerging Infectious Diseases* 3: 89-90

- Picken RN, Cheng Y, Strle F, Picken MM (1996) Patient isolates of *Borrelia burgdorferi* sensu lato with genotypic and phenotypic similarities of strain 25015. *The Journal of Infectious Diseases* 174: 1112-1115
- Piesman J, Mather TN, Sinsky RJ, and Spielman A (1987) Duration of tick attachment and *Borrelia burgdorferi* transmission. *Journal of Clinical Microbiology* 25: 557-558
- Piesman J (1993) Dynamics of *Borrelia burgdorferi* transmission by nymphal *Ixodes dammini* ticks. *The Journal of Infectious Diseases* 167: 1082-1085
- Postic D, Edlinger C, Richaud C, Grimont F, Dufresne Y, Perolat P, Baranton G, Grimont PAD (1990) Two genomic species in *Borrelia burgdorferi*. *Research in Microbiology* 141: 465-475
- Postic D, Assous M, Grimont PAD, Baranton G (1994) Diversity of *Borrelia burgdorferi* sensu lato evidenced by restriction fragment length polymorphism of rrf (5S)-rrl (23S) intergenic spacer amplicons. *International Journal of Systematic Bacteriology* 44: 742-752
- Postic D, Korenberg E, Gorelova N, Kovalevski YV, Bellenger E, Baranton G (1997) *Borrelia burgdorferi* sensu lato in Russia and neighbouring countries: high incidence of mixed isolates. *Research in Microbiology* 148: 691-702
- Postic D, Ras NM, Lane RS, Hendson M, Baranton G (1998) Expanded diversity among California *Borrelia* isolates and description of *Borrelia bissettii* sp. nov. (formerly *Borrelia* group DN127). *Journal of Clinical Microbiology* 36: 3497-3504
- Ribeiro JMC, Mather TN, Piesman J, Spielman A (1987) Dissemination and salivary delivery of Lyme disease spirochetes in vector ticks (Acari: Ixodidae). *Journal of Medical Entomology* 24: 201-205
- Richter D, Endepols S, Ohlenbusch A, Eiffert H, Spielman A, Matuschka F-R (1999) Genospecies diversity of Lyme disease spirochetes in rodent reservoirs. *Emerging Infectious Diseases* 5: 291-296
- Richter D, Spielman A, Komar N, Matuschka F-R (2000) Competence of American robins as reservoir hosts for Lyme disease spirochetes. *Emerging Infectious Diseases* 6: 133-138
- Rijpkema SGT, Molkenboer MJCH, Schouls LM, Jongejan F, Schellekens JFP (1995) Simultaneous detection and genotyping of three genomic groups of *Borrelia burgdorferi* sensu lato in Dutch *Ixodes ricinus* ticks by characterization of the amplified intergenic spacer region between 5S and 23S rRNA genes. *Journal of Clinical Microbiology* 33: 3091-3095
- Rijpkema SGT, Golubic D, Molkenboer M, Verbeek-De Kruif N, Schellekens J (1996) Identification of four genomic groups of *Borrelia burgdorferi* sensu lato in *Ixodes ricinus* ticks collected in a Lyme borreliosis endemic region of northern Croatia. *Experimental & Applied Acarology* 20: 23-30

Rijpkema SGT, Tazelaar DJ, Molkenboer M, Noordhoek GT, Plantinga G, Schouls LM, Schellekens JFP (1997) Detection of *Borrelia afzelii*, *Borrelia burgdorferi* sensu stricto, *Borrelia garinii* and group VS 116 by PCR in skin biopsies of patients with erythema migrans and acrodermatitis chronica atrophicans. *Clinical Microbiology and Infection* 3: 109-116

Rosa Brunet L, Sellitto C, Spielman A, Telford III SR (1995) Antibody response of the mouse reservoir of *Borrelia burgdorferi* in nature. *Infection and Immunity* 63: 3030-3036

Saint-Girons I, Norris SJ, Göbel U, Meyer J, Walker EM, Zuerner R (1992) Genome structure of spirochetes. *Research in Microbiology* 143: 615-621

Saint-Girons I, Gern L, Gray JS, Guy EC, Korenberg E, Nuttall PA, Rijpkema SGT, Schönberg A, Stanek G, Postic D (1998) Identification of *Borrelia burgdorferi* sensu lato species in Europe. *Zentralblatt für Bakteriologie* 287: 190-195

Sato Y, Miyamoto K, Iwaki A, Masuzawa T, Yanagihara Y, Korenberg EI, Gorelova NB, Volkov VI, Ivanov LI, Liberova RN (1996) Prevalence of Lyme disease spirochetes in *Ixodes persulcatus* and wild rodents in far eastern Russia. *Applied and Environmental Microbiology* 62: 3887-3889

Schwan TG, Piesman J, Golde WT, Dolan MC, Rosa PA (1995) Induction of an outer surface protein on *Borrelia burgdorferi* during tick feeding. *Proceedings of the National Academy of Science* 92: 2909-2913

Schwan TG and Piesman J (2000) Temporal changes in Outer Surface Proteins A and C of the Lyme disease-associated spirochete, *Borrelia burgdorferi*, during the chain of infection in ticks and mice. *Journal of Clinical Microbiology* 38: 382-388

Schwartz JJ, Gazumyan A, Schwartz I (1992) rRNA gene organization in the Lyme disease spirochete, *Borrelia burgdorferi*. *Journal of Bacteriology* 174: 3757-3765

Shih CM, Spielman A (1993) Accelerated transmission of Lyme disease spirochetes by partially fed vector ticks. *Journal of Clinical Microbiology* 31: 2878-2881

Shih CM, Liu LP, Spielman A (1995) Differential spirochetal infectivities to vector ticks of mice chronically infected by the agent of Lyme disease. *Journal of Clinical Microbiology* 33: 3164-3168

Smith RP Jr, Rand PW, Lacombe EH, Telford III SR, Rich SM, Piesman J, Spielman A (1993) Norway rats as reservoir hosts for Lyme disease spirochetes on Monhegan island, Maine. *The Journal of Infectious Diseases* 168: 687-691

Stanczak J, Kubica-Biernat B, Racewicz M, Kruminis-Lozowska W, Kur J (2000) Detection of three genospecies of *Borrelia burgdorferi* sensu lato in *Ixodes ricinus* ticks collected in different regions of Poland. *International Journal of Medical Microbiology* 290: 559-566

- Steere AC, Malawista SE, Snyderman DR, Andiman WA (1976) A cluster of arthritis in children and adults in Lyme, Connecticut. *Arthritis and Rheumatism* 19: 824
- Steere AC, Malawista SE, Snyderman DR, Shope RE, Andiman WA, Ross MR, Steele FM (1977) Lyme arthritis, an epidemic of oligoarticular arthritis in children and adults in three Connecticut communities. *Arthritis and Rheumatism* 20: 7-17
- Steere AC, Grodzicki RL, Kornblatt AN, Craft JE, Barbour AG, Burgdorfer W, Schmid GP, Johnson E, Malawista SE (1983) The spirochetal etiology of Lyme disease. *New England Journal of Medicine* 308: 733-740
- Strle F, Cheng Y, Nelson JA, Picken MM, Bouseman JK, Picken RN (1995) Infection rate of *Ixodes ricinus* ticks with *Borrelia afzelii*, *Borrelia garinii*, and *Borrelia burgdorferi* sensu stricto in Slovenia. *European Journal of Clinical Microbiology and Infectious Diseases* 14: 994-1001
- Strle F, Picken RN, Cheng Y, Cimperman J, Maraspin V, Lotric-Furlan S, Ruzic-Sabljic E, Picken MM (1997) Clinical findings for patients with Lyme borreliosis causing by *Borrelia burgdorferi* sensu lato with genotypic and phenotypic similarities to strain 25015. *Clinical Infectious Diseases* 25: 273-280
- Stünzner D, Hubalek Z, Halouzka J, Postic D, Pierer K, Marth E (1998) Prevalence of *Borrelia burgdorferi* s.l. in *Ixodes ricinus* ticks from Styria (Austria) and species identification by PCR-RFLP analysis. *Zentralblatt für Bakteriologie* 288: 471-478
- Tälleklint L und Jaenson TGT (1993) Maintenance by hares of European *Borrelia burgdorferi* in ecosystems without rodents. *Journal of Medical Entomology* 30: 273-276
- Tälleklint L, Jaenson TGT, Mather T (1993) Seasonal variation in the capacity of the bank vole to infect larval ticks (Acari: Ixodidae) with the Lyme disease spirochete, *Borrelia burgdorferi*. *Journal of Medical Entomology* 30: 812-815
- Tälleklint L und Jaenson TGT (1994) Transmission of *Borrelia burgdorferi* s.l. from mammal reservoirs to the primary vector of Lyme borreliosis, *Ixodes ricinus* (Acari: Ixodidae), in Sweden. *Journal of Medical Entomology* 31: 880-886
- Telford III SR, Mather TN, Moore SI, Wilson ML, Spielman A (1988) Incompetence of deer as reservoirs of the Lyme disease spirochete. *American Journal of Tropical Medicine and Hygiene* 39: 105-109
- Wang G, Van Dam AP, Le Fleche A, Postic D, Peter O, Baranton G, De Boer R, Spanjaard L, Dankert J (1997) Genetic and phenotypic analysis of *Borrelia valaisiana* sp. nov. (*Borrelia* genomic groups VS116 and M19). *International Journal of Systematic Bacteriology* 47: 926-932

Will G, Jauris-Heipke S, Schwab E, Busch U, Rößler D, Soutschek D, Wilske B, and Preac-Mursic V (1995) Sequence analysis of ospA genes shows homogeneity within *Borrelia burgdorferi* sensu stricto and *Borrelia afzelii* strains but reveals major subgroups within *Borrelia garinii* species. *Medical Microbiology and Immunology* 184: 73-80

Wilske B, Preac-Mursic V, Göbel UB, Graf B, Jauris S, Soutschek E, Schwab E, and Zumstein G (1993) An OspA serotyping system for *Borrelia burgdorferi* based on reactivity with monoclonal antibodies and OspA sequence analysis. *Journal of Clinical Microbiology* 31: 340-350

Zore A, Petrovec M, Prosenc K, Trilar T, Ruzic-Sabljić E, Avsic-Zupanc T (1999) Infection of small mammals with *Borrelia burgdorferi* sensu lato in Slovenia as determined by polymerase chain reaction (PCR). *Wiener Klinische Wochenschrift* 111: 997-999

Zung JL, Lewengrub S, Rudzinska MA, Spielman A, Telford III SR, and Piesman J (1989) Fine structural evidence for the penetration of the Lyme disease spirochete *Borrelia burgdorferi* through the gut and salivary tissues of *Ixodes dammini*. *Canadian Journal of Zoology* 67: 1737-1748