

4. Verzeichnis aller Referenzen

Adarichev VA, Glant TT. Experimental spondyloarthropathies: animal models of ankylosing spondylitis. *Curr Rheumatol Rep.* 2006 Aug;8(4):267-74.

Aho K, Ahvonen P, Lassus A, Sievers K, Tilikainen A: HL-A antigen 27 and reactive arthritis. *Lancet* 1972, 2: 157.

Allen RL, O'Callaghan CA, McMichael AJ, Bowness P. Cutting edge: HLA-B27 can form a novel beta 2-microglobulin-free heavy chain homodimer structure. *J Immunol.* 1999; 162: 5045-8.

Althoff CE, **Appel H**, Rudwaleit M, Sieper J, Eshed I, Hamm B, Hermann KG. Whole-body magnetic resonance imaging as a new screening tool for both axial and peripheral manifestations in spondyloarthritis. Eingereicht bei *Ann Rheum Dis*.

Altman JD, Moss PAH, Goulder PJR, Barouch DH, McHeyzer-Williams MG, Bell JI, McMichael AJ, Davis MM: Phenotypic analysis of antigen-specific T cells. *Science* 1996, 274: 94-96.

Appel H Untersuchungen zur zellulären und humoralen Immunantwort auf das 19kD Protein von *Yersinia enterocolitica* O:3 bei Patienten mit Yersinien-getriggelter Reaktiver Arthritis. Dissertation, Freie Universität Berlin 1997.

Appel H, Mertz A, Distler A, Sieper J, Braun J: The 19kDa protein of *Yersinia enterocolitica* O3 is recognized on the cellular and humoral level by patients with *Yersinia*-induced arthritis. *J Rheumatol* 1999, 26: 1964-1969.

Appel H, Gauthier L, Pyrdol J, Wucherpfennig KW: Kinetics of T-cell receptor binding by bivalent HLA-DR/Peptide complexes that activate antigen-specific human T-cells. *J Biol Chem* 2000; 275:312-21.

Appel H, Shroff N, Gauthier L, Wucherpfennig KW: Anergy induction by dimeric T cell receptor ligands. *J Immunol* 2001; 166: 5279-85.

Appel H, Rudwaleit M, Wu P, Grolms M, Sieper J, Mertz A: Synovial T cell proliferation to the *Yersinia enterocolitica* 19 kDa antigen differentiates yersinia triggered reactive arthritis (ReA) from ReA triggered by other enterobacteria. *Ann Rheum Dis* 2002; 61: 566-7.

Appel H, Neure L, Kuhne M, Braun J, Rudwaleit M, Sieper J: An elevated level of IL-10- and TGF β - secreting T cells, B cells and macrophages in the synovial membrane of patients with reactive arthritis compared to rheumatoid arthritis. *Clin Rheumatol* 2004, 23: 435-440. (a)

Appel H, Kuon W, Kuhne M, Wu P, Kuhlmann S, Thiel A, Kollnberger S, Bowness P, Sieper J: HLA-B27 restricted, peptide specific CD8+ T cell response and the use of HLA-B27 tetramers in Chlamydia triggered reactive arthritis. *Arthritis Res Ther* 2004; 6: R521-34. (b)

Appel H, Kuon W, Kuhne M, Köhler D, Kuhlmann S, Kollnberger S, Bowness P, Sieper J: The solvent-inaccessible Cys-67 residue of HLA-B27 contributes to T cell recognition of HLA-B27/peptide complexes. *J Immunol.* 2004; 173: 6564-73. (c)

Appel H, Atagunduz P, Sieper J: Reply to: A cartilage-derived self peptide presented by HLA-B27 molecules? Comment on the article by Atagunduz et al. by - Ziegler A, Loll B,

Biesiadka J, Saenger W, Kellermann T, Misselwitz R, Uchanska-Ziegler B. *Arthritis Rheum.* 2005; 52: 2581-2; author reply 2582-3.

Appel H, Kuhne M, Spiekermann S, Köhler D, Zacher J, Stein H, Sieper J, Loddenkemper C. Immunohistochemical analysis of hip arthritis in ankylosing spondylitis – an analysis of the bone/cartilage interface and subchondral bone marrow. *Arthritis Rheum* 2006; 54:1805-13. (a)

Appel H, Kuhne M, Spiekermann S, Ebhardt H, Groszdanovic Z, Köhler D, Dreimann M, Hempfing A, Rudwaleit M, Stein H, Metz-Stavenhagen P, Sieper J, Loddenkemper C. Immunohistological analysis of zygapophyseal joints in patients with ankylosing spondylitis. *Arthritis Rheum* 2006; 54: 2845-2851 (b)

Appel H, Loddenkemper C, Kuhne M, Spiekermann S, Ebhardt H, Groszdanovic Z, Köhler D, Dreimann M, Hempfing A, Metz-Stavenhagen P, Rudwaleit M, Sieper J. Inflammation in the Spine of Patients with Ankylosing Spondylitis. Correlation of Histopathology with Findings in Magnetic Resonance Imaging (MRI). *Arthritis Res Ther* 2006 Aug 22;8(5):R143 [Epub ahead of print].(c)

Atagunduz P, **Appel H**, Kuon W, Wu P, Thiel A, Sieper J: CD8+ T cell response to self-peptides in ankylosing spondylitis. *Arthritis Rheum.* 2005; 52: 892-901.

Ball J: Pathology and pathogenesis. In: Moll JMH, editor. *Ankylosing spondylitis.* Edinburgh: Churchill Livingstone; 1980. p. 96-112.

Ball EJ, Khan MA. HLA-B27 polymorphism. *Joint Bone Spine.* 2001;68:378-82. Review.
Bardos T, Szabo Z, Czipri M, Vermes C, Tunyogi-Csapo M, Urban RM, Mikecz K, Glant TT. A longitudinal study on an autoimmune murine model of ankylosing spondylitis. *Ann Rheum Dis.* 2005 Jul;64(7):981-7. Epub 2005 Jan 7.

Benjamin RJ, Madrigal JA, Parham P: Peptide binding to empty HLA-B27 molecules of viable human cells *Nature* 1991, 351: 74-77.

Bieganowska K, Hllosberg P, Buckle GJ, Lim DG, Grten TF, Schneck J, Altman JD, Jacobson S, Ledis SL, Hanchard B, Chin J, Morgan O, Roth PA, Hafler DA: Direct analysis of viral-specific CD8+ T cells with soluble HLA-A2/tax 11-19 tetramer complexes in patients with human cell lymphotropic virus-associated myelopathy. *J Immunol* 1999, 162: 1765-1771.

Bollow M, Fischer T, Reissbauer H, Backhaus M, Sieper J, Hamm B, Braun J: Quantitative analyses of sacroiliac biopsies in spondyloarthropathies: T cells and macrophages predominate in early and active sacroiliitis- cellularity correlates with the degree of enhancement detected by magnetic resonance imaging. *Ann Rheum Dis* 2000, 59: 135-40.

Bowness P, Allen RL, Barclay DN, Jones EY, McMichael AJ: Importance of a conserved TCR J α -encoded tyrosine for T cell recognition of an HLA-B27/peptide complex. *Eur J Immunol* 1998, 28: 2704-2713.

Braun J, Bollow M, Neure L, Seipelt E, Seyrekbasan F, Herbst H, Eggens U, Distler A, Sieper J. Use of immunohistologic and in situ hybridization techniques in the examination of sacroiliac joint biopsy specimens from patients with ankylosing spondylitis. *Arthritis Rheum* 1995; 38: 499-505.

Braun J, Bollow M, Remlinger G, Sieper J: Prevalence of Spondylarthropathies in HLA B27-Positive and Negative Blood Donors. *Arthritis Rheum* 1998; 41:58-67.

Braun J, Yin Z, Spiller I, Siegert S, Rudwaleit M, Liu L, Radbruch A, Sieper J: Low secretion of tumor necrosis factor alpha, but no other Th1 or Th2 cytokines, by peripheral blood mononuclear cells correlates with chronicity in reactive arthritis. *Arthritis Rheum* 1999; 42: 2039-44.

Braun J and Sieper J: Spondyloarthritides and related arthritides. In: Warrel, TM Cox, JD Firth, EJ Benz (eds): *Oxford Textbook of Medicine* (Editors: DA), 4th Edition, Oxford University Press, New York 2003: 43-53

Brewerton DA, Caffrey M, Nicholls A, Walters D, Oates JK, James DC: reiter`s disease and HL-A 27. *Lancet* 1973, 2: 996-998.

Brosterhus H, Brings S, Leyendeckers H, Manz RA, Miltenyi S, Radbruch A, Assenmacher M, Schmitz J: Enrichment and detection of live antigen-specific CD4+ and CD8+ T cells based on cytokine secretion. *Eur J Immunol* 1999, 29: 4053-4059.

Brown MA, Crane AM, Wordsworth BP. Genetic aspects of susceptibility, severity, and clinical expression in ankylosing spondylitis. *Curr Opin Rheumatol*. 2002;14:354-60.

Bywaters EGL, Olsen E: A case of early ankylosing spondylitis with fatal secondary amyloidosis: demonstrated at the Royal Postgraduate Medical School. *BMJ* 1968; 2: 412-416.

Claudepierre P, Gueguen A, Ladjouze A, Hajjaj-Hassouni N, Sellami S, Amor B, Dougados M. Predictive factors of severity of spondyloarthropathy in North Africa. *Br J Rheumatol* 1995; 34:1139-45.

Cochran JR, Cameron TO, Stern LJ. The relationship of MHC-peptide binding and T cell activation probed using chemically defined MHC class II oligomers. *Immunity*. 2000; 12: 241-50.

Corr M, Slanetz AE, Boyd LF, Jelonek MT, Khilko S, al-Ramadi BK, Kim YS, Maher SE, Bothwell AL, Margulies DH. T cell receptor-MHC class I peptide interactions: affinity, kinetics, and specificity. *Science*. 1994; 265: 946-9.

Critchfield JM, Racke MK, Zuniga-Pflucker JC, Cannella B, Raine CS, Goverman J, LenardoMJ. T cell deletion in high antigen dose therapy of autoimmune encephalomyelitis. *Science*. 1994; 263: 1139-43.

Cruickshank B: Lesions of cartilaginous joints in ankylosing spondylitis. *J Pathol Bacteriol* 1956; 71: 73-84.

Day CL, Seth NP, Lucas M, **Appel H**, Gauthier L, Lauer GM, Robbins GK, Szczepiorkowski ZM, Casson DR, Chung RT, Bell S, Harcourt G, Walker BD, Klenerman P, Wucherpfennig KW: Ex vivo analysis of human memory CD4 T cells specific for hepatitis C virus using MHC class II tetramers. *J Clin Invest*. 2003; 112: 831-42.

Deane KH, Jecock RM, Pearce JH, Gaston JS: entification and characterization of a DR4-restricted T cell epitope within chlamydia heat shock protein 60. *Clin Exp Immunol* 1997; 109: 439-445.

de Vlam K, Mielants H, Veys EM. Involvement of the zygapophyseal joint in ankylosing spondylitis: relation to the bridging syndesmophyte. *J Rheumatol* 1999; 26: 1738-45.

Dickinson AM, Wang XN, Sviland L, Vyth-Dreese FA, Jackson GH, Schumacher TNM, Haanen JBAG, Mutis T, Goulmy: In situ dissection of graft-versus-host activities of cytotoxic T cells specific for minor histocompatibility antigens. *Nature Med* 2002; 8: 410-14.

Ferlin WG, Mougneau E, Hugues S, **Appel H**, Jang MH, Cazareth J, Beaudoin L, Schricke C, Lehuen A, Wucherpfennig KW, Glaichenhaus N. Self-peptides that bind with low affinity to the diabetes-associated I-A(g7) molecule readily induce T cell tolerance in non-obese diabetic mice. *Eur J Immunol.* 2004; 34: 2656-63.

Francois R. *Le Rachis dans la spondylarthrite ankylosante* (thesis). Brussels : University of Louvain; 1976, 18.

Francois RJ, Gardner DL, Degrave EJ, Bywaters EG. Histopathologic evidence that sacroiliitis in ankylosing spondylitis is not merely enthesitis. *Arthritis Rheum.* 2000; 43: 2011-24.

Garboczi, D.N., P. Ghosh, U. Utz, Q.R. Fan, W.E. Biddison, and D.C. Wiley: Structure of the complex between human T-cell receptor, viral peptide and HLA-A2. *Nature* 1992; 384: 134-41.

Gaston JS, Life PF, Granfors K, Merilahti-Palo R, Bailey L, Consalvey S, Toivanen A, Bacon PA: Synovial T lymphocyte recognition of organisms that trigger reactive arthritis. *Clin Exp Immunol* 1989, 76: 348-353.

Gaston JS, Deane KH, Jecock RM, Pearce JH: identification of 2 *Chlamydia trachomatis* antigens recognized by synovial fluid T cells from patients with *Chlamydia* induced reactive arthritis. *J Rheumatol* 1996, 23: 130-136.

Gimmi CD, Freeman GJ, Gribben JG, Gray G, Nadler LM. Human T-cell clonal anergy is induced by antigen presentation in the absence of B7 costimulation. *Proc Natl Acad Sci.* 1993; 90: 6586-90.

Goverman J, Woods A, Larson L, Weiner LP, Hood L, Zaller DM. Transgenic mice that express a myelin basic protein-specific T cell receptor develop spontaneous autoimmunity. *Cell.* 1993; 72: 551-60.

Granfors K, Jalkanen S, von Essen R, Lahesmaa-Rantala R, Isomaki O, Pekkola Heino K et al.: *Yersinia* antigens in synovial-fluid cells from patients with reactive arthritis. *N Engl J Med* 1989, 320: 216-221.

Granfors K, Jalkanen S, Lindberg AA, Maki Ikola O, von Essen R, Lahesmaa Ranatala R et al. *Salmonella* liposaccharide in synovial cells from patients with reactive arthritis. *Lancet* 1990, 335: 685-688.

Gray CM, Lawrence J, Schapiro JM, Altman JD, Winters MA, Crompton M, Loi M, Kundu SK, Davis MM, Merigan TC: Frequency of class I HLA-restricted anti-HIV CD8+ T cells in individuals receiving highly active antiretroviral therapy (HAART) *J Immunol* 1999, 162: 1780-1788.

Haanen JB, van Oijen MG, Tirion F, Oomen LC, Kruisbeek AM, Vyth-Dreese FA, Schumacher TNM: In situ detection of virus- and tumor specific T cell immunity. *Nat Med* 2000; 6: 1056-1060.

Hammer J, Zeidler H, Klimsa S, Heesemann J: *Yersinia enterocolitica* in the synovial membrane of patients with *Yersinia*-induced arthritis. *Arthritis Rheum* 1990, 33: 1795-1800.

Hermann E, Yu DT, Meyer zum Buschenfelde KH, Fleischer B. HLA-B27-restricted CD8 T cells derived from synovial fluids of patients with reactive arthritis and ankylosing spondylitis. *Lancet* 1993; 342: 646-50.

Hermann KG, Althoff CE, Schneider U, Zuhlsdorf S, Lembcke A, Hamm B, Bollow M: Spinal changes in patients with spondyloarthritis: comparison of MR imaging and radiographic appearances. *Radiographics* 2005, 25: 559-69.

Huang F, Zhang J, Zhu J, Guo J, Yang C. Juvenile spondyloarthropathies: the Chinese experience. *Rheum Dis Clin North Am* 2003; 29:531-47.

Jevtic V, Kos-Golja M, Rozman B, McCall I: Marginal erosive discovertebral "Romanus" lesions in ankylosing spondylitis demonstrated by contrast enhanced Gd-DTPA magnetic resonance imaging. *Skeletal Radiol* 2000, 29: 27-33.

Keat A: Reiter`s Syndrome and reactive arthritis in perspective. *N Engl J Med* 1983, 309: 1606-1616.

Kern F, Khatamzas E, Surel I, Frömmel C, Reinke P, Waldrop SL, Picker LJ, Volk HD: Distribution of human CMV-specific memory T cells among CD8-positive subsets defined by CD57, CD27 and CD45 isoforms. *Eur J Immunol* 1999, 29: 2908-2915.

Khan MA Spondyloarthropathies. *Curr Opin Rheumatol* 1998; 10:279-81.

Kahn M: Spondylarthropathies. *Atlas of Rheumatology*, Editor Gene G Hunder. Current Medicine, Inc. 2.Edition; 2001; pp 5.1.-5.18.

Kuckelhorn U, Ruppert T, Strehl B, Jungblut PR, Zimny-Arndt U, Lamer S, Prinz I, Drung I, Kloetzel PM, Kaufmann SHE, Steinhoff U: Link between organ-specific antigen-processing and CD8+ T cell mediated autoimmunity. *J Exp Med*. 2002; 195: 983-990.

Kuon W, Lauster R, Böttcher U, Koroknay A, Ulbrecht M, Hartmann M, Grolms M, Ugrinovic S, Braun J, Weiss EH, Sieper J: recognition of chlamydial antigen by HLA-b27 restricted cytotoxic T cells in HLA-B2705 transgenic CBA (H-2k) mice. *Arthritis Rheum* 1997, 40: 945-954.

Kuon W, Holzhutter HG, **Appel H**, Grolms M, Kollnberger S, Traeder A, Henklein P, Weiss E, Thiel A, Lauster R, Bowness P, Radbruch A, Kloetzel PM, Sieper J. Identification of HLA-B27-restricted peptides from the *Chlamydia trachomatis* proteome with possible relevance to HLA-B27-associated diseases. *J Immunol* 2001; 167: 4738-46.

Kuon W, Sieper J Identification of HLA-B27-restricted peptides in reactive arthritis and other spondyloarthropathies: computer algorithms and fluorescent activated cell sorting analysis as tools for hunting of HLA-B27-restricted chlamydial and autologous crossreactive peptides involved in reactive arthritis and ankylosing spondylitis. *Rheum Dis Clin North Am* 2003; 29:595-611.

Kuon W, Kuhne M, Busch DH, Atagunduz P, Seipel M, Wu P, Morawietz L, Fernahl G, **Appel H**, Weiss EH, Krenn V, Sieper J. Identification of novel human aggrecan T cell epitopes in HLA-B27 transgenic mice associated with spondyloarthropathy. *J Immunol*. 2004; 173: 4859-66.

Laloux L, Voisin MC, Allain J, Martin N, Kerboull L, Chevalier X, Claudepierre P. Immunohistological study of entheses in spondyloarthropathies: comparison in rheumatoid arthritis and osteoarthritis. *Ann Rheum Dis* 2001; 60: 316-21.

Lau CS, Burgos-Vargas R, Louthrenoo W, Mok MY, Wordsworth P, Zeng QY. Features of spondyloarthritis around the world. *Rheum Dis Clin North Am* 1998; 24:753-70.

Leirisalo-Repo M. Prognosis, course of disease, and treatment of the spondyloarthropathies. *Rheum Dis Clin North Am* 1998; 24:737-51.

Madden DR, Gorga JC, Strominger JL, Wiley DC: The structure of HLA-B27 reveals nonamer self peptides bound in an extended conformation. *Nature* 1991 353: 321-325.

Maksymowych WP. Ankylosing spondylitis--at the interface of bone and cartilage. *J Rheumatol*. 2000; 27: 2295-301.

Malherbe L, Filippi C, Julia V, Foucras G, Moro M, **Appel H**, Wucherpfennig K, Guery J, Glaichenhaus N: Selective Activation and Expansion of High-Affinity CD4(+) T Cells in Resistant Mice upon Infection with *Leishmania major*. *Immunity* 2000; 13: 771-782.

Matsui K, Boniface JJ, Reay PA, Schild H, Fazekas de St Groth B, Davis MM: Low affinity interaction of peptide-MHC complexes with T cell receptors. *Science* 1991, 254: 1788-1791.

McGavern DB, Christen U, Oldstone MBA : Molecular anatomy of antigen-specific CD8+ T cell engagement and synapse formation in vivo. *Nat Immunol* 2002; 3: 918-25.

McGonagle D, Gibbon W, Emery P. Classification of inflammatory arthritis by enthesitis. *Lancet* 1998; 352: 1137-40.

McGonagle D, Gibbon W, O'Connor P, Green M, Pease C, Emery P: Characteristic magnetic resonance imaging enthesal changes of knee synovitis in spondylarthropathy. *Arthritis Rheum* 1998; 41:694-700.

McGonagle D, Emery P. Enthesitis, osteitis, microbes, biomechanics, and immune reactivity in ankylosing spondylitis. *J Rheumatol*. 2000; 27: 2302-4.

Mertz AK, Batsford SR, Curschellas E, Kist MJ, Gondolf KB. Cationic *Yersinia* antigen-induced chronic allergic arthritis in rats. A model for reactive arthritis in humans. *J Clin Invest*. 1991; 88: 632-42.

Mertz AKH, Daser A, Skurnik M, Wiesmüller KH, Braun J, **Appel H**, Batsford S, Wu P, Distler A, Sieper J: The evolutionarily conserved ribosomal protein L23 and the cationic urease β -subunit of *Yersinia enterocolitica* O:3 belong to the immunodominant antigens in *Yersinia*-induced reactive arthritis: implications for autoimmunity. *Mol Med* 1994, 1:44-55.

Mertz AKH, Ugrinovic S, Lauster R, Wu P, Böttcher U, **Appel H**, Yin Z, Schiltz E, Batsford S, Schauer-Petrowski C, Braun J, Distler A, Sieper J: *Arthritis Rheum* 1998, 41: 315-326.

Mertz AKH, Wu P, Sturniolo T, Stoll D, Rudwaleit M, Lauster R, Braun J, Sieper J: Multispecific CD4+ T cells response to a single 12-mer epitope of the immunodominant heat-shock protein 60 of *Yersinia enterocolitica* in *Yersinia*-triggered reactive arthritis: Overlap with the B27-restricted CD8 epitope, functional properties, and epitope presentation by multiple DR alleles. *J Immunol*. 2000, 164: 1529-1537.

Metz-Stavenhagen P, Krebs S, Volpel HJ. [Surgical method in treatment of total kyphosis in ankylosing spondylitis] *Orthopade*. 2001; 30: 988-95.

Mitchison NA. The dosage requirements for immunological paralysis by soluble proteins. *Immunology*. 1968; 15: 509-30.

Mielants H, Veys EM, Cuvelier C, de Vos M. Ileocolonosopic findings in seronegative spondylarthropathies. *Br J Rheumatol* 1988; 27 Suppl 2:95-105.

Murali-Krishna K, Altman JD, Suresh M, Sourdive DJ, Zajac AJ, Miller JD, Slansky J, Ahmed R: Counting antigen-specific CD8 T cells: a reevaluation of bystander activation during viral infection. *Immunity* 1998, 8: 177-187.

Olhagen B. Urogenital syndromes and spondylarthritis. *Br J Rheumatol* 1983; 22 (4 Suppl.2): 33-40.

Poole AR. The histopathology of ankylosing spondylitis: are there unifying hypotheses? *Am J Med Sci.* 1998 Oct;316(4):228-33. Review.

Purmann J, Zeidler H, Bertrams J, Juli E, Cleveland S, Berges W, Gemsa R, Specker C, Reis HE: HLA antigens in ankylosing spondylitis associated with Crohn's disease. Increased frequency of the HLA phenotype B27,B44. *J Rheumatol* 1988; 15:1658-61.

Rath HC, Schultz M, Freitag R, Dieleman LA, Li F, Linde HJ, Scholmerich J, Sartor RB. Different subsets of enteric bacteria induce and perpetuate experimental colitis in rats and mice. *Infect Immun* 2001; 69:2277-85.

Rudwaleit M, Schwarzlose S, Listing J, Brandt J, Braun J, Sieper J: Do patients with ankylosing spondylitis (AS) without elevated CRP and without active inflammation as detected by magnetic resonance imaging (MRI) respond to TNF alpha blockers?. *Ann Rheum Dis* 2005, Suppl 64: 336.

Savill DL: The manubrio-sternal joint in ankylosing spondylitis. *J Bone Joint Surg Br* 1951; 33: 56-64.

Sieper J, Kingsley G, Palacios Boix A, Pitzalis C, Treharne J, Hughes R, Keat A, Panayi GS: Synovial T lymphocyte-specific immune response to *Chlamydia trachomatis* in Reiter's disease. *Arthritis Rheum* 1991, 24: 588-598.

Sieper J, Braun J; Pathogenesis of spondylarthropathies. Persistent bacterial, auto-immunity, or both? *Arthritis Rheum* 1995, 38: 1547-1554.

Sieper J, Kingsley G: Recent advances in the pathogenesis of reactive arthritis. *Immunol today* 1996, 17:160-163

Sieper J, Braun J, Kingsley GH. Report on the Fourth International Workshop on Reactive Arthritis. *Arthritis Rheum* 2000; 43:720-34.

Skinner PJ, Haase AT: In situ tetramer staining. *J Immunol Meth* 2002; 268: 29-34.

Sourdive DJ, Murali-Krishna K, Altman JD, Zajac AJ, Whitmire JK, Pannetier C, Kourilsky P, Evavold B, Sette A, Ahmed R: Conserved T cell repertoire in primary and memory CD8 T cell responses to an acute viral infection. *J Exp Med* 1998, 188: 71-82.

Steinman L. Multiple sclerosis: a coordinated immunological attack against myelin in the central nervous system. *Cell.* 1996; 85: 299-302.

Stern, L.J., J.H. Brown, T.S. Jardetzky, J.C. Gorga, R.G. Urban, J.L. Strominger, and D.C.

Wiley: Crystal structure of the human class II MHC protein HLA-DR1 complexed with an influenza virus peptide. *Nature* 1994, 368: 215-221.

Taurog JD, Richardson JA, Croft JT, Simmons WA, Zhou M, Fernandez-Sueiro JL, Balish E, Hammer RE. The germfree state prevents development of gut and joint inflammatory disease in HLA-B27 transgenic rats. *J Exp Med* 1994; 180:2359-64.

Ugrinovic S, Mertz A, Wu P, Braun J, Sieper J: A single nonamer from *Yersinia* 60kDa heat shock protein is the target of HLA-B27-restricted CTL response in *Yersinia*-induced reactive arthritis. *J Immunol* 1997, 159: 5715-5723.

Waldrop SL, Pitcher CJ, Peterson DM, Maino VC, Picker LJ: Determination of antigen-specific memory/effector CD4+ T cell frequencies by flow cytometry: evidence for a novel, antigen-specific homeostatic mechanism in HIV-associated immunodeficiency. *Clin Invest* 1997, 99:1739-1750.

Wilkinson M, Bywaters EG: Clinical features and course of ankylosing spondylitis; as seen in a follow-up of 222 hospital referred cases. *Ann Rheum Dis* 1958; 17: 209-28.

Wilson AJ, Murphy WA, Hardy DC, Totty WG: Transient osteoporosis: transient bone marrow edema? *Radiology* 1988; 167: 757-60.

Yin Z, Braun J, Neure L, Wu P, Liu L, Eggens U, Sieper J: Crucial role of interleukin-10/interleukin-12 balance in the regulation of the type 2 T helper cytokine response in reactive arthritis. *Arthritis Rheum* 1997; 40: 1788-97.

Zamvil SS, Steinman L. The T lymphocyte in experimental allergic encephalomyelitis. *Annu Rev Immunol.* 1990; 8: 579-621.

Zanetti M, Bruder E, Romero J, Hodler J. Bone marrow edema pattern in osteoarthritic knees: correlation between MR imaging and histologic findings. *Radiology* 2000, 215: 835-40.

Zou J, Zhang Y, Thiel A, Rudwaleit M, Shi SL, Radbruch A, Poole R, Braun J, Sieper J. Predominant cellular immune response to the cartilage autoantigenic G1 aggrecan in ankylosing spondylitis and rheumatoid arthritis. *Rheumatology (Oxford)*. 2003; 42: 846-55.

Zou J, **Appel H**, Rudwaleit M, Thiel A, Sieper J: Analysis of the CD8+ T cell response to the G1 domain of aggrecan in spondylitis. *Ann Rheum Dis* 2005; 64: 722-9. Epub 2004 Nov 11