

8. Literaturverzeichnis

- Agrawal, D., Hauser P., McPherson F., Dong F., Garcia A. und Pfloderer W.J. (1996). Repression of p27^{Kip1} synthesis by platelet-derived growth factor in BALB/c 3T3 cells. *Mol Cell Biol* 16(8): 4327-4336.
- Alani R.M., Young A.Z. und Shifflett C.B. (2001). Id1 regulation of cellular senescence through transcriptional repression of p16/Ink4a. *Proc Natl Acad Sci U S A* 98(14): 7812-7816.
- Barone M.V., Pepperkok R., Peverali F.A. und Philipson L. (1994). Id proteins control growth induction in mammalian cells. *Proc Natl Acad Sci U S A* 91(11): 4985-4988.
- Benezra R., Davis R.L., Lockshon D., Turner D.L. und Weintraub H. (1990). The protein Id: a negative regulator of helix-loop-helix DNA binding proteins. *Cell* 61(1): 49-59.
- Bennett M.R., Macdonald K., Chan S.W., Boyle J.J. und Weissberg P.L. (1998). Cooperative interactions between RB and p53 regulate cell proliferation, cell senescence, and apoptosis in human vascular smooth muscle cells from atherosclerotic plaques. *Circ Res* 82(6): 704-712.
- Ben-Porath I. und Weinberg R.A. (2005). The signals and pathways activating cellular senescence. *Int. J. Biochem. Cell Biol* (5): 961-976.
- Bloethner S., Hemminki K., Thirumaran R.K., Chen B., Mueller-Berghaus J., Ugurel S., Schadendorf D. und Kumar R. (2006). Differences in global gene expression in melanoma cell lines with and without homozygous deletion of the CDKN2A locus genes. *Melanoma Res* 16(4): 297-307.
- Bond J.A., Blaydes J.P., Rowson J., Haughton M.F., Smith J.R., Wynford-Thomas D. und Wyllie F.S. (1995). Mutant p53 rescues human diploid cells from senescence without inhibiting the induction of SDI1/WAF1. *Cancer Res* 55(11): 2404-2409.
- Braun-Dullaes R.C., Mann M.J. und Dzau V.J. (1998). Cell cycle progression: new therapeutic target for vascular proliferative disease. *Circulation*. 98(1): 82-89.
- Brown J.P., Wei W. und Sedivy J.M. (1997). Bypass of senescence after disruption of p21CIP1/WAF1 gene in normal diploid human fibroblasts. *Science* 277(5327): 831-834.
- Buckley S., Shi W., Driscoll B., Ferrario A., Anderson K. und Warburton D. (2004). BMP4 signaling induces senescence and modulates the oncogenic phenotype of A549 lung adeno-carcinoma cells. *Am. J. Physiol. Lung. Cell Mol Physiol* 286(1): L81-86.

- Burke A.P., Kolodgie F.D., Farb A., Weber D.K., Malcom G.T., Smialek J. und Virmani R. (2001). Healed plaque ruptures and sudden coronary death: evidence that subclinical rupture has a role in plaque progression. *Circulation* 103(7): 934-940.
- Calbó J., Serna C., Garriga J., Graña X. und Mazo A. (2004). The fate of pancreatic tumor cell lines following p16 overexpression depends on the modulation of CDK2 activity. *Cell Death Differ* 11(10): 1055-1065.
- Cannarile M.A., Lind N.A., Rivera R., Sheridan A.D., Camfield K.A., Wu B.B., Cheung K.P., Ding Z. und Goldrath A.W. (2006). Transcriptional regulator Id2 mediates CD8+ T cell immunity. *Nat Immunol* 7(12): 1317-1325.
- Chen J.H., Hales C.N. und Ozanne S.E. (2007). DNA damage, cellular senescence and organismal ageing: causal or correlative? *Nucleic Acids Res* 35(22): 7417-7428.
- Chen Q.M. (2000). Replicative senescence and oxidant-induced premature senescence. Beyond the control of cell cycle checkpoints. *Ann NY Acad Sci* 908: 111-125.
- Chen Q.M., Bartholomew J.C., Campisi J., Acosta M., Reagan D.J. und Ames B.N. (1998). Molecular analysis of H₂O₂-induced senescent-like growth arrest in normal human fibroblasts: p53 and Rb control G1 arrest but not cell replication. *Biochem. J* 332 (Pt 1): 43-50.
- Cheng M., Olivier P., Diehl J.A., Fero M., Roussel M.F., Roberts J.M. und Sherr C.S. (1999). The p21(Cip1) and p27(Kip1) CDK 'inhibitors' are essential activators of cyclin D-dependent kinases in murine fibroblasts. *Embo J* 18(6):1571-1583.
- Child E.S. und Mann D.J. (2006). The intricacies of p21 phosphorylation: protein/protein interactions, subcellular localization and stability. *Cell Cycle* (12): 1313-1319.
- Clarke M. und Bennett M. (2006). The emerging role of vascular smooth muscle cell apoptosis in atherosclerosis and plaque stability. *Am J Nephrol* 26: 531-535.
- Coqueret O. (2003). New roles for p21 and p27 cell-cycle inhibitors: a function for each cell compartment? *Trends Cell Biol* 13(2): 65-70.
- Dimri G.P., Lee X., Basile G., Acosta M., Scott G., Roskelley C., Medrane E.E., Linskens M., Rubelj I., Pereira-Smith O., Peacocke M. und Campisi J. (1995). A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc Natl Acad Sci U S A* 92(20): 9363-9367.
- Dong Y., Chi S.L., Borowsky A.D., Fan Y. und Weiss R.H. (2003). Cytosolic p21^{Waf1/Cip1} increases cell cycle transit in vascular smooth muscle cells. *Cell Signal* 16(2): 263-269.

- Dumont P., Burton M., Chen Q.M., Gonos E.S., Fripiat C., Mazarati J.B., Eliaers F., Remacle J. und Toussaint O. (2000). Induction of replicative senescence biomarkers by sub-lethal oxidative stresses in normal human fibroblast. *Free Radic Biol Med* 28(3): 361-373.
- Duprez D.A. (2007). Aldosterone and the vasculature: mechanisms mediating resistant hypertension. *J Clin Hypertens (Greenwich)* 9(1 Suppl 1): 13-18.
- Dyson N. (1998). The regulation of E2F by pRB-family proteins. *Genes Dev* 12: 2245-2262.
- El-Deiry W.S., Tokino T., Velculescu V.E., Levy D.B., Parsons R., Trent J.M., Lin D., Mercer W.E., Kinzler K.W., und Vogelstein B. (1993). WAF1, a potential mediator of p53 tumor suppression. *Cell* 75(4): 817-25.
- Fenton M., Barker S., Kurz D. und Erusalimsky J.D. (2001). Cellular senescence after single and repeated balloon catheter denudations of rabbit carotid arteries. *Arterioscler Thromb Vasc Biol* 21: 221-226.
- Forrest S. und McNamara C. (2004). Id family of transcription factors and vascular lesion formation. *Arterioscler. Thromb Vasc Biol* 24(11): 2014-2020.
- Forrest S.T., Taylor A.M., Sarembock I.J., Perlegas D. und McNamara C.A. (1999). Phosphorylation regulates Id3 function in vascular smooth muscle cells. *Circ Res* 95(6): 557-559.
- Forsyth N.R., Evans A.P., Shay J.W. und Wright W.E. (2003). Developmental differences in the immortalization of lung fibroblasts by telomerase. *Aging Cell* 2(5): 235-243.
- Gartel A.L. und Radhakrishnan S.K. (2005). Lost in Transcription: p21 repression, mechanism, and consequences. *Cancer Res* 65(10): 3980-3985.
- Gartel A.L. und Tyner A.L. (1999). Transcriptional regulation of the p21((WAF1/CIP1)) gene. *Exp. Cell Res* 246(2): 280-289.
- Geary R.L., Williams J.K., Golden D., Brown D.G., Benjamin M.E. und Adams M.R. (1996). Time course of cellular proliferation, intimal hyperplasia, and remodeling following angioplasty in monkeys with established atherosclerosis. A nonhuman primate model of restenosis. *Arterioscler Thromb Vasc Biol* 16(1): 34-43.
- Ghesquiere S.A.I. Anatomy overview of a human artery. Maastricht, November 2005. http://en.wikipedia.org/wiki/Image:Anatomy_artery.png
- Glagov S., Weisenberg E., Zarins C.K., Stankunavicius R. und Kolettis G.J. (1987). Compensatory enlargement of human atherosclerotic coronary arteries. *N Engl J Med* 316(22): 1371-1375.

- Goldstein S. (1990). Replicative senescence: the human fibroblast comes of age. *Science* 249(4973): 1129-1133.
- Gonos E.S., Derventzi A., Kveiborg M., Agiostratidou G., Kassem M., Clark B.F., Jat P.S. und Rattan S.I. (1998). Cloning and identification of genes that associate with mammalian replicative senescence. *Exp. Cell Res* 240(1): 66-74.
- Gorenne I., Kavurma M., Scott S. und Bennett M. (2006). Vascular smooth muscle cell senescence in atherosclerosis. *Cardiovasc Res* 72: 9-17.
- Hacker C., Kirsch R.D., Ju X.S., Hieronymus T., Gust T.C., Kuhl C., Jorgas T., Kurz S.M., Rose-John S., Yokota Y. und Zenke M. (2003). Transcriptional profiling identifies Id2 function in dendritic cell development *Nat Immunol* 4(4): 380-386.
- Hahn W.C., Stewart S.A., Brooks M.W., York S.G., Eaton E., Kurachi A., Beijersbergen R.L., Knoll J.H., Meyerson M. und Weinberg R.A. (1999). Inhibition of telomerase limits the growth of human cancer cells. *Nat Med* 10: 1164-1170.
- Hara E., Hall M. und Peters G. (1997). Cdk2-dependent phosphorylation of Id2 modulates activity of E2A-related transcription factors. *Embo J* 16(2): pp.332-342.
- Hayflick L. (1965). The limited in vitro lifetime of human diploid cell strains. *Exp Cell Res* 37: 614-636.
- Hayflick L. und Moorhead P.S. (1961). The serial cultivation of human diploid cell strains. *Exp. Cell Res* 25: 585-621.
- Herbig U., Jobling W.A., Chen B.P., Chen D.J. und Sedivy J.M. (2004). Telomere shortening triggers senescence of human cells through a pathway involving ATM, p53, and p21(CIP1), but not p16(INK4a). *Mol Cell* 2004 14(4): 501-513.
- Herbig U., Wei W., Dutriaux A., Jobling W.A. und Sedivy J.M. (2003). Real-time imaging of transcriptional activation in live cells reveals rapid up-regulation of the cyclin-dependent kinase inhibitor gene CDKN1A in replicative cellular senescence. *Aging Cell* 2(6): 295-304.
- Hua H., Zhang Y.Q., Dabernat S., Kritzik M., Dietz D., Sterling L. und Sarvetnick N. (2006). BMP4 regulates pancreatic progenitor cell expansion through Id2. *J Biol Chem* 281(19): 13574-13580.
- Iavarone A., Garg P., Lasorella A., Hsu J. und Israel M.A. (1994). The helix-loop-helix protein Id2 enhances cell proliferation and binds to the retinoblastoma protein. *Genes Dev* 8(11): 1270-1284.

- Itahana K., Zou Y., Itahana Y., Martinez J.L., Beausejour C., Jacobs J.J., Van Lohuizen M., Band V., Campisi J. und Dimri G.P. (2003). Control of the replicative life span of human fibroblasts by p16 and the polycomb protein Bmi1. *Mol Cell Biol* 23: 389-401.
- Krishnamurthy J., Torrice C., Ramsey M.R., Kovalev G.I., Al-Regaiey K., Su L. und Sharpless N.E. (2004). Ink4a/Arf expression is a biomarker of aging. *J Clin Invest* 114(9): 1299-1307.
- Kumar M.S., Hendrix J.A., Johnson A.D. und Owens G.K. (2003). Smooth muscle alpha-actin gene requires two E-boxes for proper expression in vivo and is a target of class I basic helix-loop-helix proteins. *Circ Res* 92(8): 840-847.
- Kurz D.J., Decary S., Hong Y. und Erusalimsky J.D. (2000). Senescence-associated (beta)-galactosidase reflects an increase in lysosomal mass during replicative ageing of human endo-thelial cells. *J Cell Sci* 113 (Pt 20): 3613-3622.
- Lasorella A., Iavarone A. und Israel M.A. (1996). Id2 specifically alters regulation of the cell cycle by tumor suppressor proteins. *Mol Cell Biol* 16(6): 2570-2578.
- Lin A.W., Barradas M., Stone J.C., van Aelst L., Serrano M. und Lowe S.W. (1998). Premature senescence involving p53 and p16 is activated in response to constitutive MEK/MAPK mitogenic signaling. *Genes Dev* 12(19): 3008-3019.
- Lloyd R.V., Erickson L.A., Jin L., Kulig E., Qian X., Cheville J.C. und Scheithauer B.W. (1999). p27kip1: a multifunctional cyclin-dependent kinase inhibitor with prognostic significance in human cancers. *Am J Pathol* 154(2): 313-323.
- Lundberg A.S. und Weinberg R.A. (1998). Functional inactivation of the retinoblastoma protein requires sequential modification by at least two distinct cyclin-cdk complexes. *Mol Cell Biol* 18(2):753-761.
- Lutgens E., van Suylen R.J., Faber B.C., Gijbels M.J., Eurlings P.M., Bijmens A.P., Cleutjens K.B., Heeneman S. und Daemen M.J. (2003). Atherosclerotic plaque rupture: local or systemic process? *Arterioscler Thromb Vasc Biol* 23(12): 2123-2130.
- Mann J. und Davies M.J. (1999). Mechanism of progression in native coronary artery disease: role of healed plaque rupture. *Heart* 82: 265-268.
- Martens UM, Chavez EA, Poon SS, Schmoor C und Lansdorp P.M. (2000). Accumulation of short telomeres in human fibroblasts prior to replicative senescence. *Exp Cell Res* 256(1): 291-299.
- Massari M.E. und Murre C. (2000). Helix-loop-helix proteins: regulators of transcription in eucaryotic organisms. *Mol Cell Biol* 20(2): 429-440.

- Matsumara T., Zerrudo Z. und Hayflick L. (1979). Senescent human diploid cells in cul-ture: survival, DNA synthesis and morphology. *J Gerontol* 34(3): 328-334.
- Matsumura M.E., Li F., Berthoux L., Wei B., Lobe D.R., Jeon C., Hammarskjöld M.L. und McNamara C.A. (2001). Vascular injury induces posttranscriptional regulation of the Id3 gene: cloning of a novel Id3 isoform expressed during vascular lesion formation in rat and human atherosclerosis. *Arterioscler Thromb Vasc Biol* 21(5): 752-758.
- Matsumura M.E., Lobe D.R. und McNamara C.A. (2002). Contribution of the helix-loop-helix factor Id2 to regulation of vascular smooth muscle cell proliferation. *J Biol Chem* 277(9): 7293-7297.
- Matthews C., Gorenne I., Scott S., Figg N., Kirkpatrick P., Ritchie A., Goddard M. und Bennett M. (2006). Vascular smooth muscle cells undergo telomere-based senescence in human atherosclerosis: effects of telomerase and oxidative stress. *Circ Res* 99(2): 156-164.
- Mehta P.K. und Griendling K.K. (2007). Angiotensin II cell signaling: physiological and pathological effects in the cardiovascular system. *Am J Physiol Cell Physiol* 292(1): C82-97.
- Minamino T. und Komuro I. (2007). Vascular cell senescence: contribution to atherosclerosis. *Circ Res* (1): 15-26.
- Minamino T., Yoshida T., Tateno K., Miyauchi H., Zou Y., Toko H. und Komuro I. (2003). Ras induces vascular smooth muscle cell senescence and inflammation in human atherosclerosis. *Circulation* 108(18): 2264-2269.
- Mitra J., Dai C.Y., Somasundaram K., El-Deiry W.S., Satyamoorthy K., Herlyn M. und Enders G.H. (1999). Induction of p21(WAF1/CIP1) and inhibition of Cdk2 mediated by the tumor suppressor p16(INK4a). *Mol Cell Biol* 19(5): 3916-3928.
- Miyazono K. und Miyazawa K. (2002). Id: a target of BMP signaling. *Sci STKE* (151): PE40.
- Norton D.J. (2000). ID helix-loop-helix proteins in cell growth, differentiation and tumorigenesis. *J Cell Sci* 113(Pt 22): 3897-3905.
- O'Brian E.R., Alpers C.E., Stewart D.K., Ferguson M., Tran N., Gordon D., Benditt E.P., Hinohara T., Simpson J.B. und Schwartz S.M. (1993). Proliferation in Primary and Restenotic Coronary Atherectomy Tissue. Implications for antiproliferative therapy. *Circ Res* 73: 223-231.
- Ohtani N., Yamakoshi K., Takahashi A. und Hara E. (2004). The p16INK4a-RB pathway: molecular link between cellular senescence and tumor suppression. *J Med Invest* 51(3-4): 146-153.

- Ong K.L., Cheung B.M., Man Y.B., Lau C.P. und Lam K.S. (2007). Prevalence, awareness, treatment, and control of hypertension among United States adults 1999-2004. *Hypertension* 49(1): 69-75.
- Orlandi A., Bochaton-Piallat M.L., Gabbiani G. und Spagnoli L.G. (2006). Aging, smooth muscle cells and vascular pathobiology: implications for atherosclerosis. *Atherosclerosis* 188(2): 221-230.
- O'Sullivan M., Scott S.D., McCarthy N., Figg N., Shapiro L.M., Kirkpatrick P., und Bennett M.R. (2003). Differential cyclin E expression in human in-stent stenosis smooth muscle cells identifies targets for selective anti-restenosis therapy. *Cardiovasc Res* 60(3): 673-683.
- Owens G.K., Kumar M.S. und Warmhoff B. (2004). Molecular regulation of vascular smooth muscle cell differentiation in development and disease. *Physiol Rev* 84: 767-801.
- Pantoja C. und Serrano M. (1999). Murine fibroblasts lacking p21 undergo senescence and are resistant to transformation by oncogenic Ras. *Oncogene* 18(35): 4974-4982.
- Petersen S., Peto V., Rayner M., Leal J., Luengo-Fernandez R. und Gray A. (2005). European cardiovascular disease statistics 2005 edition. BHF: London.
- Prabhu S., Ignatova A., Park S.T. und Sun X.H. (1997). Regulation of the expression of cyclin-dependent kinase inhibitor p21 by E2A and Id proteins. *Mol Cell Biol* 17(10): 5888-5896.
- Raines E.W. und Ross R. (1993). Smooth muscle cell and the pathogenesis of the lesions of atherosclerosis. *Br Heart J* 69: S30-S37.
- Raines E.W. und Ross R. (1995). Biology of atherosclerotic plaque formation: possible role of growth factors in lesion development and the potential impact of soy. *J Nutr* 125(3 Suppl): 624S-630S.
- Ross R. (1999). Atherosclerosis - an inflammatory disease. *N Engl J Med* 340(2): 115-126.
- Ruas M. und Peters G. (1998). The p16INK4a/CDKN2A tumor suppressor and its relatives. *Biochem Biophys Acta* 1378(2): F115-177.
- Ruiz-Torres A., Gimeno A., Melon J., Mendez L., Munoz F.J. und Macia M. (1999). Age-related loss of proliferative activity in human vascular smooth muscle cells in culture. *Mech Ageing Dev* 110: 49-55.
- Ruiz-Torres A., Lozano R., Melon J. und Carraro R. (2003). Age-dependent decline of in vitro migration (basal and stimulated by IGF-1 or insulin) of human vascular smooth muscle cells. *J Gerontol A Biol Sci Med Sci* 58: B1074-B1077.

- Samanta J. und Kessler J.A. (2004). Interactions between ID and OLIG proteins mediate the inhibitory effects of BMP4 on oligodendroglial differentiation. *Development* 131(17): 4131-4142.
- Sanz-González S.M., Poch E., Pérez-Roger I., Díez-Juan A., Ivorra C. und Andrés V. (2000). Control of vascular smooth muscle cell growth by cyclin-dependent kinase inhibitory proteins and its implication in cardiovascular disease. *Front Biosci* 5: D619-628.
- Sherr C.J. und Roberts J.M. (1999). CDK inhibitors: positive and negative regulators of G₁-phase progression. *Genes Dev* 13: 1501-1512.
- Sherr C.J. und Roberts J.M. (2004). Living with or without cyclins and cyclin-dependent kinases. *Genes Dev* 18: 2699-2711.
- Sherwood S.W., Rush D., Ellsworth J.L. und Schmike R.T. (1988). Defining cellular senescence in IMR-90 cells: a flow cytometric analysis. *Proc Natl Acad Sci U S A* 85(23): 9086-9090.
- Springhorn J.P., Singh K., Kelly R.A. und Smith T.W. (1994). Posttranscriptional regulation of Id1 activity in cardiac muscle. Alternative splicing of novel Id1 transcript permits homodimerization. *J Biol Chem* 269(7): 5132-5136.
- Stein G.H., Drullinger L.F., Soulard A. und Dulić V. (1999). Differential roles for cyclin-dependent kinase inhibitors p21 and p16 in the mechanisms of senescence and differentiation in human fibroblasts. *Mol Cell Biol* 19(3): 2109-2117.
- Steinberg D., Parthasarathy S., Carew T.E., Khoo J.C. und Witztum J.L. (1989). Beyond Cholesterol modifications of low-density lipoprotein that increase its atherogenicity. *N Engl J Med* 320: 915-924.
- Susaki E., Nakajama K. und Nakajama K.I. (2007). Cyclin D2 translocates p27 out of the nucleus and promotes its degradation at the G₀-G₁ Transition. *Mol Cell Biol* 27(13): 4646-4640.
- Sutterlüty H., Chatelain E., Marti A., Wirbelauer C., Senften M., Müller U. und Krek W. (1999). p45SKP2 promotes p27Kip1 degradation and induces S phase in quiescent cells. *Nat Cell Biol* 1(4): 207-214.
- Takai H., Smogorzewsky A. und de Lange T. (2003). DNA damage foci at dysfunctional telomeres. *Curr Biol* 13(17): 1549-1556.
- Tanner F.C., Yang Z.Y., Duckers E., Gordon D., Nabel G.J. und Nabel E.G. (1998). Expression of cyclin-dependent kinase inhibitors in vascular disease. *Circ Res* 82(3): 396-403.

- Virmani R., Kolodgie F.D., Burke A.P., Farb A. und Schwartz S.M. (2000). Lessons from sudden coronary death: a comprehensive morphological classification scheme for atherosclerotic lesions. *Arterioscler Thromb Vasc Biol* 20(5): 1262-1275.
- Waga S., Hannon G.J., Beach D. und Stillman B. (1994). The p21 inhibitor of cyclin-dependent kinases controls DNA replication by interaction with PCNA. *Nature* 369(6481): 574-578.
- Webley K., Bond J.A., Jones C.J., Blaydes J.P., Craig A., Hupp T. und Wynford-Thomas D. (2000). Posttranslational modifications of p53 in replicative senescence overlapping but distinct from those induced by DNA damage. *Mol Cell Biol* 20(8): 2803-280
- Wei G.L., Krasinski K., Kearney M., Isner J.M., Walsh K. und Andrés V. (1997). Temporally and spatially coordinated expression of cell cycle regulatory factors after angioplasty. *Circ Res* 80(3): 418-426.
- WHO (2007). Cardiovascular diseases. http://www.who.int/cardiovascular_diseases/en
- Yokota Y., Mansouri A., Mori S., Sugawara S., Adachi S., Nishikawa S. und Gruss P. (1999). Development of peripheral lymphoid organs and natural killer cells depends on the helix-loop-helix inhibitor Id2. *Nature* 397(6721): 702-706.
- Zebedee Z. und Hara E. (2001). Id proteins in cell cycle control and cellular senescence. *Oncogene*. 20(58): 8317-8325.
- Zhang H. (2007). Molecular signaling and genetic pathways of senescence: Its role in tumorigenesis and aging. *J Cell Physiol* 210(3):567-74.
- Zhang H., Pan K.H. und Cohen S.N. (2003). Senescence-specific gene expression fingerprints reveal cell-type-dependent physical clustering of up-regulated chromosomal loci. *Proc Natl Acad Sci U S A* 100(6): 3251-3256.
- Zheng W., Wang H., Xue L., Zhang Z. und Tong T. (2004). Regulation of cellular senescence and p16(INK4a) expression by Id1 and E47 proteins in human diploid fibroblast. *J Biol Chem* 279(30): 31524-31532