

## 6 Literatur

- Baccei, M. L. und Fitzgerald, M.** (2004). Development of GABAergic and glycinergic transmission in the neonatal rat dorsal horn. *J Neurosci* **24**, 4749-57.
- Baumgardt, M., Miguel-Aliaga, I., Karlsson, D., Ekman, H. und Thor, S.** (2007). Specification of neuronal identities by feedforward combinatorial coding. *PLoS Biol* **5**, e37.
- Beatus, P. und Lendahl, U.** (1998). Notch and neurogenesis. *J Neurosci Res* **54**, 125-36.
- Beinfeld, M. C.** (2003). Biosynthesis and processing of pro CCK: recent progress and future challenges. *Life Sci* **72**, 747-57.
- Ben-Ari, Y.** (2002). Excitatory actions of gaba during development: the nature of the nurture. *Nat Rev Neurosci* **3**, 728-39.
- Berger, S. L. und Kimmel, A. R.** (1987). *Guide To Molecular Cloning Techniques*. Academic Press, San Diego, CA, USA.
- Birmingham, N. A., Hassan, B. A., Wang, V. Y., Fernandez, M., Banfi, S., Bellen, H. J., Fritzsch, B. und Zoghbi, H. Y.** (2001). Proprioceptor pathway development is dependent on Math1. *Neuron* **30**, 411-22.
- Bertrand, N., Castro, D. S. und Guillemot, F.** (2002). Proneural genes and the specification of neural cell types. *Nat Rev Neurosci* **3**, 517-30.
- Birnboim, H. C. und Doly, J.** (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* **7**, 1513-23.
- Briscoe, J., Chen, Y., Jessell, T. M. und Struhl, G.** (2001). A hedgehog-insensitive form of patched provides evidence for direct long-range morphogen activity of sonic hedgehog in the neural tube. *Mol Cell* **7**, 1279-91.
- Briscoe, J. und Ericson, J.** (2001). Specification of neuronal fates in the ventral neural tube. *Curr Opin Neurobiol* **11**, 43-9.
- Briscoe, J., Pierani, A., Jessell, T. M. und Ericson, J.** (2000). A homeodomain protein code specifies progenitor cell identity and neuronal fate in the ventral neural tube. *Cell* **101**, 435-45.
- Briscoe, J., Sussel, L., Serup, P., Hartigan-O'Connor, D., Jessell, T. M., Rubenstein, J. L. und Ericson, J.** (1999). Homeobox gene Nkx2.2 and specification of neuronal identity by graded Sonic hedgehog signalling. *Nature* **398**, 622-7.
- Brown, A. G. und Fyffe, R. E.** (1981). Form and function of dorsal horn neurones with axons ascending the dorsal columns in cat. *J Physiol* **321**, 31-47.
- Burgess, P. R. und Perl, E. R.** (1967). Myelinated afferent fibres responding specifically to noxious stimulation of the skin. *J Physiol* **190**, 541-62.
- Cai, J., Qi, Y., Hu, X., Tan, M., Liu, Z., Zhang, J., Li, Q., Sander, M. und Qiu, M.** (2005). Generation of oligodendrocyte precursor cells from mouse dorsal spinal cord independent of Nkx6 regulation and Shh signaling. *Neuron* **45**, 41-53.
- Caspary, T. und Anderson, K. V.** (2003). Patterning cell types in the dorsal spinal cord: what the mouse mutants say. *Nat Rev Neurosci* **4**, 289-97.

- Charron, F. und Tessier-Lavigne, M.** (2005). Novel brain wiring functions for classical morphogens: a role as graded positional cues in axon guidance. *Development* **132**, 2251-62.
- Chen, Z. F., Rebelo, S., White, F., Malmberg, A. B., Baba, H., Lima, D., Woolf, C. J., Basbaum, A. I. und Anderson, D. J.** (2001). The paired homeodomain protein DRG11 is required for the projection of cutaneous sensory afferent fibers to the dorsal spinal cord. *Neuron* **31**, 59-73.
- Cheng, L., Arata, A., Mizuguchi, R., Qian, Y., Karunaratne, A., Gray, P. A., Arata, S., Shirasawa, S., Bouchard, M., Luo, P. et al.** (2004). Tlx3 and Tlx1 are post-mitotic selector genes determining glutamatergic over GABAergic cell fates. *Nat Neurosci* **7**, 510-7.
- Cheng, L., Samad, O. A., Xu, Y., Mizuguchi, R., Luo, P., Shirasawa, S., Goulding, M. und Ma, Q.** (2005). Lbx1 and Tlx3 are opposing switches in determining GABAergic versus glutamatergic transmitter phenotypes. *Nat Neurosci* **8**, 1510-5.
- Chiang, C., Litingtung, Y., Lee, E., Young, K. E., Corden, J. L., Westphal, H. und Beachy, P. A.** (1996). Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. *Nature* **383**, 407-13.
- Chiou, L. C., Liao, Y. Y., Fan, P. C., Kuo, P. H., Wang, C. H., Riemer, C. und Prinssen, E. P.** (2007). Nociceptin/orphanin FQ peptide receptors: pharmacology and clinical implications. *Curr Drug Targets* **8**, 117-35.
- Conner, A. C., Simms, J., Barwell, J., Wheatley, M. und Poyner, D. R.** (2007). Ligand binding and activation of the CGRP receptor. *Biochem Soc Trans* **35**, 729-32.
- Cordero-Erausquin, M., Coull, J. A., Boudreau, D., Rolland, M. und De Koninck, Y.** (2005). Differential maturation of GABA action and anion reversal potential in spinal lamina I neurons: impact of chloride extrusion capacity. *J Neurosci* **25**, 9613-23.
- Dado, R. J., Katter, J. T. und Giesler, G. J., Jr.** (1994). Spinothalamic and spinohypothalamic tract neurons in the cervical enlargement of rats. I. Locations of antidromically identified axons in the thalamus and hypothalamus. *J Neurophysiol* **71**, 959-80.
- Dickinson, M. E., Selleck, M. A., McMahon, A. P. und Bronner-Fraser, M.** (1995). Dorsalization of the neural tube by the non-neural ectoderm. *Development* **121**, 2099-106.
- Ding, Y. Q., Yin, J., Kania, A., Zhao, Z. Q., Johnson, R. L. und Chen, Z. F.** (2004). Lmx1b controls the differentiation and migration of the superficial dorsal horn neurons of the spinal cord. *Development* **131**, 3693-703.
- Dockray, G. J.** (1987). The biosynthesis of regulatory peptides. *Am Rev Respir Dis* **136**, S9-15.
- Ericson, J., Rashbass, P., Schedl, A., Brenner-Morton, S., Kawakami, A., van Heyningen, V., Jessell, T. M. und Briscoe, J.** (1997). Pax6 controls progenitor cell identity and neuronal fate in response to graded Shh signaling. *Cell* **90**, 169-80.
- Erlander, M. G., Tillakaratne, N. J., Feldblum, S., Patel, N. und Tobin, A. J.** (1991). Two genes encode distinct glutamate decarboxylases. *Neuron* **7**, 91-100.

- Fisher, J. M., Sossin, W., Newcomb, R. und Scheller, R. H.** (1988). Multiple neuropeptides derived from a common precursor are differentially packaged and transported. *Cell* **54**, 813-22.
- Florén, A., Land, T. und Langel, U.** (2000). Galanin receptor subtypes and ligand binding. *Neuropeptides* **34**, 331-7.
- Fremeau, R. T., Jr., Troyer, M. D., Pahner, I., Nygaard, G. O., Tran, C. H., Reimer, R. J., Bellocchio, E. E., Fortin, D., Storm-Mathisen, J. und Edwards, R. H.** (2001). The expression of vesicular glutamate transporters defines two classes of excitatory synapse. *Neuron* **31**, 247-60.
- Fürst, S.** (1999). Transmitters involved in antinociception in the spinal cord. *Brain Res Bull* **48**, 129-41.
- Gentleman, R. C., Carey, V. J., Bates, D. M., Bolstad, B., Dettling, M., Dudoit, S., Ellis, B., Gautier, L., Ge, Y., Gentry, J. et al.** (2004). Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol* **5**, R80.
- Gillespie, P. G. und Walker, R. G.** (2001). Molecular basis of mechanosensory transduction. *Nature* **413**, 194-202.
- Glasgow, S. M., Henke, R. M., Macdonald, R. J., Wright, C. V. und Johnson, J. E.** (2005). Ptfla determines GABAergic over glutamatergic neuronal cell fate in the spinal cord dorsal horn. *Development* **132**, 5461-9.
- Goebbels, S., Bode, U., Pieper, A., Funfschilling, U., Schwab, M. H. und Nave, K. A.** (2005). Cre/loxP-mediated inactivation of the bHLH transcription factor gene NeuroD/BETA2. *Genesis* **42**, 247-52.
- Goebbels, S., Bormuth, I., Bode, U., Hermanson, O., Schwab, M. H. und Nave, K. A.** (2006). Genetic targeting of principal neurons in neocortex and hippocampus of NEX-Cre mice. *Genesis* **44**, 611-21.
- Goridis, C. und Rohrer, H.** (2002). Specification of catecholaminergic and serotonergic neurons. *Nat Rev Neurosci* **3**, 531-41.
- Gowan, K., Helms, A. W., Hunsaker, T. L., Collisson, T., Ebert, P. J., Odom, R. und Johnson, J. E.** (2001). Crossinhibitory activities of Ngn1 and Math1 allow specification of distinct dorsal interneurons. *Neuron* **31**, 219-32.
- Grisel, J. E. und Mogil, J. S.** (2000). Effects of supraspinal orphanin FQ/nociceptin. *Peptides* **21**, 1037-45.
- Gross, M. K., Dottori, M. und Goulding, M.** (2002). Lbx1 specifies somatosensory association interneurons in the dorsal spinal cord. *Neuron* **34**, 535-49.
- Grunwald, I. C. und Klein, R.** (2002). Axon guidance: receptor complexes and signaling mechanisms. *Curr Opin Neurobiol* **12**, 250-9.
- Hao, S., Mata, M., Goins, W., Glorioso, J. C. und Fink, D. J.** (2003). Transgene-mediated enkephalin release enhances the effect of morphine and evades tolerance to produce a sustained antiallodynic effect in neuropathic pain. *Pain* **102**, 135-42.
- Helms, A. W., Battiste, J., Henke, R. M., Nakada, Y., Simplicio, N., Guillemot, F. und Johnson, J. E.** (2005). Sequential roles for Mash1 and Ngn2 in the generation of dorsal spinal cord interneurons. *Development* **132**, 2709-19.

- Helms, A. W. und Johnson, J. E.** (2003). Specification of dorsal spinal cord interneurons. *Curr Opin Neurobiol* **13**, 42-9.
- Henderson, G. und McKnight, A. T.** (1997). The orphan opioid receptor and its endogenous ligand-nociceptin/orphanin FQ. *Trends Pharmacol Sci* **18**, 293-300.
- Hitoshi, S., Alexson, T., Tropepe, V., Donoviel, D., Elia, A. J., Nye, J. S., Conlon, R. A., Mak, T. W., Bernstein, A. und van der Kooy, D.** (2002). Notch pathway molecules are essential for the maintenance, but not the generation, of mammalian neural stem cells. *Genes Dev* **16**, 846-58.
- Hökfelt, T.** (1991). Neuropeptides in perspective: the last ten years. *Neuron* **7**, 867-79.
- Hökfelt, T., Broberger, C., Xu, Z. Q., Sergeyev, V., Ubink, R. und Diez, M.** (2000). Neuropeptides--an overview. *Neuropharmacology* **39**, 1337-56.
- Hökfelt, T., Brumovsky, P., Shi, T., Pedrazzini, T. und Villar, M.** (2007). NPY and pain as seen from the histochemical side. *Peptides* **28**, 365-72.
- Hori, K., Cholewa-Waclaw, J., Nakada, Y., Glasgow, S. M., Masui, T., Henke, R. M., Wildner, H., Martarelli, B., Beres, T. M., Epstein, J. A. et al.** (2008). A nonclassical bHLH Rbpj transcription factor complex is required for specification of GABAergic neurons independent of Notch signaling. *Genes Dev* **22**, 166-78.
- Hunt, S. P. und Mantyh, P. W.** (2001). The molecular dynamics of pain control. *Nat Rev Neurosci* **2**, 83-91.
- Hynes, M., Ye, W., Wang, K., Stone, D., Murone, M., Sauvage, F. und Rosenthal, A.** (2000). The seven-transmembrane receptor smoothened cell-autonomously induces multiple ventral cell types. *Nat Neurosci* **3**, 41-6.
- Ince-Dunn, G., Hall, B. J., Hu, S. C., Ripley, B., Huganir, R. L., Olson, J. M., Tapscott, S. J. und Ghosh, A.** (2006). Regulation of thalamocortical patterning and synaptic maturation by NeuroD2. *Neuron* **49**, 683-95.
- Innis, M. A., Gelfand, D. H. und Sninsky, J. J.** (1989). *PCR Protocols: A Guide To Methods And Applications*. Academic Press, San Diego, CA, USA.
- Inoue, H., Nojima, H. und Okayama, H.** (1990). High efficiency transformation of Escherichia coli with plasmids. *Gene* **96**, 23-8.
- Jerpseth, B., Greener, A., Short, J. M., Viola, J. und Kretz, P.** (1992). XL1-Blue MRF' E. coli cells: McrA-, McrCB-, McrF-, Mrr-, HsdR- derivative of XL1-Blue cells. *Strategies* **5**, 81-83.
- Jessell, T. M.** (2000). Neuronal specification in the spinal cord: inductive signals and transcriptional codes. *Nat Rev Genet* **1**, 20-9.
- Ji, R. R., Kohno, T., Moore, K. A. und Woolf, C. J.** (2003). Central sensitization and LTP: do pain and memory share similar mechanisms? *Trends Neurosci* **26**, 696-705.
- John, A., Wildner, H. und Britsch, S.** (2005). The homeodomain transcription factor Gbx1 identifies a subpopulation of late-born GABAergic interneurons in the developing dorsal spinal cord. *Dev Dyn* **234**, 767-71.
- Julius, D. und Basbaum, A. I.** (2001). Molecular mechanisms of nociception. *Nature* **413**, 203-10.
- Kaneko, T. und Fujiyama, F.** (2002). Complementary distribution of vesicular glutamate transporters in the central nervous system. *Neurosci Res* **42**, 243-50.

- Katter, J. T., Dado, R. J., Kostarczyk, E. und Giesler, G. J., Jr.** (1996). Spinothalamic and spinohypothalamic tract neurons in the sacral spinal cord of rats. I. Locations of antidromically identified axons in the cervical cord and diencephalon. *J Neurophysiol* **75**, 2581-605.
- Kawaguchi, Y., Cooper, B., Gannon, M., Ray, M., MacDonald, R. J. und Wright, C. V.** (2002). The role of the transcriptional regulator Ptfla in converting intestinal to pancreatic progenitors. *Nat Genet* **32**, 128-34.
- Kelly, R. B.** (1993). Storage and release of neurotransmitters. *Cell* **72 Suppl**, 43-53.
- Kerr, B. J., Cafferty, W. B., Gupta, Y. K., Bacon, A., Wynick, D., McMahon, S. B. und Thompson, S. W.** (2000). Galanin knockout mice reveal nociceptive deficits following peripheral nerve injury. *Eur J Neurosci* **12**, 793-802.
- Kessaris, N., Pringle, N. und Richardson, W. D.** (2001). Ventral neurogenesis and the neuron-glial switch. *Neuron* **31**, 677-80.
- Kim, J., Wu, H. H., Lander, A. D., Lyons, K. M., Matzuk, M. M. und Calof, A. L.** (2005). GDF11 controls the timing of progenitor cell competence in developing retina. *Science* **308**, 1927-30.
- King, M., Chang, A. und Pasternak, G. W.** (1998). Functional blockade of opioid analgesia by orphanin FQ/nociceptin. *Biochem Pharmacol* **55**, 1537-40.
- Konishi, T.** (2006). Detection and restoration of hybridization problems in affymetrix GeneChip data by parametric scanning. *Genome Inform* **17**, 100-9.
- Koster, A., Montkowski, A., Schulz, S., Stube, E. M., Knaudt, K., Jenck, F., Moreau, J. L., Nothacker, H. P., Civelli, O. und Reinscheid, R. K.** (1999). Targeted disruption of the orphanin FQ/nociceptin gene increases stress susceptibility and impairs stress adaptation in mice. *Proc Natl Acad Sci U S A* **96**, 10444-9.
- Kriks, S., Lanuza, G. M., Mizuguchi, R., Nakafuku, M. und Goulding, M.** (2005). Gsh2 is required for the repression of Ngn1 and specification of dorsal interneuron fate in the spinal cord. *Development* **132**, 2991-3002.
- Kupfermann, I.** (1991). Functional studies of cotransmission. *Physiol Rev* **71**, 683-732.
- Kwan, K. M. und Behringer, R. R.** (2002). Conditional inactivation of Lim1 function. *Genesis* **32**, 118-20.
- Lai, J., Luo, M. C., Chen, Q., Ma, S., Gardell, L. R., Ossipov, M. H. und Porreca, F.** (2006). Dynorphin A activates bradykinin receptors to maintain neuropathic pain. *Nat Neurosci* **9**, 1534-40.
- Lecci, A. und Maggi, C. A.** (2001). Tachykinins as modulators of the micturition reflex in the central and peripheral nervous system. *Regul Pept* **101**, 1-18.
- Lee, K. J., Dietrich, P. und Jessell, T. M.** (2000). Genetic ablation reveals that the roof plate is essential for dorsal interneuron specification. *Nature* **403**, 734-40.
- Lee, K. J. und Jessell, T. M.** (1999). The specification of dorsal cell fates in the vertebrate central nervous system. *Annu Rev Neurosci* **22**, 261-94.
- Lee, K. J., Mendelsohn, M. und Jessell, T. M.** (1998). Neuronal patterning by BMPs: a requirement for GDF7 in the generation of a discrete class of commissural interneurons in the mouse spinal cord. *Genes Dev* **12**, 3394-407.

- Li, H., Zeitler, P. S., Valerius, M. T., Small, K. und Potter, S. S.** (1996). Gsh-1, an orphan Hox gene, is required for normal pituitary development. *Embo J* **15**, 714-24.
- Li, M. Z., Wang, J. S., Jiang, D. J., Xiang, C. X., Wang, F. Y., Zhang, K. H., Williams, P. R. und Chen, Z. F.** (2006). Molecular mapping of developing dorsal horn-enriched genes by microarray and dorsal/ventral subtractive screening. *Dev Biol* **292**, 555-64.
- Li, S., Misra, K., Matise, M. P. und Xiang, M.** (2005). Foxn4 acts synergistically with Mash1 to specify subtype identity of V2 interneurons in the spinal cord. *Proc Natl Acad Sci USA* **102**, 10688-93.
- Liem, K. F., Jr., Tremml, G., Roelink, H. und Jessell, T. M.** (1995). Dorsal differentiation of neural plate cells induced by BMP-mediated signals from epidermal ectoderm. *Cell* **82**, 969-79.
- Lindh, B. und Hökfelt, T.** (1990). Structural and functional aspects of acetylcholine peptide coexistence in the autonomic nervous system. *Prog Brain Res* **84**, 175-91.
- Liu, B., Liu, Z., Chen, T., Li, H., Qiang, B., Yuan, J., Peng, X. und Qiu, M.** (2007). Selective expression of Blhgb5 in subsets of early-born interneurons and late-born association neurons in the spinal cord. *Dev Dyn* **236**, 829-35.
- Loh, Y. P., Kim, T., Rodriguez, Y. M. und Cawley, N. X.** (2004). Secretory granule biogenesis and neuropeptide sorting to the regulated secretory pathway in neuroendocrine cells. *J Mol Neurosci* **22**, 63-71.
- Loh, Y. P., Maldonado, A., Zhang, C., Tam, W. H. und Cawley, N.** (2002). Mechanism of sorting proopiomelanocortin and proenkephalin to the regulated secretory pathway of neuroendocrine cells. *Ann NY Acad Sci* **971**, 416-25.
- Lundberg, J. M.** (1996). Pharmacology of cotransmission in the autonomic nervous system: integrative aspects on amines, neuropeptides, adenosine triphosphate, amino acids and nitric oxide. *Pharmacol Rev* **48**, 113-78.
- Lundberg, J. M., Terenius, L., Hökfelt, T. und Goldstein, M.** (1983). High levels of neuropeptide Y in peripheral noradrenergic neurons in various mammals including man. *Neurosci Lett* **42**, 167-72.
- Manzanares, M., Trainor, P. A., Ariza-McNaughton, L., Nonchev, S. und Krumlauf, R.** (2000). Dorsal patterning defects in the hindbrain, roof plate and skeleton in the dreher (dr(J)) mouse mutant. *Mech Dev* **94**, 147-56.
- McDonell, M. W., Simon, M. N. und Studier, F. W.** (1977). Analysis of restriction fragments of T7 DNA and determination of molecular weights by electrophoresis in neutral and alkaline gels. *J Mol Biol* **110**, 119-46.
- McIntire, S. L., Reimer, R. J., Schuske, K., Edwards, R. H. und Jorgensen, E. M.** (1997). Identification and characterization of the vesicular GABA transporter. *Nature* **389**, 870-6.
- Millonig, J. H., Millen, K. J. und Hatten, M. E.** (2000). The mouse Dreher gene Lmx1a controls formation of the roof plate in the vertebrate CNS. *Nature* **403**, 764-9.
- Mirnics, K. und Koerber, H. R.** (1995). Prenatal development of rat primary afferent fibers: II. Central projections. *J Comp Neurol* **355**, 601-14.

- Mizuguchi, R., Kriks, S., Cordes, R., Gossler, A., Ma, Q. und Goulding, M.** (2006). Ascl1 and Gsh1/2 control inhibitory and excitatory cell fate in spinal sensory interneurons. *Nat Neurosci* **9**, 770-8.
- Mueller, B. K.** (1999). Growth cone guidance: first steps towards a deeper understanding. *Annu Rev Neurosci* **22**, 351-88.
- Müller, T., Anlag, K., Wildner, H., Britsch, S., Treier, M. und Birchmeier, C.** (2005). The bHLH factor Olig3 coordinates the specification of dorsal neurons in the spinal cord. *Genes Dev* **19**, 733-43.
- Müller, T., Brohmann, H., Pierani, A., Heppenstall, P. A., Lewin, G. R., Jessell, T. M. und Birchmeier, C.** (2002). The homeodomain factor lbx1 distinguishes two major programs of neuronal differentiation in the dorsal spinal cord. *Neuron* **34**, 551-62.
- Muroyama, Y., Fujihara, M., Ikeya, M., Kondoh, H. und Takada, S.** (2002). Wnt signaling plays an essential role in neuronal specification of the dorsal spinal cord. *Genes Dev* **16**, 548-53.
- Nakamura, Y., Sakakibara, S., Miyata, T., Ogawa, M., Shimazaki, T., Weiss, S., Kageyama, R. und Okano, H.** (2000). The bHLH gene hes1 as a repressor of the neuronal commitment of CNS stem cells. *J Neurosci* **20**, 283-93.
- Naya, F. J., Huang, H. P., Qiu, Y., Mutoh, H., DeMayo, F. J., Leiter, A. B. und Tsai, M. J.** (1997). Diabetes, defective pancreatic morphogenesis, and abnormal enteroendocrine differentiation in BETA2/neuroD-deficient mice. *Genes Dev* **11**, 2323-34.
- Nitsche, J. F., Schuller, A. G., King, M. A., Zengh, M., Pasternak, G. W. und Pintar, J. E.** (2002). Genetic dissociation of opiate tolerance and physical dependence in delta-opioid receptor-1 and preproenkephalin knock-out mice. *J Neurosci* **22**, 10906-13.
- Olson, J. M., Asakura, A., Snider, L., Hawkes, R., Strand, A., Stoeck, J., Hallahan, A., Pritchard, J. und Tapscott, S. J.** (2001). NeuroD2 is necessary for development and survival of central nervous system neurons. *Dev Biol* **234**, 174-87.
- Otsuka, M. und Yanagisawa, M.** (1990). Pain and neurotransmitters. *Cell Mol Neurobiol* **10**, 293-302.
- Ozaki, S. und Snider, W. D.** (1997). Initial trajectories of sensory axons toward laminar targets in the developing mouse spinal cord. *J Comp Neurol* **380**, 215-29.
- Peschanski, M., Mantyh, P. W. und Besson, J. M.** (1983). Spinal afferents to the ventrobasal thalamic complex in the rat: an anatomical study using wheat-germ agglutinin conjugated to horseradish peroxidase. *Brain Res* **278**, 240-4.
- Petruschka, M., Li, S., Gilbert, T. L., Westenbroek, R. E., Bruchas, M. R., Schreiber, S., Lowe, J., Low, M. J., Pintar, J. E. und Chavkin, C.** (2007). The absence of endogenous beta-endorphin selectively blocks phosphorylation and desensitization of mu opioid receptors following partial sciatic nerve ligation. *Neuroscience* **146**, 1795-807.
- Pierani, A., Brenner-Morton, S., Chiang, C. und Jessell, T. M.** (1999). A sonic hedgehog-independent, retinoid-activated pathway of neurogenesis in the ventral spinal cord. *Cell* **97**, 903-15.

- Pillai, A., Mansouri, A., Behringer, R., Westphal, H. und Goulding, M.** (2007). Lhx1 and Lhx5 maintain the inhibitory-neurotransmitter status of interneurons in the dorsal spinal cord. *Development* **134**, 357-66.
- Polgar, E., Shehab, S. A., Watt, C. und Todd, A. J.** (1999). GABAergic neurons that contain neuropeptide Y selectively target cells with the neurokinin 1 receptor in laminae III and IV of the rat spinal cord. *J Neurosci* **19**, 2637-46.
- Pritchard, L. E. und White, A.** (2007). Neuropeptide processing and its impact on melanocortin pathways. *Endocrinology* **148**, 4201-7.
- Rexed, B.** (1952). The cytoarchitectonic organization of the spinal cord in the cat. *J Comp Neurol* **96**, 414-95.
- Rizzi, A., Marzola, G., Bigoni, R., Guerrini, R., Salvadori, S., Mogil, J. S., Regoli, D. und Calo, G.** (2001). Endogenous nociceptin signaling and stress-induced analgesia. *Neuroreport* **12**, 3009-13.
- Roelink, H.** (1996). Tripartite signaling of pattern: interactions between Hedgehogs, BMPs and Wnts in the control of vertebrate development. *Curr Opin Neurobiol* **6**, 33-40.
- Roelink, H., Porter, J. A., Chiang, C., Tanabe, Y., Chang, D. T., Beachy, P. A. und Jessell, T. M.** (1995). Floor plate and motor neuron induction by different concentrations of the amino-terminal cleavage product of sonic hedgehog autoproteolysis. *Cell* **81**, 445-55.
- Roques, B. P. und Noble, F.** (1995). Dual inhibitors of enkephalin-degrading enzymes (neutral endopeptidase 24.11 and aminopeptidase N) as potential new medications in the management of pain and opioid addiction. *NIDA Res Monogr* **147**, 104-45.
- Rouille, Y., Duguay, S. J., Lund, K., Furuta, M., Gong, Q., Lipkind, G., Oliva, A. A., Jr., Chan, S. J. und Steiner, D. F.** (1995). Proteolytic processing mechanisms in the biosynthesis of neuroendocrine peptides: the subtilisin-like proprotein convertases. *Front Neuroendocrinol* **16**, 322-61.
- Rowan, S., Todd, A. J. und Spike, R. C.** (1993). Evidence that neuropeptide Y is present in GABAergic neurons in the superficial dorsal horn of the rat spinal cord. *Neuroscience* **53**, 537-45.
- Saiki, R. K., Scharf, S., Faloona, F., Mullis, K. B., Horn, G. T., Erlich, H. A. und Arnheim, N.** (1985). Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* **230**, 1350-4.
- Sambrook, J. und Russell, D. W.** (2001). *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor, New York, USA.
- Sander, M., Paydar, S., Ericson, J., Briscoe, J., Berber, E., German, M., Jessell, T. M. und Rubenstein, J. L.** (2000). Ventral neural patterning by Nkx homeobox genes: Nkx6.1 controls somatic motor neuron and ventral interneuron fates. *Genes Dev* **14**, 2134-9.
- Sanger, F., Nicklen, S. und Coulson, A. R.** (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* **74**, 5463-7.

- Schwab, M. H., Bartholomae, A., Heimrich, B., Feldmeyer, D., Druffel-Augustin, S., Goebbels, S., Naya, F. J., Zhao, S., Frotscher, M., Tsai, M. J. et al.** (2000). Neuronal basic helix-loop-helix proteins (NEX and BETA2/Neuro D) regulate terminal granule cell differentiation in the hippocampus. *J Neurosci* **20**, 3714-24.
- Schwab, M. H., Druffel-Augustin, S., Gass, P., Jung, M., Klugmann, M., Bartholomae, A., Rossner, M. J. und Nave, K. A.** (1998). Neuronal basic helix-loop-helix proteins (NEX, neuroD, NDRF): spatiotemporal expression and targeted disruption of the NEX gene in transgenic mice. *J Neurosci* **18**, 1408-18.
- Seidah, N. G. und Chretien, M.** (1999). Proprotein and prohormone convertases: a family of subtilases generating diverse bioactive polypeptides. *Brain Res* **848**, 45-62.
- Seidah, N. G. und Prat, A.** (2002). Precursor convertases in the secretory pathway, cytosol and extracellular milieu. *Essays Biochem* **38**, 79-94.
- Shi, T. J., Zhang, X., Berge, O. G., Erickson, J. C., Palmiter, R. D. und Hökfelt, T.** (1998). Effect of peripheral axotomy on dorsal root ganglion neuron phenotype and autonomy behaviour in neuropeptide Y-deficient mice. *Regul Pept* **75-76**, 161-73.
- Shirasaki, R. und Pfaff, S. L.** (2002). Transcriptional codes and the control of neuronal identity. *Annu Rev Neurosci* **25**, 251-81.
- Sieber, M. A., Storm, R., Martinez-de-la-Torre, M., Muller, T., Wende, H., Reuter, K., Vasyutina, E. und Birchmeier, C.** (2007). Lbx1 acts as a selector gene in the fate determination of somatosensory and viscerosensory relay neurons in the hindbrain. *J Neurosci* **27**, 4902-9.
- Siegel, G. J., Agranoff, B. W., Wayne Albers, R. W., Fisher, S. K. und D., U. M.** (1999). *Basic Neurochemistry, 6th Edition*. Lippincott-Raven, Philadelphia, USA.
- Simmons, D. R., Spike, R. C. und Todd, A. J.** (1995). Galanin is contained in GABAergic neurons in the rat spinal dorsal horn. *Neurosci Lett* **187**, 119-22.
- Smyth, G. K.** (2005). Limma: linear models for microarray data. In *Bioinformatics and Computational Biology Solutions using R and Bioconductor*, (ed. R. Gentleman V. S. Carey R. Dudoit R. Irizarry und W. Huber), pp. 397-420. Springer, New York, USA.
- Snider, W. D. und McMahon, S. B.** (1998). Tackling pain at the source: new ideas about nociceptors. *Neuron* **20**, 629-32.
- Sorge, J. A.** (1988). Bacteriophage lambda cloning vectors. *Biotechnology* **10**, 43-60.
- Steiner, D. F.** (1991). Prohormone convertases revealed at last. *Curr Biol* **1**, 375-7.
- Steiner, D. F.** (1998). The proprotein convertases. *Curr Opin Chem Biol* **2**, 31-9.
- Sun, H., Xu, J., Della Penna, K. B., Benz, R. J., Kinose, F., Holder, D. J., Koblan, K. S., Gerhold, D. L. und Wang, H.** (2002). Dorsal horn-enriched genes identified by DNA microarray, in situ hybridization and immunohistochemistry. *BMC Neurosci* **3**, 11.
- Szucsik, J. C., Witte, D. P., Li, H., Pixley, S. K., Small, K. M. und Potter, S. S.** (1997). Altered forebrain and hindbrain development in mice mutant for the Gsh-2 homeobox gene. *Dev Biol* **191**, 230-42.

- Tabor, S. und Richardson, C. C.** (1987). DNA sequence analysis with a modified bacteriophage T7 DNA polymerase. *Proc Natl Acad Sci U S A* **84**, 4767-71.
- Tessier-Lavigne, M. und Goodman, C. S.** (1996). The molecular biology of axon guidance. *Science* **274**, 1123-33.
- Todd, A. J., Hughes, D. I., Polgar, E., Nagy, G. G., Mackie, M., Ottersen, O. P. und Maxwell, D. J.** (2003). The expression of vesicular glutamate transporters VGLUT1 and VGLUT2 in neurochemically defined axonal populations in the rat spinal cord with emphasis on the dorsal horn. *Eur J Neurosci* **17**, 13-27.
- Todd, A. J. und Spike, R. C.** (1993). The localization of classical transmitters and neuropeptides within neurons in laminae I-III of the mammalian spinal dorsal horn. *Prog Neurobiol* **41**, 609-45.
- Todd, A. J., Spike, R. C., Russell, G. und Johnston, H. M.** (1992). Immunohistochemical evidence that Met-enkephalin and GABA coexist in some neurones in rat dorsal horn. *Brain Res* **584**, 149-56.
- Torres, M., Gomez-Pardo, E., Dressler, G. R. und Gruss, P.** (1995). Pax-2 controls multiple steps of urogenital development. *Development* **121**, 4057-65.
- Tsuchida, T., Ensini, M., Morton, S. B., Baldassare, M., Edlund, T., Jessell, T. M. und Pfaff, S. L.** (1994). Topographic organization of embryonic motor neurons defined by expression of LIM homeobox genes. *Cell* **79**, 957-70.
- Vallstedt, A., Klos, J. M. und Ericson, J.** (2005). Multiple dorsoventral origins of oligodendrocyte generation in the spinal cord and hindbrain. *Neuron* **45**, 55-67.
- Verhage, M., Ghijssen, W. E., Nicholls, D. G. und Wiegant, V. M.** (1991a). Characterization of the release of cholecystokinin-8 from isolated nerve terminals and comparison with exocytosis of classical transmitters. *J Neurochem* **56**, 1394-400.
- Verhage, M., McMahon, H. T., Ghijssen, W. E., Boomsma, F., Scholten, G., Wiegant, V. M. und Nicholls, D. G.** (1991b). Differential release of amino acids, neuropeptides, and catecholamines from isolated nerve terminals. *Neuron* **6**, 517-24.
- Vogelstein, B. und Gillespie, D.** (1979). Preparative and analytical purification of DNA from agarose. *Proc Natl Acad Sci U S A* **76**, 615-9.
- Walters, E. T.** (1994). Injury-related behavior and neuronal plasticity: an evolutionary perspective on sensitization, hyperalgesia, and analgesia. *Int Rev Neurobiol* **36**, 325-427.
- Wang, Z., Gardell, L. R., Ossipov, M. H., Vanderah, T. W., Brennan, M. B., Hochgeschwender, U., Hruby, V. J., Malan, T. P., Jr., Lai, J. und Porreca, F.** (2001). Pronociceptive actions of dynorphin maintain chronic neuropathic pain. *J Neurosci* **21**, 1779-86.
- Wiesenfeld-Hallin, Z. und Xu, X. J.** (1998). Galanin in somatosensory function. *Ann N Y Acad Sci* **863**, 383-9.
- Wildner, H., Müller, T., Cho, S. H., Bröhl, D., Cepko, C. L., Guillemot, F. und Birchmeier, C.** (2006). dILA neurons in the dorsal spinal cord are the product of terminal and non-terminal asymmetric progenitor cell divisions, and require Mash1 for their development. *Development* **133**, 2105-13.

- Woolf, C. J. und Salter, M. W.** (2000). Neuronal plasticity: increasing the gain in pain. *Science* **288**, 1765-9.
- Wu, S. X., Goebbels, S., Nakamura, K., Nakamura, K., Kometani, K., Minato, N., Kaneko, T., Nave, K. A. und Tamamaki, N.** (2005). Pyramidal neurons of upper cortical layers generated by NEX-positive progenitor cells in the subventricular zone. *Proc Natl Acad Sci U S A* **102**, 17172-7.
- Yamada, T., Placzek, M., Tanaka, H., Dodd, J. und Jessell, T. M.** (1991). Control of cell pattern in the developing nervous system: polarizing activity of the floor plate and notochord. *Cell* **64**, 635-47.
- Zechner, D., Müller, T., Wende, H., Walther, I., Taketo, M. M., Crenshaw, E. B., 3rd, Treier, M., Birchmeier, W. und Birchmeier, C.** (2007). Bmp and Wnt/beta-catenin signals control expression of the transcription factor Olig3 and the specification of spinal cord neurons. *Dev Biol* **303**, 181-90.
- Zhao, Y., Sheng, H. Z., Amini, R., Grinberg, A., Lee, E., Huang, S., Taira, M. und Westphal, H.** (1999). Control of hippocampal morphogenesis and neuronal differentiation by the LIM homeobox gene Lhx5. *Science* **284**, 1155-8.
- Zhijin, W., Irizarry, R. A., Gentleman, R., Martinez-Murillo, F. und Spencer, F.** (2004). A Model Based Background Adjustment for Oligonucleotide Expression Arrays. *J. Am. Stat. Assoc.* **99**, 909–917.
- Zhuang, B. und Sockanathan, S.** (2006). Dorsal-ventral patterning: a view from the top. *Curr Opin Neurobiol* **16**, 20-4.
- Zigmond, R. E. und Sun, Y.** (1997). Regulation of neuropeptide expression in sympathetic neurons. Paracrine and retrograde influences. *Ann N Y Acad Sci* **814**, 181-97.