

10 Literaturverzeichnis

- Abrams, G.D., Bauer, H., und Sprinz, H. (1963):** Influence of the normal flora on mucosal morphology and cellular renewal in the ileum: A comparison of germ-free and conventional mice. *Lab. Invest.* 12, S. 355-364.
- Allison, M.J. (1989):** Characterization of the flora of the large bowel of pigs: A status report. *J. Anim. Feed Sci. Technol.* 23, S. 79-90.
- Amtsberg, G. (1984):** Die Darmflora des Schweines: Zusammensetzung und Wirkungsmechanismen. *Prakt. Tierarzt* 65, S. 1096-1111.
- Annison, G. (1992):** Commercial enzyme supplementation of wheatbased diets raises ileal glycanase activities and improves apparent metabolisable energy, starch and pentosan digestibilities in broiler chickens. *J. Anim. Feed Sci. Technol.* 38, S. 105-121.
- Armstrong, W.D., und Cline, T.R. (1976):** Effects of various dietary nutrients levels on the incidence of colibacillary diarrhoea in pig. *J. Anim. Sci.* 42, S. 592-598.
- Arthur, D. (1970):** The determination of chromium in animal feed and excreta by atomic absorption spectrometry. *Can. J. Spectrosc.* 15, S. 1-4.
- Aulrich, K., und Flachowsky, G. (1998):** Studies on the mode of action of non-starch-polysaccharides (NSP) degrading enzymes in vitro. 1. Communication: Effects on the fractions of NSP. *Arch. Anim. Nutr.* 51, S. 293-306.
- Aulrich, K., und Flachowsky, G. (2001):** Studies on the mode of action of non-starch-polysaccharides (NSP)-degrading enzymes in vitro. 2. Communication: effects on nutrient release and hydration properties. *Arch. Anim. Nutr.* 54, S. 19-32.
- Axelsson, L. (1998):** Lactic acid bacteria. Microbiology and functional aspects. 2. Aufl. New York: Marcel Dekker Inc.
- Bach Knudsen, K.E. (1997):** Carbohydrate and lignin contents of plant materials used in animal feeding. *J. Anim. Feed Sci. Technol.* 67, S. 319-338.
- Bach Knudsen, K.E. (2001a):** Development of antibiotic resistance and options to replace antimicrobials in animal diets. *Proc. Nutr. Soc.* 60, S. 291-299.
- Bach Knudsen, K.E. (2001b):** The nutritional significance of "dietary fibre" analysis. *J. Anim. Feed Sci. Technol.* 90, S. 3-20.
- Bach Knudsen, K.E., und Canibe, N. (2000):** Breakdown of plant carbohydrates in the digestive tract of pigs fed on wheat- or oat-based rolls. *J. Sci. Food Agric.* 80, S. 1253-1261.

Literaturverzeichnis

- Bach Knudsen, K.E., und Hansen, I. (1991):** Gastrointestinal implications in pigs of wheat and oat fractions. 1. Digestibility and bulking properties of polysaccharides and other major constituents. *Br. J. Nutr.* 65, S. 217-232.
- Bach Knudsen, K.E., und Johansen, H.N. (1995):** Plant carbohydrates: Enzymatic hydrolysis and microbial fermentation in the gastrointestinal tract of pigs. Book of Abstracts, Vol. 2. Wageningen: EAAP.
- Bakker, G.C., Dekker, R.A., Jongbloed, R., und Jongbloed, A.W. (1998):** Non-starch polysaccharides in pig feeding. *Vet. Q.* 20, Supplement 3, S. 59-64.
- Barrera, M., Cervantes, M., Sauer, W.C., Araiza, A.B., Torrentra, N., und Cervantes, M. (2004):** Ileal amino acid digestibility and performance of growing pigs fed wheat-based diets supplemented with xylanase. *J. Anim. Sci.* 82.
- Bartelt, J., Jadamus, A., Wiese, F., Swiech, E., Buraczewska, L., und Simon, O. (2002):** Apparent precaecal digestibility of nutrients and level of endogenous nitrogen in digesta of the small intestine of growing pigs as affected by various digesta viscosities. *Arch. Anim. Nutr.* 56, S. 93-107.
- Baur, X., Sander, I., Posch, A., und Rauf-Heimsoth, M. (1998):** Baker's asthma due to the enzyme xylanase - A new occupational allergen. *Clin. Experim. Allergy* 28, S. 1591-1593.
- Beckmann, L., Vahjen, W., und Simon, O. (2000):** Isolation of an 1,3-1,4-beta-glucan degrading Enterococcus faecium strain from the intestinal tract of chicken and partial characterization of its novel 1,3-1,4-beta-glucanase. *J. Basic. Microbiol.* 40, S. 303-310.
- Bedford, M.R. (1995):** Mechanism of action and potential environmental benefits from the use of feed enzymes. *J. Anim. Feed Sci. Technol.* 53, S. 145-155.
- Beldman, G., Osuga, D., und Whitaker, J.R. (1996):** Some characteristics of beta-[D]-xylopyranosidases, alpha-[L]-arabinofuranohydrolase from wheat bran and germinated wheat. *J. Cereal Sci.* 23, S. 169-180.
- Bergman, E.N. (1990):** Energy contributions of volatile fatty acids from the gastrointestinal tract in various species. *Physiol. Rev.* 70, S. 567-590.
- Bertschinger, H.U., und Eggenberger, E. (1978):** Evaluation of low nutrient, high fibre diets for the prevention of porcine Escherichia coli enterotoxaemia. *Vet. Microbiol.* 3, S. 281-290.
- Böhme, H. (1996):** Einsatz von Futterenzymen in der Schweinfütterung. In: W. Drochner und B. Zaccharias (Hrsg.): LAF-Informationen. Institut für Tierernährung der Bundesforschungsanstalt für Landwirtschaft Braunschweig-Völkenrode, S. 106-118.
- Champ, M., Langkilde, A.-M.B.F., Kettilitz, B., und Collet, Y.L.B. (2003):** Advances in dietary fibre characterisation. 1. Definition of dietary fibre, physiological relevance, health benefits and analytical aspects. *Nutr. Res. Rev.* 16, S. 71-82.

- Chesson, A. (1993):** Feed enzymes. *J. Anim. Feed Sci. Technol.* 45, S. 65-79.
- Choct, M. (1997):** Feed Non-Starch Polysaccharides: Chemical Structures and Nutritional Significance. *Feed Milling International, Juni-Ausgabe*, S. 13-26.
- Choct, M., und Annison, G. (1990):** Anti-nutritive activity of wheat pentosans in broiler diets. *Br. Poult. Sci.* 31, S. 811-821.
- Choct, M., und Annison, G. (1992a):** Anti-nutritive effect of wheat pentosans in broiler chickens: roles of viscosity and gut microflora. *Br. Poult. Sci.* 33, S. 821-834.
- Choct, M., und Annison, G. (1992b):** The inhibition of nutrient digestion by wheat pentosans. *Br. J. Nutr.* 67, S. 123-132.
- Christiaens, H., Leer, R.J., Pouwels, P.H., und Verstraete, W. (1992):** Cloning and expression of a conjugated bile acid hydrolase gene from *Lactobacillus plantarum* by using a direct plate assay. *Appl. Environ. Microbiol.* 58, S. 3792-3798.
- Coates, M.E., Cole, C.B., Fuller, R., Houghton, S.B., und Yokota, Y. (1981):** The gut microflora and the uptake of glucose from the small intestine of the young chick. *Br. Poult. Sci.* 22, S. 721-726.
- Cole, C.B., und Fuller, R. (1984):** Bile acid deconjugation and attachment of chicken gut bacteria: Their possible role in growth depression. *Br. Poult. Sci.* 25, S. 227-231.
- Coleman, J.P., und Hudson, L.L. (1995):** Cloning and characterization of a conjugated bile acid hydrolase gene from *Clostridium perfringens*. *Appl. Environ. Microbiol.* 61, S. 2514-2520.
- Costigan, P., und Ellis, K.J. (1987):** Analysis of faecal chromium derived from controlled release marker devices. *N. Z. J. Technol.* 3, S. 89-92.
- Damme, K., und Möbius, C. (2005):** Geflügeljahrbuch 2005; Jahrbuch des Zentralverbandes der Deutschen Geflügelwirtschaft e.V. und seiner Mitgliedsverbände. Ulmer Verlag.
- Dänicke, S., Jeroch, H., Dusel, G., Simon, O., und Bedford, M. (1999a):** Effect of dietary fat type, pentosan level and xylanase supplementation on performance and carcass characteristics of male broilers. *Arch. Geflügelkd.* 63, S. 194-203.
- Dänicke, S., Simon, O., Jeroch, H., und Bedford, M. (1997):** Interactions between dietary fat type and xylanase supplementation when rye-based diets are fed to broiler chickens 2. Performance, nutrient digestibility and the fat-soluble vitamin status of livers. *Br. Poult. Sci.* 38, S. 546-556.
- Dänicke, S., Vahjen, W., Simon, O., und Jeroch, H. (1999b):** Effects of dietary fat type and xylanase supplementation to rye-based broiler diets on selected bacterial groups adhering to the intestinal epithelium on transit time of feed and on nutrient digestibility. *Poult Sci* 78, S. 1292-1299.
- DeVries, J.W., Prosky, L., Li, B., und Cho, S. (1999):** A historical perspective on defining dietary fiber. *Cereal Foods World* 44, S. 367-369.

Dierick, N.A. (1989): Biotechnology aids to improve feed and feed digestion: enzymes and fermentation. *Arch. Anim. Nutr.* 39, S. 241-261.

Duda, D. (2004): Fermentation ausgewählter Kohlenhydrate (insbesondere Mannanoligosaccharid und Pektin) in vivo und in vitro (Hohenheimer Futterwerttest beim Schwein). Hannover: Tierärztliche Hochschule, Fachbereich Veterinärmedizin, *Dissertation*.

Durmic, Z., Pethick, D.W., Mullan, B.P., Accioly, J.M., Schulze, H., und Hampson, D.J. (2002): Evaluation of large-intestinal parameters associated with dietary treatments designed to reduce the occurrence of swine dysentery. *Br. J. Nutr.* 88, S. 159-169.

Durmic, Z., Pethick, D.W., Pluske, J.R., und Hampson, D.J. (1998): Changes in bacterial populations in the colon of pigs fed different sources of dietary fibre, and the development of swine dysentery after experimental infection. *J. Appl. Microbiol.* 85, S. 574-582.

Dusel, G. (1998): Untersuchungen zur Variabilität des Futterwertes von Weizen unter besonderer Berücksichtigung der Arabinoxylane sowie zur Wirksamkeit von Nicht-Stärke-Polysaccharid (NSP)-hydrolysierenden Enzymen in weizenbetonten Rationen beim Broiler und Ferkel. Halle-Wittenberg: Martin-Luther-Universität, *Dissertation*.

Dusel, G., Kluge, H., Glaser, K., Simon, O., Hartmann, G., Lengerken, J., und Jeroch, H. (1997): An investigation into the variability of extract viscosity of wheat-relationship with the content of non-starch-polysaccharide fractions and metabolisable energy for broiler chickens. *Arch. Anim. Nutr.* 50, S. 121-135.

Edney, M.J., Classen, H.L., und Campbell, G.L. (1986): Application of a simple radial gel diffusion assay for endo- β -glucanase activity in dietary enzyme supplements. *Poult. Sci.* 65, S. 72-77.

Elms, J., Fishwick, D., Walker, J., Rawbone, R., Jeffrey, P., Griffin, P., Gibson, M., und Curran, A.D. (2003): Prevalence of sensitisation to cellulase and xylanase in bakery workers. *Occup. Environ. Med.* 60, S. 802-804.

Feighner, S.D., und Dashkevicz, M.P. (1987): Subtherapeutic levels of antibiotics in poultry feeds and their effects on weight gain, feed efficiency and bacterial cholytaurine hydrolase activity. *Appl. Environ. Microbiol.* 53, S. 331-336.

Feighner, S.D., und Dashkevicz, M.P. (1988): Effect of dietary carbohydrates on bacterial cholytaurine hydrolase in poultry intestinal homogenates. *Appl. Environ. Microbiol.* 54, S. 337-342.

Förster, D. (2003): Zum Einsatz von Nicht-Stärke-Polysaccharid (NSP) - hydrolysierenden Enzymen in der Legehennenfütterung. *Übers. Tierernährg.* 31, S. 1-28.

Freter, R. (1992): Factors affecting the microecology of the gut. In: R. Fuller (Hrsg.): Probiotics - The scientific basis. London: Chapman & Hall, S. 111-144.

Friendship, R.M. (2003): Gastric ulcers. *Pigs News and Information* 24, S. 45-48.

- Fuller, R. (1980):** Development and dynamics of the aerobic gut flora in gnotobiotic and conventional animals. *Fortschr. Vetmed.* 33, S. 7-15.
- Fuller, R., Barrow, P.A., und Brooker, B.E. (1977):** Bacteria associated with the gastric epithelium of neonatal pigs. *Appl. Environ. Microbiol.* 35, S. 582-591.
- Gedek, B. (1984):** Leistungssteigerung und Leistungsminderung durch Mikroorganismen und deren Stoffwechselprodukte in Futtermitteln. *Kraftfutter* 6, S. 210-211.
- Gedek, B. (1986):** Probiotika in der Tierernährung. Wirkungen auf Leistung und Tiergesundheit. *Kraftfutter* 3, S. 80-84.
- Gilliland, S.E., und Speck, M.L. (1977):** Deconjugation of bile acids by intestinal lactobacilli. *Appl. Environ. Microbiol.* 33, S. 15-18.
- Graham, H., Fadel, J.G., Newman, C.W., und Newman, R.K. (1989):** Effect of pelleting and beta-glucanase supplementation on the ileal and fecal digestibility of a barley-based diet in the pig. *J. Anim. Sci.* 67, S. 1293-1298.
- Graham, H., Hesselman, K., und Aman, P. (1986):** The influence of wheat bran and sugar-beet pulp on the digestibility of dietary components in a cereal-based pig diet. *J. Nutr.* 116, S. 242-251.
- Gruppen, H., Kormelink, F.J.M., und Voragen, A.G.J. (1993):** Differences in efficacy of xylanases in the breakdown of wheat flour arabinoxylans due to their mode of action. In: C. Wenk und M. Boessinger (Hrsg.): Enzymes in Animal Nutrition. Proceedings of the 1 st Symposium Kartause Ittingen, Switzerland, 13.-16. Okt. 1993. S. 276-280.
- Haberer, B., und Schulz, E. (1998):** Zum Einfluß NSP-hydrolysierender Enzyme in der Schweinefütterung. *Übers. Tierernährg.* 26, S. 25-64.
- Haberer, B., Schulz, E., Aulrich, K., und Flachowsky, G. (1998):** Effects of beta-glucanase and xylanase supplementation in pigs fed a diet rich in nonstarch polysaccharides: composition of digesta in different prececal segments and postprandial time. *J. Anim. Physiol. Anim. Nutr.* 78, S. 84-94.
- Harris, D.L., und Kinyon, J.M. (1974):** Significance of anaerobic spirochaetes in the intestines of animals. *Am. J. Clin. Nutr.* 27, S. 1297-1304.
- Hold, G.L., Pryde, S.E., Russel, V.J., Furrie, E., und Flint, H.J. (2002):** Assessment of microbial diversity in human colonic samples by 16 S rDNA sequence analysis. *FEMS Microbiol. Ecol.* 39, S. 33-39.
- Hopwood, D.E., Pethick, D.W., und Hampson, D.J. (2002):** Increasing the viscosity of the intestinal contents stimulates proliferation of enterotoxigenic Escherichia coli and Brachyspira pilosicoli in weaner pigs. *Br. J. Nutr.* 88, S. 523-532.
- Hopwood, D.E., Pethick, D.W., Pluske, J.R., und Hampson, D.J. (2004):** Addition of pearl barley to a rice-based diet for newly weaned piglets increases the viscosity of the intestinal contents, reduces starch digestibility and exacerbates post-weaning colibacillosis. *Br. J. Nutr.* 92, S. 419-427.

Literaturverzeichnis

- Hübener, K. (2001):** Untersuchungen zur Wirkungsweise einer Xylanase im Verdauungstrakt von Masthühnern. Berlin: Freie Universität, Fachbereich Veterinärmedizin, Institut für Tierernährung, *Dissertation*.
- Hylemon, P.B. (1985):** Metabolism of bile acid in intestinal microflora. In: H. Danielson und J. Sjövall (Hrsg.): Steroids and bile acids: New comprehensive biochemistry. Amsterdam: Elsevier Publishing, S. 331-334.
- Ikegami, S., Tsuchihashi, F., Harada, H., Tsuchihashi, N., Nishide, E., und Innami, S. (1990):** Effect of viscous indigestible polysaccharides on pancreatic-biliary secretion and digestive organs in rats. *J. Nutr.* 120, S. 353-360.
- Inborr, J. (1994):** Gastrointestinal parameters influencing performance of pigs fed enzyme-supplemented barley-based feeds. Agricultural Research Centre of Finland (Scientific Agricultural Society of Finland) (Hrsg.): In: Supplementation of pig starter diets with carbohydrate-degrading enzymes - stability, activity and mode of action. Helsinki: Jokioinen, ISBN: 9517294557.
- Inborr, J., Bedford, M.R., Patience, J.F., und Classen, H.L. (1991):** The influence of supplementary feed enzymes on nutrient disappearance and digesta characteristics in the GI-tract of early-weaned pigs. In: M.W.A. Verstegen (Hrsg.): Digestive Physiology in Pigs. Proceedings of the 5th International Symposium on Digestive Physiology in Pigs, 24. bis 26. April 1991, Wageningen Netherlands. Pudoc Wageningen, EAAP-Publikation. S. 405-410.
- Jensen, B.B. (2001):** Possible ways of modifying type and amount of products from microbial fermentation in the gut. In: A. Piva, K.E. Bach-Knudsen und J.E. Lindberg (Hrsg.): Gut Environment of Pigs. Nottingham University Press, S. 181-200.
- Jeroch, H., Drochner, W., und Simon, O. (1999):** Ernährung landwirtschaftlicher Nutztiere. Stuttgart: Eugen Ulmer-Verlag bzw. UTB für Wissenschaft.
- Jin, L., Reynolds, L.P., Redmer, D.A., Caton, J.S., und Crenshaw, J.D. (1994):** Effects of dietary fiber on intestinal growth, cell proliferation, and morphology in growing pigs. *J. Anim. Sci.* 72, S. 2270-2278.
- Jørgensen, H., und Just, A. (1988):** Effect of different dietary components on site of absorption/site of disappearance of nutrients. In: L. Buraczewska, S. Buraczewski, B. Pastuszewska und T. Zebrowska (Hrsg.): Fourth International Seminar at the Institute of Animal Physiology and Nutrition. Jabłonna, Poland. S. 230-239.
- Jørgensen, H., Zhao, X.Q., und Eggum, B.O. (1996):** The influence of dietary fibre and environmental temperature on the development of the gastrointestinal tract, digestibility, degree of fermentation in the hind-gut and energy metabolism in pigs. *Br. J. Nutr.* 75, S. 365-378.
- Juarez Tomas, M.S., Bru, E., und Nader-Macias, M.E. (2003):** Comparison of the growth and hydrogen peroxide production by vaginal probiotic lactobacilli under different culture conditions. *Am. J. Obstet. Gynecol.* 188, S. 35-44.

- Kim, J.C., Simmins, P.H., Mullan, B.P., und Pluske, J.R. (2005):** The digestible energy value of wheat for pigs, with special reference to the post-weaned animal [Review]. *J. Anim. Feed Sci. Technol.* 122, S. 257-287.
- Kirchgeßner, M. (1997):** Tierernährung. 10., neubearbeitete Aufl. Frankfurt am Main: DLG-Verlags-GmbH.
- Klingler, R.W. (1995):** Grundlagen der Getreidetechnologie. Hamburg: Behr's Verlag.
- Koolman, J., und Röhm, K.-H. (1998):** Taschenatlas der Biochemie, 2. überarbeitete und erweiterte Aufl. Stuttgart: Thieme Verlag.
- Kritchevsky, D. (1988):** Dietary fiber. *Annu. Rev. Nutr.* 8, S. 301-328.
- Lee, A. (1985):** Neglected niches the microbial ecology of the gastrointestinal tract. *Adv. Microb. Ecol.* 8, S. 115-162.
- Lehninger, A.L., Nelson, D.L., und Cox, M.M. (1998):** Prinzipien der Biochemie, 2. Aufl. Heidelberg, Berlin, Oxford: Akademischer Spektrum Verlag.
- Leser, T.D., Amenuvor, J.Z., Jensen, T.K., Lindecrona, R.H., Boye, M., und Moller, K. (2002):** Culture-independent analysis of gut bacteria: The pig gastrointestinal tract microbiota revisited. *Appl. Environ. Microbiol.* 68, S. 673-690.
- March, B.E. (1979):** The host and its microflora: An ecological unit. *J. Anim. Sci.* 49, S. 857-867.
- Masuda, N. (1981):** Deconjugation of bile salts by Bacteroides and Clostridium. *Microbiol. Immunol.* 25, S. 1-11.
- McDonald, D.E., Pethick, D.W., Mullan, B.P., und Hampson, D.J. (2001):** Increasing viscosity of the intestinal contents alters small intestinal structure and intestinal growth, and stimulates proliferation of enterotoxigenic Escherichia coli in newly-weaned pigs. *Br. J. Nutr.* 86, S. 487-498.
- Merget, R., Sander, I., Raulf-Heimsoth, M., und Baur, X. (2001):** Baker's asthma due to xylanase and cellulase without sensitization to alpha-amylase and only weak sensitization to flour. *Int. Arch. Allergy Immunol.* 124, S. 502-505.
- Meyer, H., Kamphues, J., Schneider, D., und Leibetseder, J. (1999):** Suplemente zu Vorlesungen und Übungen in der Tierernährung; 9., überarbeitete Auflage. Alfeld-Hannover: Verlag M. & H. Schaper.
- Miller, A.R. (1986):** Oxidation of cell wall polysaccharide by hydrogen peroxide: A potential mechanism for cell wall breakdown in plants. *Biochemical and Biophysical Research Communications* 141, S. 238-244.
- Montagne, L., Pluske, J.R., und Hampson, D.J. (2003):** A review of interactions between dietary fibre and the intestinal mucosa, and their consequences on digestive health in young nonruminant animals. *J. Anim. Feed Sci. Technol.* 108, S. 95-117.

Literaturverzeichnis

- Mul, A.J.A., und Perry, F.G.B. (1994):** The role of fructo-oligosaccharides in animal nutrition. In: P.C. Garnsworthy und J. Wiseman (Hrsg.): Recent Developments in Pig Nutrition 3. Nottingham, United Kingdom: Nottingham University Press, S. 79-105.
- Naumann, C., Bassler, R., Seibold, R., und Barth, C. (1988):** 2. Ergänzungslieferung, Methodenbuch Band III, Die chemische Untersuchung von Futtermitteln. Darmstadt: VDLUFA-Verlag.
- Noblet, J., und Le Goff, G. (2001):** Effect of dietary fibre on the energy value of feeds for pigs. *J. Anim Feed Sci. Technol.* 90, S. 35-52.
- Palmer, N.C., und Hulland, T.J. (1965):** Factors predisposing to the development of coliform gastroenteritis in weaned pigs. *Can. Vet. J.* 6, S. 310-316.
- Pluske, J.R., Pethick, D.W., Durmic, Z., Hampson, D.J., und Mullan, B.P. (1999):** Non-starch polysaccharides in pig diets and their influence on intestinal microflora, digestive physiology and enteric disease. In: P.C. Garnsworthy und J. Wiseman (Hrsg.): Recent Developments in Pig Nutrition 3. Nottingham, United Kingdom: Nottingham University Press, S. 123-160.
- Polizeli, M.L., Rizzatti, A.C., Monti, R., Terenzi, H.F., Jorge, J.A., und Amorim, D.S. (2005):** Xylanases from fungi: properties and industrial applications. *Appl. Microbiol. Biotechnol.* 67, S. 577-591.
- Prohaszka, L. (1986):** Antibacterial mechanism of volatile fatty acids in the intestinal tract of pigs against E.coli. *J. Vet. Med.* 33, S. 166-173.
- Rerat, A. (1981):** Verdauung und Stoffwechsel von Kohlenhydraten und Proteinen im Dickdarm von Omnivoren. [Review]. *Übers. Tierernährg.* 9, S. 177-231.
- Roche (2001a):** "Ronozyme WX - Das Xylanaseenzym ihrer Wahl für Geflügel- und Schweinefutter": Roche Vitamine GmbH, 79629 Grenzach-Wyhlen; URL:<http://www.roche.de>, *Produktinformation*, [07.05.2003].
- Roche (2001b):** "Roxazyme G2 Granular - eine neue Produktform": Roche Vitamine GmbH, 79629 Grenzach-Wyhlen; URL:<http://www.roche.de>, *Produktinformation*, [07.05.2003].
- Rozee, K.R., Copper, D., Lam, K., und Costerton, J.W. (1982):** Microbial flora of the mouse ileum, mucous layer and epithelial surface. *Appl. Environ. Microbiol.* 43, S. 1451-1463.
- Savage, D.C. (1977):** Microbial ecology of the gastrointestinal tract. *Ann. Rev. Microbiol.* 31, S. 107-133.
- Savage, D.C. (1980):** The effect of stress, diet and environment on the stability of the gastrointestinal microflora. *Fortschr. Vetmed.* 33, S. 23-31.
- Savage, D.C. (1986):** Gastrointestinal Microflora in Mammalian Nutrition. *Ann. Rev. Nutr.* 6, S. 155-178.

- Savage, D.C., Dubos, R., und Schaedler, R.W. (1968):** The gastrointestinal epithelium and its autochthonus bacterial flora. *J. Exp. Med.* 127, S. 67-76.
- Schaefer, K. (1995):** Analysis of short chain fatty acids from different intestinal samples by capillary gas chromatography. *Chromatographia* 40, S. 550-556.
- Schmitz, J.F., Mc Donald, C.E., und Gilles, K.A. (1974):** Arabinoxylanases and cellulases of wheat. *Cereal Chem.* 51, S. 809-821.
- Schneeman, B.O., Richter, B.D., und Jacobs, L.R. (1982):** Response to dietary wheat bran in the exocrine pancreas and intestine of rats. *J. Nutr.* 112, S. 283-286.
- Schnippe, F. (2003):** Neue Faserträger bringen Ruhe in die Gruppe. *Top Agrar Ausgabe* 7/2003, S. 16-20.
- Schuster, S. (2003):** In-vitro Untersuchung von zwei Enzympräparaten zur Eignung als Futterzusatzstoff. Berlin: Humboldt-Universität, Landwirtschaftlich-Gärtnerische Fakultät, *Diplomarbeit*.
- Simon, O. (1998):** Auf dem Weg zu neuen Erkenntnissen über die Wirkungsweise NSP-hydrolyzierender Enzyme. In: Aktuelle Themen der Tierernährung und Veredelungswirtschaft. Vortrag auf der wissenschaftlichen Tagung vom 26. und 27. November 1997 im Hotel Seepavillon, Cuxhaven. Lohmann Animal Health GmbH & Co. KG, Heinz-Lohmann-Str. 4, 27472 Cuxhaven, S. 9-17.
- Simon, O., Hübener, K., Hirsch, K., Beckmann, L., und Vahjen, W. (2002):** Einfluß von Xylanasen auf die Darmflora. In: Aktuelle Themen der Tierernährung und Veredelungswirtschaft. Vortrag auf der wissenschaftlichen Tagung vom 21. und 22. November 2001 im Hotel Seepavillon, Cuxhaven. Lohmann Animal Health GmbH & Co. KG, Heinz-Lohmann-Str. 4, 27472 Cuxhaven, S. 68-80.
- Slade, A.M., Hoj, P.B., Morrice, N.A., und Fincher, G.B. (1989):** Purification and characterization of three (1-4)-beta-D-xylan endohydrolases from germinated barley. *Eur. J. Biochem.* 185, S. 533-539.
- Smith, H.W., und Halls, S. (1968):** The production of oedema disease and diarrhoea in weaned pigs by the oral administration of Escherichia coli: factors that influence the course of the experimental disease. *J. Med. Microbiol.* 1, S. 45-59.
- Sommer, W., und Maier-Loepke (2004):** Hohe Getreidepreise-jetzt auf Nebenprodukte umstellen?, *Schweinezucht und Schweinemast, Ausgabe: 1/2004*.
- Song, Y.-L., Kato, N., Matsumiya, Y., Liu, C.-X., Kato, H., und Watanabe, K. (1999):** Identification of and hydrogen peroxide production by fecal and vaginal Lactobacilli isolated from Japanese women and newborn infants. *J. Clin. Microbiol.* 37, S. 3062-3064.
- Steenfeldt, S. (2003):** Wheat quality-a continuing important issue in poultry nutrition-results from broiler studies in Denmark. In: 14 th European Symposium on Poultry Nutrition. August 10-14, 2003, Lillehammer, Norway. URL:<http://www.feedinfo.com>. S. 1-9.

Literaturverzeichnis

- Steenfeldt, S., und Pettersson, D. (2001):** Improvements in nutrient digestibility and performance of broiler chickens fed a wheat-and-rye based diet supplemented with enzymes. *J. Anim. Feed Sci. Technol.* 10, S. 143-157.
- Sudendey, C., und Kamphues, J. (1995):** Effekte einer Enzymzulage (alpha-Amylase, Xylanase, β -Glucanase) auf Verdauungsvorgänge im Magen-Darmtrakt von Absetzferkeln unter den Bedingungen einer forcierteren Futteraufnahme. *Proc. Soc. Nutr. Physiol.* 4, S. 108.
- Taiz, L., und Honigman, W.A. (1976):** Production of cell wall hydrolyzing enzymes by barley aleurone layers in response to gibberellic acid. *Plant Physiol.* 58, S. 380-386.
- Tannock, G.W., Fuller, R., und Pedersen, K. (1990):** Lactobacillus succession in the piglet digestive tract demonstrated by plasmid profiling. *Appl. Environ. Microbiol.* 56, S. 1310-1316.
- Teather, R.M., und Wood, P.J. (1982):** Use of congo red-polysaccharide interactions in enumeration and characterization of cellulolytic bacteria from the bovine rumen. *Appl. Environ. Microbiol.* 43, S. 777-780.
- Theander, O., Westerlund, E., Aman, P., und Graham, H. (1989):** Plant cell walls and monogastric diets. *J. Anim. Feed Sci. Technol.* 23, S. 205-225.
- Thomlinson, J.R., und Lawrence, T.L. (1981):** Dietary manipulation of gastric pH in the prophylaxis of enteric disease in weaned pigs: some field observations. *Vet. Rec.* 109, S. 120-122.
- Vahjen, W., Glaser, K., Froeck, M., und Simon, O. (1997):** Non starch polysaccharide hydrolyzing enzymes as feed additives: detection of enzyme activities and problems encountered with quantitative determination in complex samples. *Arch. Anim. Nutr.* 50, S. 331-345.
- Vahjen, W., und Simon, O. (1997a):** Mögliche Wirkungsebenen NSP-hydrolysierender Enzyme auf intestinale Mikroorganismenpopulationen bei Monogastriden. In: R. Schubert, G. Flachowsky, R. Bitsch und G. Jahreis (Hrsg.): Vitamine und Zusatzstoffe in der Ernährung von Mensch und Tier; 6. Symposium 24./25.09.1997 in Jena/Thüringen. Buch- und Kunstdruckerei Keßler GmbH, Weimar; ISBN 3-00-002381-X., S. 91-100.
- Visek, W.J. (1978):** The mode of growth promotion by antibiotics. *J. Anim. Sci.* 46, S. 1447-1469.
- Wagner, D.D., und Thomas, O.P. (1978):** Influence of diets containing rye or pectin on the intestinal flora of chicks. *Poult. Sci.* 57, S. 971-975.
- Walsh, G.A., Murphy, R.A., Killeen, G.F., Headon, D.R., und Power, R.F. (1995):** Technical note: Detection and quantification of supplemental fungal beta-glucanase activity in animal feed. *J. Anim. Sci.* 73, S. 1074-1076.

- Ward, A.T., und Marquardt, R.R. (1983):** The effect of saturation, chain length of pure triglycerides and age of bird on the utilization of rye diets. *Poult. Sci.* 62, S. 1054-1062.
- Warren, R.A. (1996):** Microbial hydrolysis of polysaccharides. *Annu. Rev. Microbiol.* 50, S. 183-212.
- Wenk, C. (2001):** The role of dietary fibre in the digestive physiology of the pig. *J. Anim. Feed Sci. Technol.* 90, S. 21-33.
- Wenk, C., und Zurcher, U. (1990):** Energetische Verwertung nahrungsfaserreicher Nebenprodukte aus der Müllerei- und Nahrungsmittelindustrie beim Schwein. *Arch. Anim. Nutr.* 40, S. 423-430.
- Wood, P.J. (1981):** The use of dye-polysaccharide interactions in β -glucanase assay. *Carbohydr. Research* 94, S. C19-C23.
- Wood, P.J., Erfle, J.D., und Teather, R.M. (1988):** Use of complex formation between congo red in detection and assay of polysaccharide hydrolases. *Methods Enzymol.* 160, S. 59-74.
- Zhu, X.Y., Zhong, T.Y., Pandya, Y., und Joerger, R.D. (2002):** 16S rRNA-based analysis of microbiota from the caecum of broiler chickens. *Appl. Environ. Microbiol.* 68, S. 124-137.
- Zoetendal, E.G., Akkermans, A.D.L., und De Vos, W.M. (1998):** Temperature gradient gel electrophoresis analysis of 16S rRNA from human faecal samples reveals stable and host specific communities of active bacteria. *Appl. Environ. Microbiol.* 64, S. 3854-3859.