

CHAPTER 13

References

1. Aasland, R. *et al.* Normalization of nomenclature for peptide motifs as ligands of modular protein domains. *FEBS Lett.* **513**, 141-144 (2002).
2. Pawson, T. New impressions of Src and Hck. *Nature* **385**, 582-3, 585 (1997).
3. Mayer, B. J. SH3 domains: complexity in moderation. *J. Cell Sci.* **114**, 1253-1263 (2001).
4. Pawson, T. & Nash, P. Protein-protein interactions define specificity in signal transduction. *Genes Dev.* **14**, 1027-1047 (2000).
5. Apic, G., Gough, J. & Teichmann, S. A. An insight into domain combinations. *Bioinformatics*. **17 Suppl 1**, S83-S89 (2001).
6. Pawson, T. & Nash, P. Assembly of cell regulatory systems through protein interaction domains. *Science* **300**, 445-452 (2003).
7. Cohen, G. B., Ren, R. & Baltimore, D. Modular binding domains in signal transduction proteins. *Cell* **80**, 237-248 (1995).
8. Copley, R. R., Doerks, T., Letunic, I. & Bork, P. Protein domain analysis in the era of complete genomes. *FEBS Lett.* **513**, 129-134 (2002).
9. Das, S. & Smith, T. F. Identifying nature's protein Lego set. *Adv. Protein Chem.* **54**, 159-183 (2000).
10. Sadowski, I., Stone, J. C. & Pawson, T. A noncatalytic domain conserved among cytoplasmic protein-tyrosine kinases modifies the kinase function and transforming activity of Fujinami sarcoma virus P130gag-fps. *Mol Cell Biol.* **6**, 4396-4408 (1986).
11. Anderson, D. *et al.* Binding of SH2 domains of phospholipase C gamma 1, GAP, and Src to activated growth factor receptors. *Science* **250**, 979-982 (1990).
12. Margolis, B. *et al.* The tyrosine phosphorylated carboxyterminus of the EGF receptor is a binding site for GAP and PLC-gamma. *EMBO J.* **9**, 4375-4380 (1990).
13. Moran, M. F. *et al.* Src homology region 2 domains direct protein-protein interactions in signal transduction. *Proc. Natl. Acad. Sci. U. S. A* **87**, 8622-8626 (1990).
14. Matsuda, M., Mayer, B. J., Fukui, Y. & Hanafusa, H. Binding of transforming protein, P47gag-crk, to a broad range of phosphotyrosine-containing proteins. *Science* **248**, 1537-1539 (1990).
15. Kuriyan, J. & Cowburn, D. Modular peptide recognition domains in eukaryotic signaling. *Annu. Rev. Biophys. Biomol. Struct.* **26**, 259-288 (1997).

16. Bradshaw, J. M. & Waksman, G. Molecular recognition by SH2 domains. *Adv. Protein Chem.* **61**, 161-210 (2002).
17. Cesareni, G., Gimona, m., Sudol, M. & Yaffe, M. Modular protein domains. Wiley (2004, whole book).
18. Cesareni, G., Panni, S., Nardelli, G. & Castagnoli, L. Can we infer peptide recognition specificity mediated by SH3 domains? *FEBS Lett.* **513**, 38-44 (2002).
19. Zarrinpar, A., Bhattacharyya, R. P. & Lim, W. A. The structure and function of proline recognition domains. *Sci. STKE*. **2003**, RE8 (2003).
20. Neduvia, V. *et al.* Systematic Discovery of New Recognition Peptides Mediating Protein Interaction Networks. *PLoS. Biol.* **3**, e405 (2005).
21. Mahoney, N. M., Janmey, P. A. & Almo, S. C. Structure of the profilin-poly-L-proline complex involved in morphogenesis and cytoskeletal regulation. *Nat. Struct. Biol.* **4**, 953-960 (1997).
22. Ball, L. J. *et al.* Dual epitope recognition by the VASP EVH1 domain modulates polyproline ligand specificity and binding affinity. *EMBO J.* **19**, 4903-4914 (2000).
23. Freund, C. *et al.* Dynamic interaction of CD2 with the GYF and the SH3 domain of compartmentalized effector molecules. *EMBO J.* **21**, 5985-5995 (2002).
24. Garrus, J. E. *et al.* Tsg101 and the vacuolar protein sorting pathway are essential for HIV-1 budding. *Cell* **107**, 55-65 (2001).
25. Otte, L. *et al.* WW domain sequence activity relationships identified using ligand recognition propensities of 42 WW domains. *Protein Sci.* **12**, 491-500 (2003).
26. Vaynberg, J. *et al.* Structure of an ultraweak protein-protein complex and its crucial role in regulation of cell morphology and motility. *Mol Cell* **17**, 513-523 (2005).
27. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool. *J. Mol Biol.* **215**, 403-410 (1990).
28. Hubbard, T. *et al.* Ensembl 2005. *Nucleic Acids Res.* **33**, D447-D453 (2005).
29. Liu, B. A. *et al.* The human and mouse complement of SH2 domain proteins-establishing the boundaries of phosphotyrosine signaling. *Mol. Cell* **22**, 851-868 (2006).
30. Schultz, J., Milpetz, F., Bork, P. & Ponting, C. P. SMART, a simple modular architecture research tool: identification of signaling domains. *Proc. Natl. Acad. Sci. U. S. A* **95**, 5857-5864 (1998).
31. Peri, S. *et al.* Development of human protein reference database as an initial platform for approaching systems biology in humans. *Genome Res.* **13**, 2363-2371 (2003).
32. Rhee, S. Y. *et al.* The Arabidopsis Information Resource (TAIR): a model organism database providing a centralized, curated gateway to Arabidopsis biology, research materials and community. *Nucleic Acids Res.* **31**, 224-228 (2003).

33. Welchman, R. L., Gordon, C. & Mayer, R. J. Ubiquitin and ubiquitin-like proteins as multifunctional signals. *Nat. Rev. Mol Cell Biol.* **6**, 599-609 (2005).
34. Lee, C. H., Saksela, K., Mirza, U. A., Chait, B. T. & Kuriyan, J. Crystal structure of the conserved core of HIV-1 Nef complexed with a Src family SH3 domain. *Cell* **85**, 931-942 (1996).
35. Ghose, R., Shekhtman, A., Goger, M. J., Ji, H. & Cowburn, D. A novel, specific interaction involving the Csk SH3 domain and its natural ligand. *Nat. Struct. Biol.* **8**, 998-1004 (2001).
36. Kami, K., Takeya, R., Sumimoto, H. & Kohda, D. Diverse recognition of non-PxxP peptide ligands by the SH3 domains from p67(phox), Grb2 and Pex13p. *EMBO J.* **21**, 4268-4276 (2002).
37. Karkkainen, S. *et al.* Identification of preferred protein interactions by phage-display of the human Src homology-3 proteome. *EMBO Rep.* (2005).
38. Ladbury, J. E. & Arold, S. Searching for specificity in SH domains. *Chem. Biol.* **7**, R3-R8 (2000).
39. Li, S. S. Specificity and versatility of SH3 and other proline-recognition domains: structural basis and implications for cellular signal transduction. *Biochem. J.* **390**, 641-653 (2005).
40. Landgraf, C. *et al.* Protein interaction networks by proteome peptide scanning. *PLoS. Biol.* **2**, E14 (2004).
41. Zarrinpar, A., Park, S. H. & Lim, W. A. Optimization of specificity in a cellular protein interaction network by negative selection. *Nature* **426**, 676-680 (2003).
42. Nishizawa, K., Freund, C., Li, J., Wagner, G. & Reinherz, E. L. Identification of a proline-binding motif regulating CD2-triggered T lymphocyte activation. *Proc. Natl. Acad. Sci. U. S. A* **95**, 14897-14902 (1998).
43. Freund, C., Dotsch, V., Nishizawa, K., Reinherz, E. L. & Wagner, G. The GYF domain is a novel structural fold that is involved in lymphoid signaling through proline-rich sequences. *Nat. Struct. Biol.* **6**, 656-660 (1999).
44. Rickles, R. J. *et al.* Identification of Src, Fyn, Lyn, PI3K and Abl SH3 domain ligands using phage display libraries. *EMBO J.* **13**, 5598-5604 (1994).
45. Sparks, A. B. *et al.* Distinct ligand preferences of Src homology 3 domains from Src, Yes, Abl, Cortactin, p53bp2, PLCgamma, Crk, and Grb2. *Proc. Natl. Acad. Sci. U. S. A* **93**, 1540-1544 (1996).
46. Tong, A. H. *et al.* A combined experimental and computational strategy to define protein interaction networks for peptide recognition modules. *Science* **295**, 321-324 (2002).
47. Mongiovi, A. M. *et al.* A novel peptide-SH3 interaction. *EMBO J.* **18**, 5300-5309 (1999).

48. Chen, H. I. & Sudol, M. The WW domain of Yes-associated protein binds a proline-rich ligand that differs from the consensus established for Src homology 3-binding modules. *Proc. Natl. Acad. Sci. U. S. A* **92**, 7819-7823 (1995).
49. Pires, J. R. *et al.* Solution structures of the YAP65 WW domain and the variant L30 K in complex with the peptides GTPPPPYTVG, N-(n-octyl)-GPPPY and PLPPY and the application of peptide libraries reveal a minimal binding epitope. *J. Mol Biol.* **314**, 1147-1156 (2001).
50. Bedford, M. T., Sarbassova, D., Xu, J., Leder, P. & Yaffe, M. B. A novel pro-Arg motif recognized by WW domains. *J. Biol. Chem.* **275**, 10359-10369 (2000).
51. Bedford, M. T., Chan, D. C. & Leder, P. FBP WW domains and the Abl SH3 domain bind to a specific class of proline-rich ligands. *EMBO J.* **16**, 2376-2383 (1997).
52. Yaffe, M. B. *et al.* Sequence-specific and phosphorylation-dependent proline isomerization: a potential mitotic regulatory mechanism. *Science* **278**, 1957-1960 (1997).
53. Lu, P. J., Zhou, X. Z., Shen, M. & Lu, K. P. Function of WW domains as phosphoserine- or phosphothreonine-binding modules. *Science* **283**, 1325-1328 (1999).
54. Niebuhr, K. *et al.* A novel proline-rich motif present in ActA of *Listeria monocytogenes* and cytoskeletal proteins is the ligand for the EVH1 domain, a protein module present in the Ena/VASP family. *EMBO J.* **16**, 5433-5444 (1997).
55. Tu, J. C. *et al.* Homer binds a novel proline-rich motif and links group 1 metabotropic glutamate receptors with IP₃ receptors. *Neuron* **21**, 717-726 (1998).
56. Volkman, B. F., Prehoda, K. E., Scott, J. A., Peterson, F. C. & Lim, W. A. Structure of the N-WASP EVH1 domain-WIP complex: insight into the molecular basis of Wiskott-Aldrich Syndrome. *Cell* **111**, 565-576 (2002).
57. Carlsson, L., Nystrom, L. E., Sundkvist, I., Markey, F. & Lindberg, U. Actin polymerizability is influenced by profilin, a low molecular weight protein in non-muscle cells. *J. Mol Biol.* **115**, 465-483 (1977).
58. Mahoney, N. M., Rozwarski, D. A., Fedorov, E., Fedorov, A. A. & Almo, S. C. Profilin binds proline-rich ligands in two distinct amide backbone orientations. *Nat. Struct. Biol.* **6**, 666-671 (1999).
59. Myllyharju, J. & Kivirikko, K. I. Identification of a novel proline-rich peptide-binding domain in prolyl 4-hydroxylase. *EMBO J.* **18**, 306-312 (1999).
60. Pornillos, O., Alam, S. L., Davis, D. R. & Sundquist, W. I. Structure of the Tsg101 UEV domain in complex with the PTAP motif of the HIV-1 p6 protein. *Nat. Struct. Biol.* **9**, 812-817 (2002).
61. Gilbert, W. Why genes in pieces? *Nature* **271**, 501 (1978).
62. Roy, S. W. Recent evidence for the exon theory of genes. *Genetica* **118**, 251-266 (2003).
63. Patthy, L. Genome evolution and the evolution of exon-shuffling--a review. *Gene* **238**, 103-114 (1999).

64. Kolkman, J. A. & Stemmer, W. P. Directed evolution of proteins by exon shuffling. *Nat. Biotechnol.* **19**, 423-428 (2001).
65. Liu, M. & Grigoriev, A. Protein domains correlate strongly with exons in multiple eukaryotic genomes--evidence of exon shuffling? *Trends Genet.* **20**, 399-403 (2004).
66. Liu, M., Wu, S., Walch, H. & Grigoriev, A. Exon-domain correlation and its corollaries. *Bioinformatics.* **21**, 3213-3216 (2005).
67. Dibb, N. J. & Newman, A. J. Evidence that introns arose at proto-splice sites. *EMBO J.* **8**, 2015-2021 (1989).
68. Perler, F. *et al.* The evolution of genes: the chicken preproinsulin gene. *Cell* **20**, 555-566 (1980).
69. Lonberg, N. & Gilbert, W. Intron/exon structure of the chicken pyruvate kinase gene. *Cell* **40**, 81-90 (1985).
70. Seed, B. & Aruffo, A. Molecular cloning of the CD2 antigen, the T-cell erythrocyte receptor, by a rapid immunoselection procedure. *Proc. Natl. Acad. Sci. U. S. A* **84**, 3365-3369 (1987).
71. Dustin, M. L. *et al.* A novel adaptor protein orchestrates receptor patterning and cytoskeletal polarity in T-cell contacts. *Cell* **94**, 667-677 (1998).
72. Hutchings, N. J., Clarkson, N., Chalkley, R., Barclay, A. N. & Brown, M. H. Linking the T cell surface protein CD2 to the actin-capping protein CAPZ via CMS and CIN85. *J. Biol. Chem.* **278**, 22396-22403 (2003).
73. Li, J. *et al.* A cdc15-like adaptor protein (CD2BP1) interacts with the CD2 cytoplasmic domain and regulates CD2-triggered adhesion. *EMBO J.* **17**, 7320-7336 (1998).
74. Tibaldi, E. V. & Reinherz, E. L. CD2BP3, CIN85 and the structurally related adaptor protein CMS bind to the same CD2 cytoplasmic segment, but elicit divergent functional activities. *Int. Immunol.* **15**, 313-329 (2003).
75. Lin, H. *et al.* Association of p59(fyn) with the T lymphocyte costimulatory receptor CD2. Binding of the Fyn Src homology (SH) 3 domain is regulated by the Fyn SH2 domain. *J. Biol. Chem.* **273**, 19914-19921 (1998).
76. Holdorf, A. D. *et al.* Proline residues in CD28 and the Src homology (SH)3 domain of Lck are required for T cell costimulation. *J. Exp. Med.* **190**, 375-384 (1999).
77. Xu, W., Doshi, A., Lei, M., Eck, M. J. & Harrison, S. C. Crystal structures of c-Src reveal features of its autoinhibitory mechanism. *Mol. Cell* **3**, 629-638 (1999).
78. Sicheri, F., Moarefi, I. & Kuriyan, J. Crystal structure of the Src family tyrosine kinase Hck. *Nature* **385**, 602-609 (1997).
79. Dikic, I. CIN85/CMS family of adaptor molecules. *FEBS Lett.* **529**, 110-115 (2002).

80. Kowanetz, K. *et al.* Identification of a novel proline-arginine motif involved in CIN85-dependent clustering of Cbl and down-regulation of epidermal growth factor receptors. *J. Biol. Chem.* **278**, 39735-39746 (2003).
81. Kurakin, A. V., Wu, S. & Bredesen, D. E. Atypical recognition consensus of CIN85/SETA/Ruk SH3 domains revealed by target-assisted iterative screening. *J. Biol. Chem.* **278**, 34102-34109 (2003).
82. Williamson, M. P. The structure and function of proline-rich regions in proteins. *Biochem. J.* **297** (Pt 2), 249-260 (1994).
83. COWAN, P. M., MCGAVIN, S. & NORTH, A. C. The polypeptide chain configuration of collagen. *Nature* **176**, 1062-1064 (1955).
84. Kay, B. K., Williamson, M. P. & Sudol, M. The importance of being proline: the interaction of proline-rich motifs in signaling proteins with their cognate domains. *FASEB J.* **14**, 231-241 (2000).
85. Rubin, G. M. *et al.* Comparative genomics of the eukaryotes. *Science* **287**, 2204-2215 (2000).
86. Mayer, B. J., Hamaguchi, M. & Hanafusa, H. A novel viral oncogene with structural similarity to phospholipase C. *Nature* **332**, 272-275 (1988).
87. Stahl, M. L., Ferenz, C. R., Kelleher, K. L., Kriz, R. W. & Knopf, J. L. Sequence similarity of phospholipase C with the non-catalytic region of src. *Nature* **332**, 269-272 (1988).
88. Bork, P. & Sudol, M. The WW domain: a signalling site in dystrophin? *Trends Biochem. Sci.* **19**, 531-533 (1994).
89. Sancho, E. *et al.* Role of UEV-1, an inactive variant of the E2 ubiquitin-conjugating enzymes, in vitro differentiation and cell cycle behavior of HT-29-M6 intestinal mucosecretory cells. *Mol. Cell Biol.* **18**, 576-589 (1998).
90. Pekkala, M. *et al.* The peptide-substrate-binding domain of collagen prolyl 4-hydroxylases is a tetratricopeptide repeat domain with functional aromatic residues. *J. Biol. Chem.* **279**, 52255-52261 (2004).
91. Saito, K. *et al.* The CAP-Gly domain of CYLD associates with the proline-rich sequence in NEMO/IKKgamma. *Structure. (Camb.)* **12**, 1719-1728 (2004).
92. Bednarek, A. K. *et al.* WWOX, a novel WW domain-containing protein mapping to human chromosome 16q23.3-24.1, a region frequently affected in breast cancer. *Cancer Res.* **60**, 2140-2145 (2000).
93. Wulf, G., Finn, G., Suizu, F. & Lu, K. P. Phosphorylation-specific prolyl isomerization: is there an underlying theme? *Nat. Cell Biol.* **7**, 435-441 (2005).
94. Staub, O. *et al.* WW domains of Nedd4 bind to the proline-rich PY motifs in the epithelial Na⁺ channel deleted in Liddle's syndrome. *EMBO J.* **15**, 2371-2380 (1996).
95. Buschdorf, J. P. & Stratling, W. H. A WW domain binding region in methyl-CpG-binding protein MeCP2: impact on Rett syndrome. *J. Mol Med.* **82**, 135-143 (2004).

96. Roberts, R. G., Gardner, R. J. & Bobrow, M. Searching for the 1 in 2,400,000: a review of dystrophin gene point mutations. *Hum. Mutat.* **4**, 1-11 (1994).
97. Liou, Y. C. *et al.* Role of the prolyl isomerase Pin1 in protecting against age-dependent neurodegeneration. *Nature* **424**, 556-561 (2003).
98. Lu, P. J., Wulf, G., Zhou, X. Z., Davies, P. & Lu, K. P. The prolyl isomerase Pin1 restores the function of Alzheimer-associated phosphorylated tau protein. *Nature* **399**, 784-788 (1999).
99. Passani, L. A. *et al.* Huntington's WW domain partners in Huntington's disease post-mortem brain fulfill genetic criteria for direct involvement in Huntington's disease pathogenesis. *Hum. Mol. Genet.* **9**, 2175-2182 (2000).
100. Faber, P. W. *et al.* Huntington interacts with a family of WW domain proteins. *Hum. Mol. Genet.* **7**, 1463-1474 (1998).
101. Saunier, S. *et al.* Characterization of the NPHP1 locus: mutational mechanism involved in deletions in familial juvenile nephronophthisis. *Am. J. Hum. Genet.* **66**, 778-789 (2000).
102. le Maire, A. *et al.* Solution NMR structure of the SH3 domain of human nephrocystin and analysis of a mutation-causing juvenile nephronophthisis. *Proteins* **59**, 347-355 (2005).
103. Hildebrandt, F. *et al.* A novel gene encoding an SH3 domain protein is mutated in nephronophthisis type 1. *Nat. Genet.* **17**, 149-153 (1997).
104. Derry, J. M., Ochs, H. D. & Francke, U. Isolation of a novel gene mutated in Wiskott-Aldrich syndrome. *Cell* **78**, 635-644 (1994).
105. Ball, L. J., Kuhne, R., Schneider-Mergener, J. & Oschkinat, H. Recognition of Proline-Rich Motifs by Protein-Protein-Interaction Domains. *Angew. Chem. Int. Ed Engl.* **44**, 2852-2869 (2005).
106. Yu, H. *et al.* Solution structure of the SH3 domain of Src and identification of its ligand-binding site. *Science* **258**, 1665-1668 (1992).
107. Yu, H. *et al.* Structural basis for the binding of proline-rich peptides to SH3 domains. *Cell* **76**, 933-945 (1994).
108. Musacchio, A., Saraste, M. & Wilmanns, M. High-resolution crystal structures of tyrosine kinase SH3 domains complexed with proline-rich peptides. *Nat. Struct. Biol.* **1**, 546-551 (1994).
109. Huang, X. *et al.* Structure of a WW domain containing fragment of dystrophin in complex with beta-dystroglycan. *Nat. Struct. Biol.* **7**, 634-638 (2000).
110. Prehoda, K. E., Lee, D. J. & Lim, W. A. Structure of the enabled/VASP homology 1 domain-peptide complex: a key component in the spatial control of actin assembly. *Cell* **97**, 471-480 (1999).
111. Fedorov, A. A., Fedorov, E., Gertler, F. & Almo, S. C. Structure of EVH1, a novel proline-rich ligand-binding module involved in cytoskeletal dynamics and neural function. *Nat. Struct. Biol.* **6**, 661-665 (1999).

112. Zimmermann, J. *et al.* Design of N-substituted peptomer ligands for EVH1 domains. *J. Biol. Chem.* **278**, 36810-36818 (2003).
113. Beneken, J. *et al.* Structure of the Homer EVH1 domain-peptide complex reveals a new twist in polyproline recognition. *Neuron* **26**, 143-154 (2000).
114. Barzik, M. *et al.* The N-terminal domain of Homer/Vesl is a new class II EVH1 domain. *J. Mol Biol.* **309**, 155-169 (2001).
115. Nguyen, J. T., Turck, C. W., Cohen, F. E., Zuckermann, R. N. & Lim, W. A. Exploiting the basis of proline recognition by SH3 and WW domains: design of N-substituted inhibitors. *Science* **282**, 2088-2092 (1998).
116. Hietala, R. *et al.* The peptide-substrate-binding domain of human collagen prolyl 4-hydroxylases. Backbone assignments, secondary structure, and binding of proline-rich peptides. *J. Biol. Chem.* **278**, 34966-34974 (2003).
117. Gough, J., Karplus, K., Hughey, R. & Chothia, C. Assignment of homology to genome sequences using a library of hidden Markov models that represent all proteins of known structure. *J. Mol Biol.* **313**, 903-919 (2001).
118. Murzin, A. G., Brenner, S. E., Hubbard, T. & Chothia, C. SCOP: a structural classification of proteins database for the investigation of sequences and structures. *J. Mol Biol.* **247**, 536-540 (1995).
119. Eddy, S. R. What is a hidden Markov model? *Nat. Biotechnol.* **22**, 1315-1316 (2004).
120. Nielsen, T. K., Liu, S., Luhrmann, R. & Ficner, R. The crystal structure of a binary U5 snRNP complex. (Oct 2005, **unpublished work**).
121. Freund, C. *et al.* Structural investigations of a GYF domain covalently linked to a proline-rich peptide. *J. Biomol. NMR* **27**, 143-149 (2003).
122. Pawson, T. & Linding, R. Synthetic modular systems--reverse engineering of signal transduction. *FEBS Lett.* **579**, 1808-1814 (2005).
123. Viguera, A. R., Arrondo, J. L., Musacchio, A., Saraste, M. & Serrano, L. Characterization of the interaction of natural proline-rich peptides with five different SH3 domains. *Biochemistry* **33**, 10925-10933 (1994).
124. Larson, S. M. & Davidson, A. R. The identification of conserved interactions within the SH3 domain by alignment of sequences and structures. *Protein Sci.* **9**, 2170-2180 (2000).
125. Pires, J. R. *et al.* Structural basis for APPTPPPLPP peptide recognition by the FBP11WW1 domain. *J. Mol Biol.* **348**, 399-408 (2005).
126. Nameki, N., Koshiba, S., Inoue, M., Kigawa, T. & Yokoyama, S. Solution Structure Of The Gyf Domain Of A Hypothetical Protein From Arabidopsis Thaliana. (May 2004, **unpublished work**).
127. Holm, L. & Sander, C. Mapping the protein universe. *Science* **273**, 595-603 (1996).

128. Lillie, S. H. & Brown, S. S. Suppression of a myosin defect by a kinesin-related gene. *Nature* **356**, 358-361 (1992).
129. Cullen, C. F., May, K. M., Hagan, I. M., Glover, D. M. & Ohkura, H. A new genetic method for isolating functionally interacting genes: high plo1(+) -dependent mutants and their suppressors define genes in mitotic and septation pathways in fission yeast. *Genetics* **155**, 1521-1534 (2000).
130. Pyke, K. W., Rawlings, G. A. & Gelfand, E. W. Isolation and characterization of the sheep erythrocyte receptor in man. *J. Immunol.* **115**, 211-215 (1975).
131. Meuer, S. C. *et al.* An alternative pathway of T-cell activation: a functional role for the 50 kd T11 sheep erythrocyte receptor protein. *Cell* **36**, 897-906 (1984).
132. Siliciano, R. F., Pratt, J. C., Schmidt, R. E., Ritz, J. & Reinherz, E. L. Activation of cytolytic T lymphocyte and natural killer cell function through the T11 sheep erythrocyte binding protein. *Nature* **317**, 428-430 (1985).
133. Sayre, P. H. *et al.* Molecular cloning and expression of T11 cDNAs reveal a receptor-like structure on human T lymphocytes. *Proc. Natl. Acad. Sci. U. S. A* **84**, 2941-2945 (1987).
134. Moingeon, P. *et al.* CD2-mediated adhesion facilitates T lymphocyte antigen recognition function. *Nature* **339**, 312-314 (1989).
135. Freund, C. (Sep 2005, **personal communication**).
136. Davis, S. J., Ikemizu, S., Wild, M. K. & van der Merwe, P. A. CD2 and the nature of protein interactions mediating cell-cell recognition. *Immunol. Rev.* **163**, 217-236 (1998).
137. Wang, J. H. *et al.* Structure of a heterophilic adhesion complex between the human CD2 and CD58 (LFA-3) counterreceptors. *Cell* **97**, 791-803 (1999).
138. Janeway, C. A., Travers, P., Walport, M. & Shlomchik, M. J. *Immunobiology: the immune system in health and disease*. Garland Science Publishing (2006, **whole book**).
139. Bachmann, M. F., Barner, M. & Kopf, M. CD2 sets quantitative thresholds in T cell activation. *J. Exp. Med.* **190**, 1383-1392 (1999).
140. Wingren, A. G. *et al.* Monocyte-regulated IFN-gamma production in human T cells involves CD2 signaling. *J. Immunol.* **151**, 1328-1336 (1993).
141. Gollob, J. A. & Ritz, J. CD2-CD58 interaction and the control of T-cell interleukin-12 responsiveness. Adhesion molecules link innate and acquired immunity. *Ann. N. Y. Acad. Sci.* **795**, 71-81 (1996).
142. Gollob, J. A. *et al.* Molecular interaction between CD58 and CD2 counter-receptors mediates the ability of monocytes to augment T cell activation by IL-12. *J. Immunol.* **157**, 1886-1893 (1996).
143. Tibaldi, E. V., Salgia, R. & Reinherz, E. L. CD2 molecules redistribute to the uropod during T cell scanning: implications for cellular activation and immune surveillance. *Proc. Natl. Acad. Sci. U. S. A* **99**, 7582-7587 (2002).

144. Hahn, W. C. & Bierer, B. E. Separable portions of the CD2 cytoplasmic domain involved in signaling and ligand avidity regulation. *J. Exp. Med.* **178**, 1831-1836 (1993).
145. Reem, G. H., Carding, S. & Reinherz, E. L. Lymphokine synthesis is induced in human thymocytes by activation of the CD 2 (T11) pathway. *J. Immunol.* **139**, 130-134 (1987).
146. Gassmann, M., Amrein, K. E., Flint, N. A., Schraven, B. & Burn, P. Identification of a signaling complex involving CD2, zeta chain and p59fyn in T lymphocytes. *Eur. J. Immunol.* **24**, 139-144 (1994).
147. Fukai, I., Hussey, R. E., Sunder-Plassmann, R. & Reinherz, E. L. A critical role for p59(fyn) in CD2-based signal transduction. *Eur. J. Immunol.* **30**, 3507-3515 (2000).
148. Chang, H. C. *et al.* Involvement of the PPPGHR motif in T cell activation via CD2. *J. Exp. Med.* **172**, 351-355 (1990).
149. Chang, H. C. *et al.* Dissection of the human CD2 intracellular domain. Identification of a segment required for signal transduction and interleukin 2 production. *J. Exp. Med.* **169**, 2073-2083 (1989).
150. Allard, L., Peseckis, S. M., Atherton, R. E., Berthiaume, L. & Resh, M. D. Dual myristylation and palmitylation of Src family member p59fyn affects subcellular localization. *J. Biol. Chem.* **269**, 16701-16705 (1994).
151. van't Hof, W. & Resh, M. D. Rapid plasma membrane anchoring of newly synthesized p59fyn: selective requirement for NH₂-terminal myristylation and palmitoylation at cysteine-3. *J. Cell Biol.* **136**, 1023-1035 (1997).
152. Yang, H. & Reinherz, E. L. Dynamic recruitment of human CD2 into lipid rafts. Linkage to T cell signal transduction. *J. Biol. Chem.* **276**, 18775-18785 (2001).
153. Berget, S. M., Moore, C. & Sharp, P. A. Spliced segments at the 5' terminus of adenovirus 2 late mRNA. *Proc. Natl. Acad. Sci. U. S. A* **74**, 3171-3175 (1977).
154. Witkowski, J. A. The discovery of 'split' genes: a scientific revolution. *Trends Biochem. Sci.* **13**, 110-113 (1988).
155. Darnell, J. E., Jr. Implications of RNA-RNA splicing in evolution of eukaryotic cells. *Science* **202**, 1257-1260 (1978).
156. Doolittle, R. F. Genes in pieces: were they ever together? *Nature* **272**, 581-582 (1978).
157. Crick, F. Split genes and RNA splicing. *Science* **204**, 264-271 (1979).
158. Moore, M. J. Gene expression. When the junk isn't junk. *Nature* **379**, 402-403 (1996).
159. Weber, M. J. New human and mouse microRNA genes found by homology search. *FEBS J.* **272**, 59-73 (2005).
160. Delahodde, A. *et al.* Site-specific DNA endonuclease and RNA maturase activities of two homologous intron-encoded proteins from yeast mitochondria. *Cell* **56**, 431-441 (1989).

161. Lopez, A. J. Alternative splicing of pre-mRNA: developmental consequences and mechanisms of regulation. *Annu. Rev. Genet.* **32**, 279-305 (1998).
162. Maniatis, T. & Tasic, B. Alternative pre-mRNA splicing and proteome expansion in metazoans. *Nature* **418**, 236-243 (2002).
163. Sorek, R. & Amitai, M. Piecing together the significance of splicing. *Nat. Biotechnol.* **19**, 196 (2001).
164. Modrek, B. & Lee, C. A genomic view of alternative splicing. *Nat. Genet.* **30**, 13-19 (2002).
165. Black, D. L. Protein diversity from alternative splicing: a challenge for bioinformatics and post-genome biology. *Cell* **103**, 367-370 (2000).
166. Graveley, B. R. Alternative splicing: increasing diversity in the proteomic world. *Trends Genet.* **17**, 100-107 (2001).
167. Senapathy, P., Shapiro, M. B. & Harris, N. L. Splice junctions, branch point sites, and exons: sequence statistics, identification, and applications to genome project. *Methods Enzymol.* **183**, 252-278 (1990).
168. Lopez, P. J. & Seraphin, B. Genomic-scale quantitative analysis of yeast pre-mRNA splicing: implications for splice-site recognition. *RNA* **5**, 1135-1137 (1999).
169. Grabowski, P. J., Padgett, R. A. & Sharp, P. A. Messenger RNA splicing in vitro: an excised intervening sequence and a potential intermediate. *Cell* **37**, 415-427 (1984).
170. Padgett, R. A., Konarska, M. M., Grabowski, P. J., Hardy, S. F. & Sharp, P. A. Lariat RNA's as intermediates and products in the splicing of messenger RNA precursors. *Science* **225**, 898-903 (1984).
171. Ruskin, B., Krainer, A. R., Maniatis, T. & Green, M. R. Excision of an intact intron as a novel lariat structure during pre-mRNA splicing in vitro. *Cell* **38**, 317-331 (1984).
172. Cech, T. R. The generality of self-splicing RNA: relationship to nuclear mRNA splicing. *Cell* **44**, 207-210 (1986).
173. Kramer, A. The structure and function of proteins involved in mammalian pre-mRNA splicing. *Annu. Rev. Biochem.* **65**, 367-409 (1996).
174. Jurica, M. S. & Moore, M. J. Pre-mRNA splicing: awash in a sea of proteins. *Mol Cell* **12**, 5-14 (2003).
175. Gesteland, R., F., Cech, T. R. & Atkins, J. F. The RNA World. Cold Spring Harbor (2005, **whole book**).
176. Nilsen, T. W. RNA-RNA interactions in the spliceosome: unraveling the ties that bind. *Cell* **78**, 1-4 (1994).
177. Will, C. L. & Luhrmann, R. Spliceosomal UsnRNP biogenesis, structure and function. *Curr. Opin. Cell Biol.* **13**, 290-301 (2001).
178. Guthrie, C. & Patterson, B. Spliceosomal snRNAs. *Annu. Rev. Genet.* **22**, 387-419 (1988).

179. Kambach, C., Walke, S. & Nagai, K. Structure and assembly of the spliceosomal small nuclear ribonucleoprotein particles. *Curr. Opin. Struct. Biol.* **9**, 222-230 (1999).
180. Stark, H., Dube, P., Luehrmann, R. & Kastner, B. Arrangement of RNA and proteins in the spliceosomal U1 small nuclear ribonucleoprotein particle. *Nature* **409**, 539-542 (2001).
181. Seraphin, B. & Rosbash, M. Identification of functional U1 snRNA-pre-mRNA complexes committed to spliceosome assembly and splicing. *Cell* **59**, 349-358 (1989).
182. Seraphin, B. & Rosbash, M. The yeast branchpoint sequence is not required for the formation of a stable U1 snRNA-pre-mRNA complex and is recognized in the absence of U2 snRNA. *EMBO J.* **10**, 1209-1216 (1991).
183. Michaud, S. & Reed, R. An ATP-independent complex commits pre-mRNA to the mammalian spliceosome assembly pathway. *Genes Dev.* **5**, 2534-2546 (1991).
184. Makarov, E. M. *et al.* Small nuclear ribonucleoprotein remodeling during catalytic activation of the spliceosome. *Science* **298**, 2205-2208 (2002).
185. Makarova, O. V. *et al.* A subset of human 35S U5 proteins, including Prp19, function prior to catalytic step 1 of splicing. *EMBO J.* **23**, 2381-2391 (2004).
186. Rosbash, M. & Seraphin, B. Who's on first? The U1 snRNP-5' splice site interaction and splicing. *Trends Biochem. Sci.* **16**, 187-190 (1991).
187. Puig, O., Gottschalk, A., Fabrizio, P. & Seraphin, B. Interaction of the U1 snRNP with nonconserved intronic sequences affects 5' splice site selection. *Genes Dev.* **13**, 569-580 (1999).
188. Zhang, D. & Rosbash, M. Identification of eight proteins that cross-link to pre-mRNA in the yeast commitment complex. *Genes Dev.* **13**, 581-592 (1999).
189. Query, C. C., Moore, M. J. & Sharp, P. A. Branch nucleophile selection in pre-mRNA splicing: evidence for the bulged duplex model. *Genes Dev.* **8**, 587-597 (1994).
190. Liu, Z. *et al.* Structural basis for recognition of the intron branch site RNA by splicing factor 1. *Science* **294**, 1098-1102 (2001).
191. Pikielny, C. W., Rymond, B. C. & Rosbash, M. Electrophoresis of ribonucleoproteins reveals an ordered assembly pathway of yeast splicing complexes. *Nature* **324**, 341-345 (1986).
192. Konarska, M. M. & Sharp, P. A. Interactions between small nuclear ribonucleoprotein particles in formation of spliceosomes. *Cell* **49**, 763-774 (1987).
193. Cheng, S. C. & Abelson, J. Spliceosome assembly in yeast. *Genes Dev.* **1**, 1014-1027 (1987).
194. Konforti, B. B., Koziolkiewicz, M. J. & Konarska, M. M. Disruption of base pairing between the 5' splice site and the 5' end of U1 snRNA is required for spliceosome assembly. *Cell* **75**, 863-873 (1993).
195. Kandels-Lewis, S. & Seraphin, B. Involvement of U6 snRNA in 5' splice site selection. *Science* **262**, 2035-2039 (1993).

196. Lesser, C. F. & Guthrie, C. Mutations in U6 snRNA that alter splice site specificity: implications for the active site. *Science* **262**, 1982-1988 (1993).
197. Luukkonen, B. G. & Seraphin, B. Genetic interaction between U6 snRNA and the first intron nucleotide in *Saccharomyces cerevisiae*. *RNA* **4**, 167-180 (1998).
198. Madhani, H. D. & Guthrie, C. A novel base-pairing interaction between U2 and U6 snRNAs suggests a mechanism for the catalytic activation of the spliceosome. *Cell* **71**, 803-817 (1992).
199. Sun, J. S. & Manley, J. L. A novel U2-U6 snRNA structure is necessary for mammalian mRNA splicing. *Genes Dev.* **9**, 843-854 (1995).
200. Newman, A. & Norman, C. Mutations in yeast U5 snRNA alter the specificity of 5' splice-site cleavage. *Cell* **65**, 115-123 (1991).
201. Newman, A. J. & Norman, C. U5 snRNA interacts with exon sequences at 5' and 3' splice sites. *Cell* **68**, 743-754 (1992).
202. Sontheimer, E. J. & Steitz, J. A. The U5 and U6 small nuclear RNAs as active site components of the spliceosome. *Science* **262**, 1989-1996 (1993).
203. Newman, A. J. The role of U5 snRNP in pre-mRNA splicing. *EMBO J.* **16**, 5797-5800 (1997).
204. Staley, J. P. & Guthrie, C. Mechanical devices of the spliceosome: Motors, clocks, springs, and things. *Cell* **92**, 315-326 (1998).
205. Hertel, K. J. & Graveley, B. R. RS domains contact the pre-mRNA throughout spliceosome assembly. *Trends Biochem. Sci.* **30**, 115-118 (2005).
206. Graveley, B. R. A protein interaction domain contacts RNA in the prespliceosome. *Mol. Cell* **13**, 302-304 (2004).
207. Bedford, M. T., Reed, R. & Leder, P. WW domain-mediated interactions reveal a spliceosome-associated protein that binds a third class of proline-rich motif: the proline glycine and methionine-rich motif. *Proc. Natl. Acad. Sci. U. S. A* **95**, 10602-10607 (1998).
208. Stevens, S. W. *et al.* Composition and functional characterization of the yeast spliceosomal penta-snRNP. *Mol Cell* **9**, 31-44 (2002).
209. Bialkowska, A. & Kurlandzka, A. Proteins interacting with Lin 1p, a putative link between chromosome segregation, mRNA splicing and DNA replication in *Saccharomyces cerevisiae*. *Yeast* **19**, 1323-1333 (2002).
210. Fromont-Racine, M., Rain, J. C. & Legrain, P. Toward a functional analysis of the yeast genome through exhaustive two-hybrid screens. *Nat. Genet.* **16**, 277-282 (1997).
211. Kramer, A. & Utans, U. Three protein factors (SF1, SF3 and U2AF) function in pre-splicing complex formation in addition to snRNPs. *EMBO J.* **10**, 1503-1509 (1991).

212. Abovich, N., Liao, X. C. & Rosbash, M. The yeast MUD2 protein: an interaction with PRP11 defines a bridge between commitment complexes and U2 snRNP addition. *Genes Dev.* **8**, 843-854 (1994).
213. Staknis, D. & Reed, R. Direct interactions between pre-mRNA and six U2 small nuclear ribonucleoproteins during spliceosome assembly. *Mol Cell Biol.* **14**, 2994-3005 (1994).
214. Abovich, N. & Rosbash, M. Cross-intron bridging interactions in the yeast commitment complex are conserved in mammals. *Cell* **89**, 403-412 (1997).
215. Rain, J. C., Rafi, Z., Rhani, Z., Legrain, P. & Kramer, A. Conservation of functional domains involved in RNA binding and protein-protein interactions in human and *Saccharomyces cerevisiae* pre-mRNA splicing factor SF1. *RNA* **4**, 551-565 (1998).
216. Hartmuth, K. *et al.* Protein composition of human prespliceosomes isolated by a tobramycin affinity-selection method. *Proc. Natl. Acad. Sci. U. S. A* **99**, 16719-16724 (2002).
217. Laggerbauer, B. *et al.* The human U5 snRNP 52K protein (CD2BP2) interacts with U5-102K (hPrp6), a U4/U6.U5 tri-snRNP bridging protein, but dissociates upon tri-snRNP formation. *RNA* **11**, 598-608 (2005).
218. Beggs, J. D., Teigelkamp, S. & Newman, A. J. The role of PRP8 protein in nuclear pre-mRNA splicing in yeast. *J. Cell Sci. Suppl.* **19**, 101-105 (1995).
219. Grainger, R. J. & Beggs, J. D. Prp8 protein: at the heart of the spliceosome. *RNA* **11**, 533-557 (2005).
220. Bailis, J. M., Bernard, P., Antonelli, R., Allshire, R. C. & Forsburg, S. L. Hsk1-Dfp1 is required for heterochromatin-mediated cohesion at centromeres. *Nat. Cell Biol.* **5**, 1111-1116 (2003).
221. Giovannone, B. *et al.* Two novel proteins that are linked to insulin-like growth factor (IGF-I) receptors by the Grb10 adapter and modulate IGF-I signaling. *J. Biol. Chem.* **278**, 31564-31573 (2003).
222. Tong, A. H. *et al.* Global mapping of the yeast genetic interaction network. *Science* **303**, 808-813 (2004).
223. Bensen, E. S., Yeung, B. G. & Payne, G. S. Ric1p and the Ypt6p GTPase function in a common pathway required for localization of trans-Golgi network membrane proteins. *Mol Biol. Cell* **12**, 13-26 (2001).
224. Siniossoglou, S., Peak-Chew, S. Y. & Pelham, H. R. Ric1p and Rgp1p form a complex that catalyses nucleotide exchange on Ypt6p. *EMBO J.* **19**, 4885-4894 (2000).
225. Andoh, T., Azad, A. K., Shigematsu, A., Ohshima, Y. & Tani, T. The fission yeast ptr1+ gene involved in nuclear mRNA export encodes a putative ubiquitin ligase. *Biochem. Biophys. Res. Commun.* **317**, 1138-1143 (2004).
226. Kistler, A. L. & Guthrie, C. Deletion of MUD2, the yeast homolog of U2AF65, can bypass the requirement for sub2, an essential spliceosomal ATPase. *Genes Dev.* **15**, 42-49 (2001).

227. Strasser, K. & Hurt, E. Splicing factor Sub2p is required for nuclear mRNA export through its interaction with Yra1p. *Nature* **413**, 648-652 (2001).
228. Espejo, A., Cote, J., Bednarek, A., Richard, S. & Bedford, M. T. A protein-domain microarray identifies novel protein-protein interactions. *Biochem. J.* **367**, 697-702 (2002).
229. Hu, H. *et al.* A map of WW domain family interactions. *Proteomics*. **4**, 643-655 (2004).
230. Wiedemann, U. *et al.* Quantification of PDZ domain specificity, prediction of ligand affinity and rational design of super-binding peptides. *J. Mol Biol.* **343**, 703-718 (2004).
231. Boisguerin, P. *et al.* An improved method for the synthesis of cellulose membrane-bound peptides with free C termini is useful for PDZ domain binding studies. *Chem. Biol.* **11**, 449-459 (2004).
232. Smith, G. P. Filamentous fusion phage: novel expression vectors that display cloned antigens on the virion surface. *Science* **228**, 1315-1317 (1985).
233. Frank, R. Spot-synthesis: an easy technique for the positionally addressable, parallel chemical synthesis on a membrane support. *Tetrahedron* **48**, 9217-9232 (1992).
234. Scott, J. K. & Smith, G. P. Searching for peptide ligands with an epitope library. *Science* **249**, 386-390 (1990).
235. Sambrook, J., Fritsch, E. F. & Maniatis, T. Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, Plainview, NY (1989, **whole book**).
236. Chen, H. I. *et al.* Characterization of the WW domain of human yes-associated protein and its polyproline-containing ligands. *J. Biol. Chem.* **272**, 17070-17077 (1997).
237. Toepert, F. *et al.* Combining SPOT synthesis and native peptide ligation to create large arrays of WW protein domains. *Angew. Chem. Int. Ed Engl.* **42**, 1136-1140 (2003).
238. Weng, Z., Taylor, J. A., Turner, C. E., Brugge, J. S. & Seidel-Dugan, C. Detection of Src homology 3-binding proteins, including paxillin, in normal and v-Src-transformed Balb/c 3T3 cells. *J. Biol. Chem.* **268**, 14956-14963 (1993).
239. Rigaut, G. *et al.* A generic protein purification method for protein complex characterization and proteome exploration. *Nat. Biotechnol.* **17**, 1030-1032 (1999).
240. Knuesel, M. *et al.* Identification of novel protein-protein interactions using a versatile mammalian tandem affinity purification expression system. *Mol. Cell Proteomis.* **2**, 1225-1233 (2003).
241. Zhou, Z., Licklider, L. J., Gygi, S. P. & Reed, R. Comprehensive proteomic analysis of the human spliceosome. *Nature* **419**, 182-185 (2002).
242. Gavin, A. C. *et al.* Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature* **415**, 141-147 (2002).
243. Ho, Y. *et al.* Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. *Nature* **415**, 180-183 (2002).

244. Fields, S. & Song, O. A novel genetic system to detect protein-protein interactions. *Nature* **340**, 245-246 (1989).
245. Serrano, M., Hannon, G. J. & Beach, D. A new regulatory motif in cell-cycle control causing specific inhibition of cyclin D/CDK4. *Nature* **366**, 704-707 (1993).
246. Finley, R. L., Jr. & Brent, R. Interaction mating reveals binary and ternary connections between Drosophila cell cycle regulators. *Proc. Natl. Acad. Sci. U. S. A* **91**, 12980-12984 (1994).
247. Allen, J. B., Walberg, M. W., Edwards, M. C. & Elledge, S. J. Finding prospective partners in the library: the two-hybrid system and phage display find a match. *Trends Biochem. Sci.* **20**, 511-516 (1995).
248. Ito, T. *et al.* A comprehensive two-hybrid analysis to explore the yeast protein interactome. *Proc. Natl. Acad. Sci. U. S. A* **98**, 4569-4574 (2001).
249. Giot, L. *et al.* A protein interaction map of Drosophila melanogaster. *Science* **302**, 1727-1736 (2003).
250. Rual, J. F. *et al.* Towards a proteome-scale map of the human protein-protein interaction network. *Nature* **437**, 1173-1178 (2005).
251. Stelzl, U. *et al.* A human protein-protein interaction network: a resource for annotating the proteome. *Cell* **122**, 957-968 (2005).
252. Macgregor, P. F., Abate, C. & Curran, T. Direct cloning of leucine zipper proteins: Jun binds cooperatively to the CRE with CRE-BP1. *Oncogene* **5**, 451-458 (1990).
253. Zhu, H. *et al.* Global analysis of protein activities using proteome chips. *Science* **293**, 2101-2105 (2001).
254. Mosser, E. A. *et al.* Physical and functional interactions between the transactivation domain of the hematopoietic transcription factor NF-E2 and WW domains. *Biochemistry* **37**, 13686-13695 (1998).
255. Tommasi di Vigiano, A., Di Zenzo, G., Sudol, M., Cesareni, G. & Dente, L. Contribution of the different modules in the utrophin carboxy-terminal region to the formation and regulation of the DAP complex. *FEBS Lett.* **471**, 229-234 (2000).
256. Dalby, P. A., Hoess, R. H. & DeGrado, W. F. Evolution of binding affinity in a WW domain probed by phage display. *Protein Sci.* **9**, 2366-2376 (2000).
257. Kasanov, J., Pirozzi, G., Uveges, A. J. & Kay, B. K. Characterizing Class I WW domains defines key specificity determinants and generates mutant domains with novel specificities. *Chem. Biol.* **8**, 231-241 (2001).
258. Gram, H., Schmitz, R., Zuber, J. F. & Baumann, G. Identification of phosphopeptide ligands for the Src-homology 2 (SH2) domain of Grb2 by phage display. *Eur. J. Biochem.* **246**, 633-637 (1997).
259. Schmitz, R., Baumann, G. & Gram, H. Catalytic specificity of phosphotyrosine kinases Blk, Lyn, c-Src and Syk as assessed by phage display. *J. Mol Biol.* **260**, 664-677 (1996).

260. Cochrane, D., Webster, C., Masih, G. & McCafferty, J. Identification of natural ligands for SH2 domains from a phage display cDNA library. *J. Mol Biol.* **297**, 89-97 (2000).
261. Fuh, G. *et al.* Analysis of PDZ domain-ligand interactions using carboxyl-terminal phage display. *J. Biol. Chem.* **275**, 21486-21491 (2000).
262. Marvin, D. A. Filamentous phage structure, infection and assembly. *Curr. Opin. Struct. Biol.* **8**, 150-158 (1998).
263. Roth, T. A., Weiss, G. A., Eigenbrot, C. & Sidhu, S. S. A minimized M13 coat protein defines the requirements for assembly into the bacteriophage particle. *J. Mol Biol.* **322**, 357-367 (2002).
264. Rodi, D. J. & Makowski, L. Phage-display technology--finding a needle in a vast molecular haystack. *Curr. Opin. Biotechnol.* **10**, 87-93 (1999).
265. Mullaney, B. P. & Pallavicini, M. G. Protein-protein interactions in hematology and phage display. *Exp. Hematol.* **29**, 1136-1146 (2001).
266. Smothers, J. F., Henikoff, S. & Carter, P. Tech.Sight. Phage display. Affinity selection from biological libraries. *Science* **298**, 621-622 (2002).
267. Hoogenboom, H. R. Overview of antibody phage-display technology and its applications. *Methods Mol Biol.* **178**, 1-37 (2002).
268. Mattheakis, L. C., Bhatt, R. R. & Dower, W. J. An in vitro polysome display system for identifying ligands from very large peptide libraries. *Proc. Natl. Acad. Sci. U. S. A* **91**, 9022-9026 (1994).
269. Roberts, R. W. & Szostak, J. W. RNA-peptide fusions for the in vitro selection of peptides and proteins. *Proc. Natl. Acad. Sci. U. S. A* **94**, 12297-12302 (1997).
270. Nemoto, N., Miyamoto-Sato, E., Husimi, Y. & Yanagawa, H. In vitro virus: bonding of mRNA bearing puromycin at the 3'-terminal end to the C-terminal end of its encoded protein on the ribosome in vitro. *FEBS Lett.* **414**, 405-408 (1997).
271. Amstutz, P., Forrer, P., Zahnd, C. & Pluckthun, A. In vitro display technologies: novel developments and applications. *Curr. Opin. Biotechnol.* **12**, 400-405 (2001).
272. Frank, R. The SPOT-synthesis technique. Synthetic peptide arrays on membrane supports--principles and applications. *J. Immunol. Methods* **267**, 13-26 (2002).
273. Songyang, Z. *et al.* SH2 domains recognize specific phosphopeptide sequences. *Cell* **72**, 767-778 (1993).
274. Yaffe, M. B. *et al.* The structural basis for 14-3-3:phosphopeptide binding specificity. *Cell* **91**, 961-971 (1997).
275. Santamaria, F., Wu, Z., Boulegue, C., Pal, G. & Lu, W. Reexamination of the recognition preference of the specificity pocket of the Abl SH3 domain. *J. Mol Recognit.* **16**, 131-138 (2003).

276. Chan, W. & White, P. Fmoc Solid Phase Peptide Synthesis: A Practical Approach. Oxford University Press (1999, **whole book**).
277. Edman, P. & Begg, G. A protein sequenator. *Eur. J. Biochem.* **1**, 80-91 (1967).
278. Lam, K. S. *et al.* A new type of synthetic peptide library for identifying ligand-binding activity. *Nature* **354**, 82-84 (1991).
279. Chen, J. K., LANE, W. S., Brauer, A. W., TANAKA, A. & Schreiber, S. Biased combinatorial Libraries- Novel Ligands for the SH3 domain of Phosphatidylinositol 3-Kinase. *Journal of the American Chemical Society* **115**, 12591-12592 (1993).
280. Lebl, M. *et al.* One-bead-one-structure combinatorial libraries. *Biopolymers* **37**, 177-198 (1995).
281. Beebe, K. D., Wang, P., Arabaci, G. & Pei, D. Determination of the binding specificity of the SH2 domains of protein tyrosine phosphatase SHP-1 through the screening of a combinatorial phosphotyrosyl peptide library. *Biochemistry* **39**, 13251-13260 (2000).
282. Frank, R. High-density synthetic peptide microarrays: emerging tools for functional genomics and proteomics. *Comb. Chem. High Throughput. Screen.* **5**, 429-440 (2002).
283. Frank, R. & Overwin, H. SPOT synthesis. Epitope analysis with arrays of synthetic peptides prepared on cellulose membranes. *Methods Mol Biol.* **66**, 149-169 (1996).
284. Fodor, S. P. *et al.* Light-directed, spatially addressable parallel chemical synthesis. *Science* **251**, 767-773 (1991).
285. Ermekova, K. S. *et al.* The WW domain of neural protein FE65 interacts with proline-rich motifs in Mena, the mammalian homolog of Drosophila enabled. *J. Biol. Chem.* **272**, 32869-32877 (1997).
286. Drees, B. *et al.* Characterization of the interaction between zyxin and members of the Ena/vasodilator-stimulated phosphoprotein family of proteins. *J. Biol. Chem.* **275**, 22503-22511 (2000).
287. Berry, D. M., Nash, P., Liu, S. K., Pawson, T. & McGlade, C. J. A high-affinity Arg-X-X-Lys SH3 binding motif confers specificity for the interaction between Gads and SLP-76 in T cell signaling. *Curr. Biol.* **12**, 1336-1341 (2002).
288. Cestra, G. *et al.* The SH3 domains of endophilin and amphiphysin bind to the proline-rich region of synaptojanin 1 at distinct sites that display an unconventional binding specificity. *J. Biol. Chem.* **274**, 32001-32007 (1999).
289. Rodriguez, M., Li, S. S., Harper, J. W. & Songyang, Z. An oriented peptide array library (OPAL) strategy to study protein-protein interactions. *J. Biol. Chem.* **279**, 8802-8807 (2004).
290. Cavanagh, J., Fairbrother, W., Palmer III, A. & Skelton N. PROTEIN NMR SPECTROSCOPY:Principles and Practice. Elsevier (2005, **whole book**).
291. Yu, H. Extending the size limit of protein nuclear magnetic resonance. *Proc. Natl. Acad. Sci. U. S. A* **96**, 332-334 (1999).

292. Neumann, E., Schaefer-Ridder, M., Wang, Y. & Hofsneider, P. H. Gene transfer into mouse lyoma cells by electroporation in high electric fields. *EMBO J.* **1**, 841-845 (1982).
293. Taketo, A. DNA transfection of Escherichia coli by electroporation. *Biochim. Biophys. Acta* **949**, 318-324 (1988).
294. Cohen, S. N., Chang, A. C. & Hsu, L. Nonchromosomal antibiotic resistance in bacteria: genetic transformation of Escherichia coli by R-factor DNA. *Proc. Natl. Acad. Sci. U. S. A* **69**, 2110-2114 (1972).
295. Freund, C., Ross, A., Guth, B., Pluckthun, A. & Holak, T. A. Characterization of the linker peptide of the single-chain Fv fragment of an antibody by NMR spectroscopy. *FEBS Lett.* **320**, 97-100 (1993).
296. Zech, T. Functional studies of the adaptor protein CD2BP2.(Nov 2003, **Diplomarbeit**).
297. Felici, F., Castagnoli, L., Musacchio, A., Jappelli, R. & Cesareni, G. Selection of antibody ligands from a large library of oligopeptides expressed on a multivalent exposition vector. *J. Mol. Biol.* **222**, 301-310 (1991).
298. Kramer, A. & Schneider-Mergener, J. Synthesis and screening of peptide libraries on continuous cellulose membrane supports. *Methods Mol. Biol.* **87**, 25-39 (1998).
299. Kramer, A. *et al.* Spot synthesis: observations and optimizations. *Journal of Peptide Research* **54**, 319-327 (1999).
300. Heuer, K., Kofler, M., Langdon, G., Thiemke, K. & Freund, C. Structure of a helically extended SH3 domain of the T cell adapter protein ADAP. *Structure (Camb.)* **12**, 603-610 (2004).
301. Yamamoto, K. R., Alberts, B. M., Benzinger, R., Lawhorne, L. & Treiber, G. Rapid bacteriophage sedimentation in the presence of polyethylene glycol and its application to large-scale virus purification. *Virology* **40**, 734-744 (1970).
302. Beyermann, M., Fechner, K., Furtkert, J., Krause, E. & Bienert, M. A single-point slight alteration set as a tool for structure-activity relationship studies of ovine corticotropin releasing factor. *J. Med. Chem.* **39**, 3324-3330 (1996).
303. Eccles, C., Guntert, P., Billeter, M. & Wuthrich, K. Efficient analysis of protein 2D NMR spectra using the software package EASY. *J. Biomol. NMR* **1**, 111-130 (1991).
304. Goddard, T. D. & Kneller, D. G. SPARKY. (2006, **computer program**).
305. Grzesiek, S. & Bax, A. Correlating Backbone Amide and Side-Chain Resonances in Larger Proteins by Multiple Relayed Triple Resonance Nmr. *Journal of the American Chemical Society* **114**, 6291-6293 (1992).
306. Grzesiek, S. & Bax, A. An Efficient Experiment for Sequential Backbone Assignment of Medium-Sized Isotopically Enriched Proteins. *Journal of Magnetic Resonance* **99**, 201-207 (1992).

307. Kay, L. E., Ikura, M., Tschudin, R. & Bax, A. 3-Dimensional Triple-Resonance Nmr-Spectroscopy of Isotopically Enriched Proteins. *Journal of Magnetic Resonance* **89**, 496-514 (1990).
308. Cosentino, G. P. *et al.* Eap1p, a novel eukaryotic translation initiation factor 4E-associated protein in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* **20**, 4604-4613 (2000).
309. Fan, H. Y., Hu, Y., Tudor, M. & Ma, H. Specific interactions between the K domains of AG and AGLs, members of the MADS domain family of DNA binding proteins. *Plant J.* **12**, 999-1010 (1997).
310. Koradi, R., Billeter, M. & Wuthrich, K. MOLMOL: a program for display and analysis of macromolecular structures. *J. Mol. Graph.* **14**, 51-32 (1996).
311. Aivazian, D. & Stern, L. J. Phosphorylation of T cell receptor zeta is regulated by a lipid dependent folding transition. *Nat. Struct. Biol.* **7**, 1023-1026 (2000).
312. Goldberg, M. B. Actin-based motility of intracellular microbial pathogens. *Microbiol. Mol Biol. Rev.* **65**, 595-626, table (2001).
313. Pornillos, O. *et al.* Structure and functional interactions of the Tsg101 UEV domain. *EMBO J.* **21**, 2397-2406 (2002).
314. Bouamr, F. *et al.* PPPYVEPTAP motif is the late domain of human T-cell leukemia virus type 1 Gag and mediates its functional interaction with cellular proteins Nedd4 and Tsg101 [corrected]. *J. Virol.* **77**, 11882-11895 (2003).
315. Timmins, J. *et al.* Ebola virus matrix protein VP40 interaction with human cellular factors Tsg101 and Nedd4. *J. Mol Biol.* **326**, 493-502 (2003).
316. Harty, R. N., Brown, M. E., Wang, G., Huibregtse, J. & Hayes, F. P. A PPxY motif within the VP40 protein of Ebola virus interacts physically and functionally with a ubiquitin ligase: implications for filovirus budding. *Proc. Natl. Acad. Sci. U. S. A* **97**, 13871-13876 (2000).
317. Beer, B. E. *et al.* Characterization of novel simian immunodeficiency viruses from red-capped mangabeys from Nigeria (SIVrcmNG409 and -NG411). *J. Virol.* **75**, 12014-12027 (2001).
318. Guyader, M. *et al.* Genome organization and transactivation of the human immunodeficiency virus type 2. *Nature* **326**, 662-669 (1987).
319. Nash, P. *et al.* Multisite phosphorylation of a CDK inhibitor sets a threshold for the onset of DNA replication. *Nature* **414**, 514-521 (2001).
320. Uetz, P. *et al.* A comprehensive analysis of protein-protein interactions in *Saccharomyces cerevisiae*. *Nature* **403**, 623-627 (2000).
321. de Hoog, C. L., Foster, L. J. & Mann, M. RNA and RNA binding proteins participate in early stages of cell spreading through spreading initiation centers. *Cell* **117**, 649-662 (2004).

322. Stevens, S. W. *et al.* Biochemical and genetic analyses of the U5, U65 and U4/U6•U5 small nuclear ribonucleoproteins from *Saccharomyces cerevisiae*. *Rna-A Publication of the Rna Society* **7**, 1543-1553 (2001).
323. Behrens, S. E. & Luhrmann, R. Immunoaffinity Purification of A [U4/U6-U5] Tri-Snrnp from Human-Cells. *Genes & Development* **5**, 1439-1452 (1991).
324. Giaever, G. *et al.* Functional profiling of the *Saccharomyces cerevisiae* genome. *Nature* **418**, 387-391 (2002).
325. Huh, W. K. *et al.* Global analysis of protein localization in budding yeast. *Nature* **425**, 686-691 (2003).
326. Llorian, M. *et al.* Nucleocytoplasmic shuttling of the splicing factor SIPP1. *J. Biol. Chem.* (2005).
327. Matsuo, R. *et al.* The yeast eIF4E-associated protein Eap1p attenuates GCN4 translation upon TOR-inactivation. *FEBS Lett.* **579**, 2433-2438 (2005).
328. Deloche, O., de la, C. J., Kressler, D., Doere, M. & Linder, P. A membrane transport defect leads to a rapid attenuation of translation initiation in *Saccharomyces cerevisiae*. *Mol. Cell* **13**, 357-366 (2004).
329. Lang, B. D. & Fridovich-Keil, J. L. Scp160p, a multiple KH-domain protein, is a component of mRNP complexes in yeast. *Nucleic Acids Res.* **28**, 1576-1584 (2000).
330. Mendelsohn, B. A. *et al.* Genetic and biochemical interactions between SCP160 and EAP1 in yeast. *Nucleic Acids Res.* **31**, 5838-5847 (2003).
331. von Der, H. T., Ball, P. D. & McCarthy, J. E. Stabilization of eukaryotic initiation factor 4E binding to the mRNA 5'-Cap by domains of eIF4G. *J. Biol. Chem.* **275**, 30551-30555 (2000).
332. Verlhac, M. H., Chen, R. H., Hanachi, P., Hershey, J. W. & Deryck, R. Identification of partners of TIF34, a component of the yeast eIF3 complex, required for cell proliferation and translation initiation. *EMBO J.* **16**, 6812-6822 (1997).
333. Naranda, T., Kainuma, M., MacMillan, S. E. & Hershey, J. W. The 39-kilodalton subunit of eukaryotic translation initiation factor 3 is essential for the complex's integrity and for cell viability in *Saccharomyces cerevisiae*. *Mol Cell Biol.* **17**, 145-153 (1997).
334. Hershey,J.W. & Merrick,W.C. Translational Control of Gene Expression. Sonenberg,N., Hershey,J.W. & Mathews,M.B. (eds.), pp. 33-88 (Cold Spring Harbor Laboratory Press,2000).
335. Valasek, L., Nielsen, K. H. & Hinnebusch, A. G. Direct eIF2-eIF3 contact in the multifactor complex is important for translation initiation in vivo. *EMBO J.* **21**, 5886-5898 (2002).
336. Hinnebusch,A.G. Translational Control of Gene Expression. Sonenberg,N., Hershey,J.W. & Mathews,M.B. (eds.), pp. 185-243 (Cold Spring Harbor Laboratory Press,2000).

337. McCutchen-Maloney, S. L. *et al.* cDNA cloning, expression, and functional characterization of PI31, a proline-rich inhibitor of the proteasome. *J. Biol. Chem.* **275**, 18557-18565 (2000).
338. Albert, T. K. *et al.* Identification of a ubiquitin-protein ligase subunit within the CCR4-NOT transcription repressor complex. *EMBO J.* **21**, 355-364 (2002).
339. Collart, M. A. Global control of gene expression in yeast by the Ccr4-Not complex. *Gene* **313**, 1-16 (2003).
340. Rutz, B. & Seraphin, B. A dual role for BBP/ScSF1 in nuclear pre-mRNA retention and splicing. *EMBO J.* **19**, 1873-1886 (2000).
341. Rousseau, D., Kaspar, R., Rosenwald, I., Gehrke, L. & Sonenberg, N. Translation initiation of ornithine decarboxylase and nucleocytoplasmic transport of cyclin D1 mRNA are increased in cells overexpressing eukaryotic initiation factor 4E. *Proc. Natl. Acad. Sci. U. S. A* **93**, 1065-1070 (1996).
342. Carnahan, R. H. *et al.* Dim1p is required for efficient splicing and export of mRNA encoding lid1p, a component of the fission yeast anaphase-promoting complex. *Eukaryot. Cell* **4**, 577-587 (2005).
343. Pornillos, O., Garrus, J. E. & Sundquist, W. I. Mechanisms of enveloped RNA virus budding. *Trends Cell Biol.* **12**, 569-579 (2002).
344. Timmins, J., Ruigrok, R. W. & Weissenhorn, W. Structural studies on the Ebola virus matrix protein VP40 indicate that matrix proteins of enveloped RNA viruses are analogues but not homologues. *FEMS Microbiol. Lett.* **233**, 179-186 (2004).
345. Gottlinger, H. G., Dorfman, T., Sodroski, J. G. & Haseltine, W. A. Effect of mutations affecting the p6 gag protein on human immunodeficiency virus particle release. *Proc. Natl. Acad. Sci. U. S. A* **88**, 3195-3199 (1991).
346. Huang, M., Orenstein, J. M., Martin, M. A. & Freed, E. O. p6Gag is required for particle production from full-length human immunodeficiency virus type 1 molecular clones expressing protease. *J. Virol.* **69**, 6810-6818 (1995).
347. Pornillos, O. *et al.* HIV Gag mimics the Tsg101-recruiting activity of the human Hrs protein. *J. Cell Biol.* **162**, 425-434 (2003).
348. Guyader, M., Emerman, M., Montagnier, L. & Peden, K. VPX mutants of HIV-2 are infectious in established cell lines but display a severe defect in peripheral blood lymphocytes. *EMBO J.* **8**, 1169-1175 (1989).
349. Fletcher, T. M., III *et al.* Nuclear import and cell cycle arrest functions of the HIV-1 Vpr protein are encoded by two separate genes in HIV-2/SIV(SM). *EMBO J.* **15**, 6155-6165 (1996).
350. Pancio, H. A., Vander, H. N. & Ratner, L. The C-terminal proline-rich tail of human immunodeficiency virus type 2 Vpx is necessary for nuclear localization of the viral preintegration complex in nondividing cells. *J. Virol.* **74**, 6162-6167 (2000).

351. Mueller, S. M., Jung, R., Weiler, S. & Lang, S. M. Vpx proteins of SIVmac239 and HIV-2ROD interact with the cytoskeletal protein alpha-actinin 1. *J. Gen. Virol.* **85**, 3291-3303 (2004).
352. Otey, C. A. & Carpen, O. Alpha-actinin revisited: a fresh look at an old player. *Cell Motil. Cytoskeleton* **58**, 104-111 (2004).
353. Macias, M. J., Gervais, V., Civera, C. & Oschkinat, H. Structural analysis of WW domains and design of a WW prototype. *Nat. Struct. Biol.* **7**, 375-379 (2000).
354. Jager, M., Nguyen, H., Crane, J. C., Kelly, J. W. & Gruebele, M. The folding mechanism of a beta-sheet: the WW domain. *J. Mol Biol.* **311**, 373-393 (2001).
355. van't Hof, W. & Resh, M. D. Dual fatty acylation of p59(Fyn) is required for association with the T cell receptor zeta chain through phosphotyrosine-Src homology domain-2 interactions. *J. Cell Biol.* **145**, 377-389 (1999).
356. Bedford, M. T. *et al.* Arginine methylation inhibits the binding of proline-rich ligands to Src homology 3, but not WW, domains. *J. Biol. Chem.* **275**, 16030-16036 (2000).
357. Burge, C. & Karlin, S. Prediction of complete gene structures in human genomic DNA. *J. Mol Biol.* **268**, 78-94 (1997).
358. Zhang, Y. *et al.* Evidence that dim1 associates with proteins involved in pre-mRNA splicing, and delineation of residues essential for dim1 interactions with hnRNP F and Npw38/PQBP-1. *Gene* **257**, 33-43 (2000).
359. Waragai, M. *et al.* PQBP-1/Npw38, a nuclear protein binding to the polyglutamine tract, interacts with U5-15kD/dim1p via the carboxyl-terminal domain. *Biochem. Biophys. Res. Commun.* **273**, 592-595 (2000).
360. Chan, B. *et al.* SAP couples Fyn to SLAM immune receptors. *Nat. Cell Biol.* **5**, 155-160 (2003).
361. Komuro, A., Saeki, M. & Kato, S. Association of two nuclear proteins, Npw38 and NpwBP, via the interaction between the WW domain and a novel proline-rich motif containing glycine and arginine. *J. Biol. Chem.* **274**, 36513-36519 (1999).
362. Waragai, M. *et al.* PQBP-1, a novel polyglutamine tract-binding protein, inhibits transcription activation by Brn-2 and affects cell survival. *Hum. Mol. Genet.* **8**, 977-987 (1999).
363. Okazawa, H. *et al.* Interaction between mutant ataxin-1 and PQBP-1 affects transcription and cell death. *Neuron* **34**, 701-713 (2002).
364. Blom, N., Gammeltoft, S. & Brunak, S. Sequence and structure-based prediction of eukaryotic protein phosphorylation sites. *J. Mol Biol.* **294**, 1351-1362 (1999).