
1	Table of Abbreviations	1
2	Introduction	3
2.1	Heat shock proteins	3
2.2	The innate and the adaptive immune response to hsp	5
2.3	Lymphocytes	6
2.4	T cell receptor signaling	11
2.5	T cell autoimmunity	13
2.6	The T cell clone UZ3/4	16
2.7	Inflammatory bowel disease	18
2.8	TCR transgenic mice	20
3	Aims of the study	22
4	Materials	24
4.1	Apparatus	24
4.2	Software	24
4.3	Chemicals, enzymes and kits	25
4.4	Organisms	25
4.5	Mammalian cell lines	26
4.6	Mice	26
4.7	Media and Buffers	26
4.8	Antibodies	26
4.9	Plasmids	28
4.10	Oligonucleotides	29

5	Methods	31
5.1	Molecular biological methods	31
5.1.1	<i>E.coli</i> heat shock transformation	31
5.1.2	Electroporation of <i>E. coli</i>	31
5.1.3	Polymerase chain reaction (PCR)	32
5.1.4	Southern Blot	32
5.1.5	RT-PCR	32
5.2	Cell culture and transfection	33
	Electroporation of cultured cells	33
5.2.2	Retroviral Transduction	34
5.3	Biochemical methods	35
5.3.1	Western Blot	35
5.3.2	Pulse-chase metabolic labeling and immunoprecipitation.	35
5.4	Generation of transgenic mice	36
5.4.1	Cloning strategies for TCR expression constructs	36
5.4.2	Preparation of transgenic vector samples	37
5.4.3	Pronucleus injection of transgenic vector samples into fertilized eggs	37
5.4.4	Screening of potential transgenic founders	38
5.5	Animal procedures	39
5.5.1	BCG infection	39
5.5.2	Peptide immunization	39
5.5.3	Heterozygous breeding	39
5.6	Immunological methods	39
5.6.1	Flow Cytometry and Cell Sorting	39
5.6.2	Generation of MHC class I tetramers and staining of cells with tetramers	40
5.6.3	⁵¹ Cr-release assay	41
5.6.4	Cell isolation from different tissues	42
5.6.5	CD69 upregulation assay	42
5.7	Histology.	43

6 Results	44
6.1 Expression of two TCR alpha chains in the CD8+ T cell clone UZ3/4	45
6.2 Surface expression of the TCR $\alpha 8$ chain but not the TCR $\alpha 7$ chain	47
6.3 Promiscuous peptide recognition of mycobacterial and murine peptides by the $\alpha 8/\beta 8$ TCR combination	51
6.4 Generation of TCR $\alpha 8$ and TCR $\beta 8$ transgenic mice	53
6.5 Peripheral TCR $\alpha 8$ and $\beta 8$ double transgenic mice are specific for the peptide SALQNAASIA from mycobacterial hsp60	58
6.6 Induction of CD8+ T cells specific for SALQNAASIA mycobacterial hsp60 peptide during mycobacterial infection	61
Generation of TCR $\alpha 7$ transgenic mice	64
Augmented levels of CD4+ TCR $\alpha^{-}\beta^{+}$ T cells in $\alpha 7.2$ mice	67
6.9 CD4+ TCR $\alpha^{-}\beta^{+}$ T cells show an activated phenotype	68
Accelerated course of IBD in $\alpha 7.2$ mice	69
6.11 Stabilization of endogenous TCR β chains by non-pairing TCR $\alpha 7.2$	72
7 Discussion	74
7.1 Promiscuous peptide recognition of a single TCR combination is responsible for the induction of autoimmune pathology	74
7.2 Hsp60 specific TCR transgenic mice	78
7.3 TCR V$\alpha 7$ chain transgenic mice	80
7.4 Concluding remarks and outlook:	84

8 Summary	87
9 Zusammenfassung	89
10 Reference List	91
Danksagung	111
Curriculum	112
Publications	114
Erklärung	115
Appendix	116
A. DNA sequences of the UZ3/4 TCR chains	116
TCR alpha 7 chain	116
TCR alpha 8 chain	117
TCR beta 8 chain	118
B. Plasmid charts	119
pP14 α AR	119
pP142 β 8AR	120
pHSE3'	121
pMSCV2.2-IRES-GFP	122
pCR2.1	123
pCMV/Zeo	124
C. Suppliers	125
D. Reprint	127