

7. Referenzen

- Ahmed, K., Gerber, D.A. und Cochet, C. (2002) Joining the cell survival squad: an emerging role for protein kinase CK2. *Trends Cell Biol*, **12**, 226-230.
- Alarcon-Vargas, D. und Ronai, Z. (2002) p53-Mdm2--the affair that never ends. *Carcinogenesis*, **23**, 541-547.
- Ang, L.H., Chattopadhyay, S., Wei, N., Oyama, T., Okada, K., Batschauer, A. und Deng, X.W. (1998) Molecular interaction between COP1 and HY5 defines a regulatory switch for light control of Arabidopsis development. *Mol Cell*, **1**, 213-222.
- Ang, L.H. und Deng, X.W. (1994) Regulatory hierarchy of photomorphogenic loci: allele-specific and light-dependent interaction between the HY5 and COP1 loci. *Plant Cell*, **6**, 613-628.
- Aravind, L. und Ponting, C.P. (1998) Homologues of 26S proteasome subunits are regulators of transcription and translation. *Protein Sci*, **7**, 1250-1254.
- Arbiser, J.L., Klauber, N., Rohan, R., van Leeuwen, R., Huang, M.T., Fisher, C., Flynn, E. und Byers, H.R. (1998) Curcumin is an in vivo inhibitor of angiogenesis. *Mol Med*, **4**, 376-383.
- Arora, R.B., Kapoor, V., Basu, N. und Jain, A.P. (1971) Anti-inflammatory studies on *Curcuma longa* (turmeric). *Indian J Med Res*, **59**, 1289-1295.
- Banin, S., Moyal, L., Shieh, S., Taya, Y., Anderson, C.W., Chessa, L., Smorodinsky, N.I., Prives, C., Reiss, Y., Shiloh, Y. und Ziv, Y. (1998) Enhanced phosphorylation of p53 by ATM in response to DNA damage. *Science*, **281**, 1674-1677.
- Barak, Y., Juven, T., Haffner, R. und Oren, M. (1993) mdm2 expression is induced by wild type p53 activity. *Embo J*, **12**, 461-468.
- Baron, C.L. und Malhotra, V. (2002) Role of diacylglycerol in PKD recruitment to the TGN and protein transport to the plasma membrane. *Science*, **295**, 325-328.
- Battistutta, R., Sarno, S., De Moliner, E., Papinutto, E., Zanotti, G. und Pinna, L.A. (2000) The replacement of ATP by the competitive inhibitor emodin induces conformational modifications in the catalytic site of protein kinase CK2. *J Biol Chem*, **275**, 29618-29622.
- Bech-Otschir, D., Kraft, R., Huang, X., Henklein, P., Kapelari, B., Pollmann, C. und Dubiel, W. (2001) COP9 signalosome-specific phosphorylation targets p53 to degradation by the ubiquitin system. *Embo J*, **20**, 1630-1639.
- Bech-Otschir, D., Seeger, M. und Dubiel, W. (2002) The COP9 signalosome: at the interface between signal transduction and ubiquitin-dependent proteolysis. *J Cell Sci*, **115**, 467-473.
- Blattner, C., Hay, T., Meek, D.W. und Lane, D.P. (2002) Hypophosphorylation of Mdm2 augments p53 stability. *Mol Cell Biol*, **22**, 6170-6182.
- Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*, **72**, 248-254.
- Braun, B.C., Glickman, M., Kraft, R., Dahlmann, B., Kloetzel, P.M., Finley, D. und Schmidt, M. (1999) The base of the proteasome regulatory particle exhibits chaperone-like activity. *Nat Cell Biol*, **1**, 221-226.
- Burnett, G. und Kennedy, E.P. (1954) The enzymatic phosphorylation of proteins. *J Biol Chem*, **211**, 969-980.
- Buschmann, T., Fuchs, S.Y., Lee, C.G., Pan, Z.Q. und Ronai, Z. (2000) SUMO-1 modification of Mdm2 prevents its self-ubiquitination and increases Mdm2 ability to ubiquitinate p53. *Cell*, **101**, 753-762.

- Cahilly-Snyder, L., Yang-Feng, T., Francke, U. und George, D.L. (1987) Molecular analysis and chromosomal mapping of amplified genes isolated from a transformed mouse 3T3 cell line. *Somat Cell Mol Genet*, **13**, 235-244.
- Canton, D.A., Zhang, C. und Litchfield, D.W. (2001) Assembly of protein kinase CK2: investigation of complex formation between catalytic and regulatory subunits using a zinc-finger-deficient mutant of CK2beta. *Biochem J*, **358**, 87-94.
- Chamovitz, D.A., Wei, N., Osterlund, M.T., von Arnim, A.G., Staub, J.M., Matsui, M. und Deng, X.W. (1996) The COP9 complex, a novel multisubunit nuclear regulator involved in light control of a plant developmental switch. *Cell*, **86**, 115-121.
- Chehab, N.H., Malikzay, A., Appel, M. und Halazonetis, T.D. (2000) Chk2/hCds1 functions as a DNA damage checkpoint in G(1) by stabilizing p53. *Genes Dev*, **14**, 278-288.
- Chen, Y.R. und Tan, T.H. (1998) Inhibition of the c-Jun N-terminal kinase (JNK) signaling pathway by curcumin. *Oncogene*, **17**, 173-178.
- Cope, G.A., Suh, G.S., Aravind, L., Schwarz, S.E., Zipursky, S.L., Koonin, E.V. und Deshaies, R.J. (2002) Role of predicted metalloprotease motif of Jab1/Csn5 in cleavage of Nedd8 from Cul1. *Science*, **298**, 608-611.
- Daujat, S., Neel, H. und Piette, J. (2001) MDM2: life without p53. *Trends Genet*, **17**, 459-464.
- Deng, X.W., Dubiel, W., Wei, N., Hofmann, K., Mundt, K., Colicelli, J., Kato, J., Naumann, M., Segal, D., Seeger, M., Carr, A., Glickman, M. und Chamovitz, D.A. (2000) Unified nomenclature for the COP9 signalosome and its subunits: an essential regulator of development. *Trends Genet*, **16**, 202-203.
- Derijard, B., Hibi, M., Wu, I.H., Barrett, T., Su, B., Deng, T., Karin, M. und Davis, R.J. (1994) JNK1: a protein kinase stimulated by UV light and Ha-Ras that binds and phosphorylates the c-Jun activation domain. *Cell*, **76**, 1025-1037.
- Desagher, S., Osen-Sand, A., Montessuit, S., Magnenat, E., Vilbois, F., Hochmann, A., Journot, L., Antonsson, B. und Martinou, J.C. (2001) Phosphorylation of bid by casein kinases I and II regulates its cleavage by caspase 8. *Mol Cell*, **8**, 601-611.
- Deveraux, Q., Ustrell, V., Pickart, C. und Rechsteiner, M. (1994) A 26 S protease subunit that binds ubiquitin conjugates. *J Biol Chem*, **269**, 7059-7061.
- Dornan, D., Wertz, I., Shimizu, H., Arnott, D., Frantz, G.D., Dowd, P., K, O.R., Koepfen, H. und Dixit, V.M. (2004) The ubiquitin ligase COP1 is a critical negative regulator of p53. *Nature*.
- Doronkin, S., Djagaeva, I. und Beckendorf, S.K. (2002) CSN5/Jab1 mutations affect axis formation in the Drosophila oocyte by activating a meiotic checkpoint. *Development*, **129**, 5053-5064.
- Dubiel, W. und Gordon, C. (1999) Ubiquitin pathway: another link in the polyubiquitin chain? *Curr Biol*, **9**, R554-557.
- Elenbaas, B., Dobbstein, M., Roth, J., Shenk, T. und Levine, A.J. (1996) The MDM2 oncoprotein binds specifically to RNA through its RING finger domain. *Mol Med*, **2**, 439-451.
- Endo, K., Oki, E., Biedermann, V., Kojima, H., Yoshida, K., Johannes, F.J., Kufe, D. und Datta, R. (2000) Proteolytic cleavage and activation of protein kinase C [micro] by caspase-3 in the apoptotic response of cells to 1-beta -D-arabinofuranosylcytosine and other genotoxic agents. *J Biol Chem*, **275**, 18476-18481.

- Fakharzadeh, S.S., Trusko, S.P. und George, D.L. (1991) Tumorigenic potential associated with enhanced expression of a gene that is amplified in a mouse tumor cell line. *Embo J*, **10**, 1565-1569.
- Fang, S., Jensen, J.P., Ludwig, R.L., Vousden, K.H. und Weissman, A.M. (2000) Mdm2 is a RING finger-dependent ubiquitin protein ligase for itself and p53. *J Biol Chem*, **275**, 8945-8951.
- Faust, M. und Montenarh, M. (2000) Subcellular localization of protein kinase CK2. A key to its function? *Cell Tissue Res*, **301**, 329-340.
- Feldman, R.M., Correll, C.C., Kaplan, K.B. und Deshaies, R.J. (1997) A complex of Cdc4p, Skp1p, and Cdc53p/cullin catalyzes ubiquitination of the phosphorylated CDK inhibitor Sic1p. *Cell*, **91**, 221-230.
- Ferrell, K., Wilkinson, C.R., Dubiel, W. und Gordon, C. (2000) Regulatory subunit interactions of the 26S proteasome, a complex problem. *Trends Biochem Sci*, **25**, 83-88.
- Ficarro, S.B., McClelland, M.L., Stukenberg, P.T., Burke, D.J., Ross, M.M., Shabanowitz, J., Hunt, D.F. und White, F.M. (2002) Phosphoproteome analysis by mass spectrometry and its application to *Saccharomyces cerevisiae*. *Nat Biotechnol*, **20**, 301-305.
- Freedman, D.A. und Levine, A.J. (1998) Nuclear export is required for degradation of endogenous p53 by MDM2 and human papillomavirus E6. *Mol Cell Biol*, **18**, 7288-7293.
- Freilich, S., Oron, E., Kapp, Y., Nevo-Caspi, Y., Orgad, S., Segal, D. und Chamovitz, D.A. (1999) The COP9 signalosome is essential for development of *Drosophila melanogaster*. *Curr Biol*, **9**, 1187-1190.
- Fuchs, S.Y., Chen, A., Xiong, Y., Pan, Z.Q. und Ronai, Z. (1999) HOS, a human homolog of Slimb, forms an SCF complex with Skp1 and Cullin1 and targets the phosphorylation-dependent degradation of I κ B and beta-catenin. *Oncogene*, **18**, 2039-2046.
- Gavin, A.C., Bosche, M., Krause, R., Grandi, P., Marzioch, M., Bauer, A., Schultz, J., Rick, J.M., Michon, A.M., Cruciat, C.M., Remor, M., Hofert, C., Schelder, M., Brajenovic, M., Ruffner, H., Merino, A., Klein, K., Hudak, M., Dickson, D., Rudi, T., Gnau, V., Bauch, A., Bastuck, S., Huhse, B., Leutwein, C., Heurtier, M.A., Copley, R.R., Edelmann, A., Querfurth, E., Rybin, V., Drewes, G., Raida, M., Bouwmeester, T., Bork, P., Seraphin, B., Kuster, B., Neubauer, G. und Superti-Furga, G. (2002) Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature*, **415**, 141-147.
- Geyer, R.K., Yu, Z.K. und Maki, C.G. (2000) The MDM2 RING-finger domain is required to promote p53 nuclear export. *Nat Cell Biol*, **2**, 569-573.
- Giaccia, A.J. und Kastan, M.B. (1998) The complexity of p53 modulation: emerging patterns from divergent signals. *Genes Dev*, **12**, 2973-2983.
- Glickman, M.H., Rubin, D.M., Coux, O., Wefes, I., Pfeifer, G., Cjeka, Z., Baumeister, W., Fried, V.A. und Finley, D. (1998) A subcomplex of the proteasome regulatory particle required for ubiquitin-conjugate degradation and related to the COP9-signalosome and eIF3. *Cell*, **94**, 615-623.
- Glover, C.V., 3rd. (1998) On the physiological role of casein kinase II in *Saccharomyces cerevisiae*. *Prog Nucleic Acid Res Mol Biol*, **59**, 95-133.
- Goldberg, Z., Vogt Sionov, R., Berger, M., Zwang, Y., Perets, R., Van Etten, R.A., Oren, M., Taya, Y. und Haupt, Y. (2002) Tyrosine phosphorylation of Mdm2 by c-Abl: implications for p53 regulation. *Embo J*, **21**, 3715-3727.

- Götz, C., Kartarius, S., Scholtes, P., Nastainczyk, W. und Montenarh, M. (1999) Identification of a CK2 phosphorylation site in mdm2. *Eur J Biochem*, **266**, 493-501.
- Graham, K.C. und Litchfield, D.W. (2000) The regulatory beta subunit of protein kinase CK2 mediates formation of tetrameric CK2 complexes. *J Biol Chem*, **275**, 5003-5010.
- Grossman, S.R., Perez, M., Kung, A.L., Joseph, M., Mansur, C., Xiao, Z.X., Kumar, S., Howley, P.M. und Livingston, D.M. (1998) p300/MDM2 complexes participate in MDM2-mediated p53 degradation. *Mol Cell*, **2**, 405-415.
- Guerra, B. und Issinger, O.G. (1999) Protein kinase CK2 and its role in cellular proliferation, development and pathology. *Electrophoresis*, **20**, 391-408.
- Hardtke, C.S. und Deng, X.W. (2000) The cell biology of the COP/DET/FUS proteins. Regulating proteolysis in photomorphogenesis and beyond? *Plant Physiol*, **124**, 1548-1557.
- Hardtke, C.S., Gohda, K., Osterlund, M.T., Oyama, T., Okada, K. und Deng, X.W. (2000) HY5 stability and activity in arabidopsis is regulated by phosphorylation in its COP1 binding domain. *Embo J*, **19**, 4997-5006.
- Hasmeda, M. und Polya, G.M. (1996) Inhibition of cyclic AMP-dependent protein kinase by curcumin. *Phytochemistry*, **42**, 599-605.
- Haupt, S., Loria-Hayon, I. und Haupt, Y. (2003) P53 licensed to kill? Operating the assassin. *J Cell Biochem*, **88**, 76-82.
- Haupt, Y., Maya, R., Kazaz, A. und Oren, M. (1997) Mdm2 promotes the rapid degradation of p53. *Nature*, **387**, 296-299.
- Hausser, A., Storz, P., Link, G., Stoll, H., Liu, Y.C., Altman, A., Pfizenmaier, K. und Johannes, F.J. (1999) Protein kinase C mu is negatively regulated by 14-3-3 signal transduction proteins. *J Biol Chem*, **274**, 9258-9264.
- Haworth, R.S. und Avkiran, M. (2001) Inhibition of protein kinase D by resveratrol. *Biochem Pharmacol*, **62**, 1647-1651.
- Hayashi, A., Seki, N., Hattori, A., Kozuma, S. und Saito, T. (1999) PKCnu, a new member of the protein kinase C family, composes a fourth subfamily with PKCmu. *Biochim Biophys Acta*, **1450**, 99-106.
- Henke, W., Ferrell, K., Bech-Otschir, D., Seeger, M., Schade, R., Jungblut, P., Naumann, M. und Dubiel, W. (1999) Comparison of human COP9 signalsome and 26S proteasome lid'. *Mol Biol Rep*, **26**, 29-34.
- Hershko, A. und Ciechanover, A. (1998) The ubiquitin system. *Annu Rev Biochem*, **67**, 425-479.
- Hirao, A., Kong, Y.Y., Matsuoka, S., Wakeham, A., Ruland, J., Yoshida, H., Liu, D., Elledge, S.J. und Mak, T.W. (2000) DNA damage-induced activation of p53 by the checkpoint kinase Chk2. *Science*, **287**, 1824-1827.
- Hjerrild, M., Milne, D., Dumaz, N., Hay, T., Issinger, O.G. und Meek, D. (2001) Phosphorylation of murine double minute clone 2 (MDM2) protein at serine-267 by protein kinase CK2 in vitro and in cultured cells. *Biochem J*, **355**, 347-356.
- Ho, J.H., Kallstrom, G. und Johnson, A.W. (2000) Nmd3p is a Crm1p-dependent adapter protein for nuclear export of the large ribosomal subunit. *J Cell Biol*, **151**, 1057-1066.
- Hochstrasser, M. (1996) Ubiquitin-dependent protein degradation. *Annu Rev Genet*, **30**, 405-439.
- Hofmann, K. und Bucher, P. (1998) The PCI domain: a common theme in three multiprotein complexes. *Trends Biochem Sci*, **23**, 204-205.
- Honda, R., Tanaka, H. und Yasuda, H. (1997) Oncoprotein MDM2 is a ubiquitin ligase E3 for tumor suppressor p53. *FEBS Lett*, **420**, 25-27.

- Honda, R. und Yasuda, H. (1999) Association of p19(ARF) with Mdm2 inhibits ubiquitin ligase activity of Mdm2 for tumor suppressor p53. *Embo J*, **18**, 22-27.
- Honda, R. und Yasuda, H. (2000) Activity of MDM2, a ubiquitin ligase, toward p53 or itself is dependent on the RING finger domain of the ligase. *Oncogene*, **19**, 1473-1476.
- Huang, M.T., Lou, Y.R., Ma, W., Newmark, H.L., Reuhl, K.R. und Conney, A.H. (1994) Inhibitory effects of dietary curcumin on forestomach, duodenal, and colon carcinogenesis in mice. *Cancer Res*, **54**, 5841-5847.
- Hurd, C. und Rozengurt, E. (2001) Protein kinase D is sufficient to suppress EGF-induced c-Jun Ser 63 phosphorylation. *Biochem Biophys Res Commun*, **282**, 404-408.
- Hurd, C., Waldron, R.T. und Rozengurt, E. (2002) Protein kinase D complexes with C-Jun N-terminal kinase via activation loop phosphorylation and phosphorylates the C-Jun N-terminus. *Oncogene*, **21**, 2154-2160.
- Iglesias, T., Cabrera-Poch, N., Mitchell, M.P., Naven, T.J., Rozengurt, E. und Schiavo, G. (2000) Identification and cloning of Kidins220, a novel neuronal substrate of protein kinase D. *J Biol Chem*, **275**, 40048-40056.
- Iglesias, T., Waldron, R.T. und Rozengurt, E. (1998) Identification of in vivo phosphorylation sites required for protein kinase D activation. *J Biol Chem*, **273**, 27662-27667.
- Issinger, O.G. (1977) Purification and properties of a ribosomal casein kinase from rabbit reticulocytes. *Biochem J*, **165**, 511-518.
- Iwakuma, T. und Lozano, G. (2003) MDM2, an introduction. *Mol Cancer Res*, **1**, 993-1000.
- Jackson, P.K., Eldridge, A.G., Freed, E., Furstenthal, L., Hsu, J.Y., Kaiser, B.K. und Reimann, J.D. (2000) The lore of the RINGs: substrate recognition and catalysis by ubiquitin ligases. *Trends Cell Biol*, **10**, 429-439.
- Jamora, C., Takizawa, P.A., Zaarour, R.F., Denesvre, C., Faulkner, D.J. und Malhotra, V. (1997) Regulation of Golgi structure through heterotrimeric G proteins. *Cell*, **91**, 617-626.
- Jamora, C., Yamanouye, N., Van Lint, J., Laudenslager, J., Vandenheede, J.R., Faulkner, D.J. und Malhotra, V. (1999) Gbetagamma-mediated regulation of Golgi organization is through the direct activation of protein kinase D. *Cell*, **98**, 59-68.
- Jesenberger, V. und Jentsch, S. (2002) Deadly encounter: ubiquitin meets apoptosis. *Nat Rev Mol Cell Biol*, **3**, 112-121.
- Johannes, F.J., Prestle, J., Eis, S., Oberhagemann, P. und Pfizenmaier, K. (1994) PKCu is a novel, atypical member of the protein kinase C family. *J Biol Chem*, **269**, 6140-6148.
- Jones, S.N., Roe, A.E., Donehower, L.A. und Bradley, A. (1995) Rescue of embryonic lethality in Mdm2-deficient mice by absence of p53. *Nature*, **378**, 206-208.
- Kapelari, B., Bech-Otschir, D., Hegerl, R., Schade, R., Dumdey, R. und Dubiel, W. (2000) Electron microscopy and subunit-subunit interaction studies reveal a first architecture of COP9 signalosome. *J Mol Biol*, **300**, 1169-1178.
- Karniol, B., Malec, P. und Chamovitz, D.A. (1999) Arabidopsis FUSCA5 encodes a novel phosphoprotein that is a component of the COP9 complex. *Plant Cell*, **11**, 839-848.
- Kasahara, H. und Izumo, S. (1999) Identification of the in vivo casein kinase II phosphorylation site within the homeodomain of the cardiac tissue-specifying homeobox gene product Csx/Nkx2.5. *Mol Cell Biol*, **19**, 526-536.

- Kipreos, E.T., Lander, L.E., Wing, J.P., He, W.W. und Hedgecock, E.M. (1996) *cul-1* is required for cell cycle exit in *C. elegans* and identifies a novel gene family. *Cell*, **85**, 829-839.
- Koegl, M., Hoppe, T., Schlenker, S., Ulrich, H.D., Mayer, T.U. und Jentsch, S. (1999) A novel ubiquitination factor, E4, is involved in multiubiquitin chain assembly. *Cell*, **96**, 635-644.
- Krek, W., Maridor, G. und Nigg, E.A. (1992) Casein kinase II is a predominantly nuclear enzyme. *J Cell Biol*, **116**, 43-55.
- Krippner-Heidenreich, A., Talanian, R.V., Sekul, R., Kraft, R., Thole, H., Oettleben, H. und Luscher, B. (2001) Targeting of the transcription factor Max during apoptosis: phosphorylation-regulated cleavage by caspase-5 at an unusual glutamic acid residue in position P1. *Biochem J*, **358**, 705-715.
- Kubbutat, M.H., Jones, S.N. und Vousden, K.H. (1997) Regulation of p53 stability by Mdm2. *Nature*, **387**, 299-303.
- Kubbutat, M.H., Ludwig, R.L., Ashcroft, M. und Vousden, K.H. (1998) Regulation of Mdm2-directed degradation by the C terminus of p53. *Mol Cell Biol*, **18**, 5690-5698.
- Kurokawa, K., Tanaka, T. und Kato, J. (1999) p19ARF prevents G1 cyclin-dependent kinase activation by interacting with MDM2 and activating p53 in mouse fibroblasts. *Oncogene*, **18**, 2718-2727.
- Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, **227**, 680-685.
- Lai, Z., Ferry, K.V., Diamond, M.A., Wee, K.E., Kim, Y.B., Ma, J., Yang, T., Benfield, P.A., Copeland, R.A. und Auger, K.R. (2001) Human mdm2 mediates multiple mono-ubiquitination of p53 by a mechanism requiring enzyme isomerization. *J Biol Chem*, **276**, 31357-31367.
- Lam, Y.A., Lawson, T.G., Velayutham, M., Zweier, J.L. und Pickart, C.M. (2002) A proteasomal ATPase subunit recognizes the polyubiquitin degradation signal. *Nature*, **416**, 763-767.
- Leng, R.P., Lin, Y., Ma, W., Wu, H., Lemmers, B., Chung, S., Parant, J.M., Lozano, G., Hakem, R. und Benchimol, S. (2003) Pirh2, a p53-induced ubiquitin-protein ligase, promotes p53 degradation. *Cell*, **112**, 779-791.
- Linares, L.K., Hengstermann, A., Ciechanover, A., Muller, S. und Scheffner, M. (2003) HdmX stimulates Hdm2-mediated ubiquitination and degradation of p53. *Proc Natl Acad Sci U S A*, **100**, 12009-12014.
- Litchfield, D.W. (2003) Protein kinase CK2: structure, regulation and role in cellular decisions of life and death. *Biochem J*, **369**, 1-15.
- Lohrum, M.A., Ashcroft, M., Kubbutat, M.H. und Vousden, K.H. (2000) Identification of a cryptic nucleolar-localization signal in MDM2. *Nat Cell Biol*, **2**, 179-181.
- Lohrum, M.A., Ludwig, R.L., Kubbutat, M.H., Hanlon, M. und Vousden, K.H. (2003) Regulation of HDM2 activity by the ribosomal protein L11. *Cancer Cell*, **3**, 577-587.
- Lyapina, S., Cope, G., Shevchenko, A., Serino, G., Tsuge, T., Zhou, C., Wolf, D.A., Wei, N. und Deshaies, R.J. (2001) Promotion of NEDD-CUL1 conjugate cleavage by COP9 signalosome. *Science*, **292**, 1382-1385.
- Maeda, Y., Beznoussenko, G.V., Van Lint, J., Mironov, A.A. und Malhotra, V. (2001) Recruitment of protein kinase D to the trans-Golgi network via the first cysteine-rich domain. *Embo J*, **20**, 5982-5990.
- Maki, C.G. (1999) Oligomerization is required for p53 to be efficiently ubiquitinated by MDM2. *J Biol Chem*, **274**, 16531-16535.

- Marin, O., Meggio, F. und Pinna, L.A. (1999) Structural features underlying the unusual mode of calmodulin phosphorylation by protein kinase CK2: A study with synthetic calmodulin fragments. *Biochem Biophys Res Commun*, **256**, 442-446.
- Martel, V., Filhol, O., Nueda, A., Gerber, D., Benitez, M.J. und Cochet, C. (2001) Visualization and molecular analysis of nuclear import of protein kinase CK2 subunits in living cells. *Mol Cell Biochem*, **227**, 81-90.
- Matthews, S., Iglesias, T., Cantrell, D. und Rozengurt, E. (1999) Dynamic redistribution of protein kinase D (PKD) as revealed by a GFP-PKD fusion protein: dissociation from PKD activation. *FEBS Lett*, **457**, 515-521.
- Mauxion, F., Le Borgne, R., Munier-Lehmann, H. und Hoflack, B. (1996) A casein kinase II phosphorylation site in the cytoplasmic domain of the cation-dependent mannose 6-phosphate receptor determines the high affinity interaction of the AP-1 Golgi assembly proteins with membranes. *J Biol Chem*, **271**, 2171-2178.
- Mayo, L.D. und Donner, D.B. (2001) A phosphatidylinositol 3-kinase/Akt pathway promotes translocation of Mdm2 from the cytoplasm to the nucleus. *Proc Natl Acad Sci U S A*, **98**, 11598-11603.
- Mayo, L.D., Turchi, J.J. und Berberich, S.J. (1997) Mdm-2 phosphorylation by DNA-dependent protein kinase prevents interaction with p53. *Cancer Res*, **57**, 5013-5016.
- Maytal-Kivity, V., Piran, R., Pick, E., Hofmann, K. und Glickman, M.H. (2002) COP9 signalosome components play a role in the mating pheromone response of *S. cerevisiae*. *EMBO Rep*, **3**, 1215-1221.
- Meek, D.W. und Knippschild, U. (2003) Posttranslational modification of MDM2. *Mol Cancer Res*, **1**, 1017-1026.
- Meggio, F., Boldyreff, B., Marin, O., Marchiori, F., Perich, J.W., Issinger, O.G. und Pinna, L.A. (1992) The effect of polylysine on casein-kinase-2 activity is influenced by both the structure of the protein/peptide substrates and the subunit composition of the enzyme. *Eur J Biochem*, **205**, 939-945.
- Meggio, F., Marin, O. und Pinna, L.A. (1994) Substrate specificity of protein kinase CK2. *Cell Mol Biol Res*, **40**, 401-409.
- Meggio, F. und Pinna, L.A. (2003) One-thousand-and-one substrates of protein kinase CK2? *Faseb J*, **17**, 349-368.
- Mehta, K., Pantazis, P., McQueen, T. und Aggarwal, B.B. (1997) Antiproliferative effect of curcumin (diferuloylmethane) against human breast tumor cell lines. *Anticancer Drugs*, **8**, 470-481.
- Michael, D. und Oren, M. (2002) The p53 and Mdm2 families in cancer. *Curr Opin Genet Dev*, **12**, 53-59.
- Midgley, C.A., Desterro, J.M., Saville, M.K., Howard, S., Sparks, A., Hay, R.T. und Lane, D.P. (2000) An N-terminal p14ARF peptide blocks Mdm2-dependent ubiquitination in vitro and can activate p53 in vivo. *Oncogene*, **19**, 2312-2323.
- Midgley, C.A. und Lane, D.P. (1997) p53 protein stability in tumour cells is not determined by mutation but is dependent on Mdm2 binding. *Oncogene*, **15**, 1179-1189.
- Momand, J., Zambetti, G.P., Olson, D.C., George, D. und Levine, A.J. (1992) The mdm-2 oncogene product forms a complex with the p53 protein and inhibits p53-mediated transactivation. *Cell*, **69**, 1237-1245.
- Montes de Oca Luna, R., Wagner, D.S. und Lozano, G. (1995) Rescue of early embryonic lethality in mdm2-deficient mice by deletion of p53. *Nature*, **378**, 203-206.

- Mundt, K.E., Liu, C. und Carr, A.M. (2002) Deletion mutants in COP9/signalosome subunits in fission yeast *Schizosaccharomyces pombe* display distinct phenotypes. *Mol Biol Cell*, **13**, 493-502.
- Mundt, K.E., Porte, J., Murray, J.M., Brikos, C., Christensen, P.U., Caspari, T., Hagan, I.M., Millar, J.B., Simanis, V., Hofmann, K. und Carr, A.M. (1999) The COP9/signalosome complex is conserved in fission yeast and has a role in S phase. *Curr Biol*, **9**, 1427-1430.
- Nateri, A.S., Riera-Sans, L., Da Costa, C. und Behrens, A. (2004) The ubiquitin ligase SCFFbw7 antagonizes apoptotic JNK signaling. *Science*, **303**, 1374-1378.
- Naumann, M., Bech-Otschir, D., Huang, X., Ferrell, K. und Dubiel, W. (1999) COP9 signalosome-directed c-Jun activation/stabilization is independent of JNK. *J Biol Chem*, **274**, 35297-35300.
- Niefind, K., Guerra, B., Ermakowa, I. und Issinger, O.G. (2001) Crystal structure of human protein kinase CK2: insights into basic properties of the CK2 holoenzyme. *Embo J*, **20**, 5320-5331.
- Nishikawa, K., Toker, A., Johannes, F.J., Songyang, Z. und Cantley, L.C. (1997) Determination of the specific substrate sequence motifs of protein kinase C isozymes. *J Biol Chem*, **272**, 952-960.
- Nishikawa, K., Toker, A., Wong, K., Marignani, P.A., Johannes, F.J. und Cantley, L.C. (1998) Association of protein kinase Cmu with type II phosphatidylinositol 4-kinase and type I phosphatidylinositol-4-phosphate 5-kinase. *J Biol Chem*, **273**, 23126-23133.
- Okumura, N., Saji, S., Eguchi, H., Nakashima, S. und Hayashi, S. (2002) Distinct promoter usage of mdm2 gene in human breast cancer. *Oncol Rep*, **9**, 557-563.
- Oliner, J.D., Kinzler, K.W., Meltzer, P.S., George, D.L. und Vogelstein, B. (1992) Amplification of a gene encoding a p53-associated protein in human sarcomas. *Nature*, **358**, 80-83.
- Oliner, J.D., Pietenpol, J.A., Thiagalingam, S., Gyuris, J., Kinzler, K.W. und Vogelstein, B. (1993) Oncoprotein MDM2 conceals the activation domain of tumour suppressor p53. *Nature*, **362**, 857-860.
- Olson, D.C., Marechal, V., Momand, J., Chen, J., Romocki, C. und Levine, A.J. (1993) Identification and characterization of multiple mdm-2 proteins and mdm-2-p53 protein complexes. *Oncogene*, **8**, 2353-2360.
- Oren, M., Damalas, A., Gottlieb, T., Michael, D., Taplick, J., Leal, J.F., Maya, R., Moas, M., Seger, R., Taya, Y. und Ben-Ze'Ev, A. (2002) Regulation of p53: intricate loops and delicate balances. *Ann N Y Acad Sci*, **973**, 374-383.
- Oron, E., Mannervik, M., Rencus, S., Harari-Steinberg, O., Neuman-Silberberg, S., Segal, D. und Chamovitz, D.A. (2002) COP9 signalosome subunits 4 and 5 regulate multiple pleiotropic pathways in *Drosophila melanogaster*. *Development*, **129**, 4399-4409.
- Osterlund, M.T., Ang, L.H. und Deng, X.W. (1999) The role of COP1 in repression of *Arabidopsis* photomorphogenic development. *Trends Cell Biol*, **9**, 113-118.
- Osterlund, M.T., Hardtke, C.S., Wei, N. und Deng, X.W. (2000) Targeted destabilization of HY5 during light-regulated development of *Arabidopsis*. *Nature*, **405**, 462-466.
- Padmanabha, R., Chen-Wu, J.L., Hanna, D.E. und Glover, C.V. (1990) Isolation, sequencing, and disruption of the yeast CKA2 gene: casein kinase II is essential for viability in *Saccharomyces cerevisiae*. *Mol Cell Biol*, **10**, 4089-4099.

- Pearson, M., Carbone, R., Sebastiani, C., Cioce, M., Fagioli, M., Saito, S., Higashimoto, Y., Appella, E., Minucci, S., Pandolfi, P.P. und Pelicci, P.G. (2000) PML regulates p53 acetylation and premature senescence induced by oncogenic Ras. *Nature*, **406**, 207-210.
- Pollmann, C., Huang, X., Mall, J., Bech-Otschir, D., Naumann, M. und Dubiel, W. (2001) The constitutive photomorphogenesis 9 signalosome directs vascular endothelial growth factor production in tumor cells. *Cancer Res*, **61**, 8416-8421.
- Pomerantz, J., Schreiber-Agus, N., Liegeois, N.J., Silverman, A., Alland, L., Chin, L., Potes, J., Chen, K., Orlow, I., Lee, H.W., Cordon-Cardo, C. und DePinho, R.A. (1998) The Ink4a tumor suppressor gene product, p19Arf, interacts with MDM2 and neutralizes MDM2's inhibition of p53. *Cell*, **92**, 713-723.
- Ponting, C.P., Aravind, L., Schultz, J., Bork, P. und Koonin, E.V. (1999) Eukaryotic signalling domain homologues in archaea and bacteria. Ancient ancestry and horizontal gene transfer. *J Mol Biol*, **289**, 729-745.
- Poyurovsky, M.V., Jacq, X., Ma, C., Karni-Schmidt, O., Parker, P.J., Chalfie, M., Manley, J.L. und Prives, C. (2003) Nucleotide binding by the Mdm2 RING domain facilitates Arf-independent Mdm2 nucleolar localization. *Mol Cell*, **12**, 875-887.
- Rechsteiner, M., Hoffman, L. und Dubiel, W. (1993) The multicatalytic and 26 S proteases. *J Biol Chem*, **268**, 6065-6068.
- Reddy, A.C. und Lokesh, B.R. (1994) Studies on the inhibitory effects of curcumin and eugenol on the formation of reactive oxygen species and the oxidation of ferrous iron. *Mol Cell Biochem*, **137**, 1-8.
- Roach, P.J. (1991) Multisite and hierarchical protein phosphorylation. *J Biol Chem*, **266**, 14139-14142.
- Roth, J., Dobbstein, M., Freedman, D.A., Shenk, T. und Levine, A.J. (1998) Nucleocytoplasmic shuttling of the hdm2 oncoprotein regulates the levels of the p53 protein via a pathway used by the human immunodeficiency virus rev protein. *Embo J*, **17**, 554-564.
- Sarrouilhe, D., Filhol, O., Leroy, D., Bonello, G., Baudry, M., Chambaz, E.M. und Cochet, C. (1998) The tight association of protein kinase CK2 with plasma membranes is mediated by a specific domain of its regulatory beta-subunit. *Biochim Biophys Acta*, **1403**, 199-210.
- Seeger, M., Kraft, R., Ferrell, K., Bech-Otschir, D., Dumdey, R., Schade, R., Gordon, C., Naumann, M. und Dubiel, W. (1998) A novel protein complex involved in signal transduction possessing similarities to 26S proteasome subunits. *Faseb J*, **12**, 469-478.
- Sharp, D.A., Kratowicz, S.A., Sank, M.J. und George, D.L. (1999) Stabilization of the MDM2 oncoprotein by interaction with the structurally related MDMX protein. *J Biol Chem*, **274**, 38189-38196.
- Shi, X., Potvin, B., Huang, T., Hilgard, P., Spray, D.C., Suadicani, S.O., Wolkoff, A.W., Stanley, P. und Stockert, R.J. (2001) A novel casein kinase 2 alpha-subunit regulates membrane protein traffic in the human hepatoma cell line HuH-7. *J Biol Chem*, **276**, 2075-2082.
- Shieh, S.Y., Ikeda, M., Taya, Y. und Prives, C. (1997) DNA damage-induced phosphorylation of p53 alleviates inhibition by MDM2. *Cell*, **91**, 325-334.
- Shirangi, T.R., Zaika, A. und Moll, U.M. (2002) Nuclear degradation of p53 occurs during down-regulation of the p53 response after DNA damage. *Faseb J*, **16**, 420-422.

- Smith, P., Leung-Chiu, W.M., Montgomery, R., Orsborn, A., Kuznicki, K., Gressman-Coberly, E., Mutapcic, L. und Bennett, K. (2002) The GLH proteins, *Caenorhabditis elegans* P granule components, associate with CSN-5 and KGB-1, proteins necessary for fertility, and with ZYX-1, a predicted cytoskeletal protein. *Dev Biol*, **251**, 333-347.
- Sreejayan und Rao, M.N. (1994) Curcuminoids as potent inhibitors of lipid peroxidation. *J Pharm Pharmacol*, **46**, 1013-1016.
- Stad, R., Little, N.A., Xirodimas, D.P., Frenk, R., van der Eb, A.J., Lane, D.P., Saville, M.K. und Jochemsen, A.G. (2001) Mdmx stabilizes p53 and Mdm2 via two distinct mechanisms. *EMBO Rep*, **2**, 1029-1034.
- Stommel, J.M. und Wahl, G.M. (2004) Accelerated MDM2 auto-degradation induced by DNA-damage kinases is required for p53 activation. *Embo J*, **23**, 1547-1556.
- Stott, F.J., Bates, S., James, M.C., McConnell, B.B., Starborg, M., Brookes, S., Palmero, I., Ryan, K., Hara, E., Vousden, K.H. und Peters, G. (1998) The alternative product from the human CDKN2A locus, p14(ARF), participates in a regulatory feedback loop with p53 and MDM2. *Embo J*, **17**, 5001-5014.
- Sturany, S., Van Lint, J., Muller, F., Wilda, M., Hameister, H., Hocker, M., Brey, A., Gern, U., Vandenheede, J., Gress, T., Adler, G. und Seufferlein, T. (2001) Molecular cloning and characterization of the human protein kinase D2. A novel member of the protein kinase D family of serine threonine kinases. *J Biol Chem*, **276**, 3310-3318.
- Suh, G.S., Poeck, B., Chouard, T., Oron, E., Segal, D., Chamovitz, D.A. und Zipursky, S.L. (2002) *Drosophila* JAB1/CSN5 acts in photoreceptor cells to induce glial cells. *Neuron*, **33**, 35-46.
- Sun, Y., Wilson, M.P. und Majerus, P.W. (2002) Inositol 1,3,4-trisphosphate 5/6-kinase associates with the COP9 signalosome by binding to CSN1. *J Biol Chem*, **277**, 45759-45764.
- Suzuki, G., Yanagawa, Y., Kwok, S.F., Matsui, M. und Deng, X.W. (2002) Arabidopsis COP10 is a ubiquitin-conjugating enzyme variant that acts together with COP1 and the COP9 signalosome in repressing photomorphogenesis. *Genes Dev*, **16**, 554-559.
- Tanimura, S., Ohtsuka, S., Mitsui, K., Shirouzu, K., Yoshimura, A. und Ohtsubo, M. (1999) MDM2 interacts with MDMX through their RING finger domains. *FEBS Lett*, **447**, 5-9.
- Tao, W. und Levine, A.J. (1999) Nucleocytoplasmic shuttling of oncoprotein Hdm2 is required for Hdm2-mediated degradation of p53. *Proc Natl Acad Sci U S A*, **96**, 3077-3080.
- Taylor, J.A., Bren, G.D., Pennington, K.N., Trushin, S.A., Asin, S. und Paya, C.V. (1999) Serine 32 and serine 36 of I κ B α are directly phosphorylated by protein kinase CKII in vitro. *J Mol Biol*, **290**, 839-850.
- te Poele, R.H., Okorokov, A.L. und Joel, S.P. (1999) RNA synthesis block by 5, 6-dichloro-1- β -D-ribofuranosylbenzimidazole (DRB) triggers p53-dependent apoptosis in human colon carcinoma cells. *Oncogene*, **18**, 5765-5772.
- Thrower, J.S., Hoffman, L., Rechsteiner, M. und Pickart, C.M. (2000) Recognition of the polyubiquitin proteolytic signal. *Embo J*, **19**, 94-102.
- Tsuge, T., Matsui, M. und Wei, N. (2001) The subunit 1 of the COP9 signalosome suppresses gene expression through its N-terminal domain and incorporates into the complex through the PCI domain. *J Mol Biol*, **305**, 1-9.
- Uhle, S., Medalia, O., Waldron, R., Dumdey, R., Henklein, P., Bech-Otschir, D., Huang, X., Berse, M., Sperling, J., Schade, R. und Dubiel, W. (2003) Protein

- kinase CK2 and protein kinase D are associated with the COP9 signalosome. *Embo J*, **22**, 1302-1312.
- Valverde, A.M., Sinnett-Smith, J., Van Lint, J. und Rozengurt, E. (1994) Molecular cloning and characterization of protein kinase D: a target for diacylglycerol and phorbol esters with a distinctive catalytic domain. *Proc Natl Acad Sci U S A*, **91**, 8572-8576.
- van den Heuvel, S. (2004) Protein degradation: CUL-3 and BTB--partners in proteolysis. *Curr Biol*, **14**, R59-61.
- Van Lint, J., Rykx, A., Maeda, Y., Vantus, T., Sturany, S., Malhotra, V., Vandenhede, J.R. und Seufferlein, T. (2002) Protein kinase D: an intracellular traffic regulator on the move. *Trends Cell Biol*, **12**, 193-200.
- Verma, R., Aravind, L., Oania, R., McDonald, W.H., Yates, J.R., 3rd, Koonin, E.V. und Deshaies, R.J. (2002) Role of Rpn11 metalloprotease in deubiquitination and degradation by the 26S proteasome. *Science*, **298**, 611-615.
- Vertommen, D., Rider, M., Ni, Y., Waelkens, E., Merlevede, W., Vandenhede, J.R. und Van Lint, J. (2000) Regulation of protein kinase D by multisite phosphorylation. Identification of phosphorylation sites by mass spectrometry and characterization by site-directed mutagenesis. *J Biol Chem*, **275**, 19567-19576.
- Waldron, R.T., Rey, O., Iglesias, T., Tugal, T., Cantrell, D. und Rozengurt, E. (2001) Activation loop Ser744 and Ser748 in protein kinase D are transphosphorylated in vivo. *J Biol Chem*, **276**, 32606-32615.
- Waldron, R.T. und Rozengurt, E. (2003) Protein kinase C phosphorylates protein kinase D activation loop Ser744 and Ser748 and releases autoinhibition by the pleckstrin homology domain. *J Biol Chem*, **278**, 154-163.
- Wang, Y. und Colicelli, J. (2001) RAS interaction with effector target RIN1. *Methods Enzymol*, **332**, 139-151.
- Warburg, O. und Christian, W. (1941) Isolierung und Kristallisation des Gärungsferments Enolase. *Biochem. Z.*, **310**, 384-423.
- Weber, J.D., Taylor, L.J., Roussel, M.F., Sherr, C.J. und Bar-Sagi, D. (1999) Nuclear Arf sequesters Mdm2 and activates p53. *Nat Cell Biol*, **1**, 20-26.
- Webley, K.M., Shorthouse, A.J. und Royds, J.A. (2000) Effect of mutation and conformation on the function of p53 in colorectal cancer. *J Pathol*, **191**, 361-367.
- Wee, S., Hetfeld, B., Dubiel, W. und Wolf, D.A. (2002) Conservation of the COP9/signalosome in budding yeast. *BMC Genet*, **3**, 15.
- Wei, N., Chamovitz, D.A. und Deng, X.W. (1994) Arabidopsis COP9 is a component of a novel signaling complex mediating light control of development. *Cell*, **78**, 117-124.
- Wei, N. und Deng, X.W. (1998) Characterization and purification of the mammalian COP9 complex, a conserved nuclear regulator initially identified as a repressor of photomorphogenesis in higher plants. *Photochem Photobiol*, **68**, 237-241.
- Wei, N., Tsuge, T., Serino, G., Dohmae, N., Takio, K., Matsui, M. und Deng, X.W. (1998) The COP9 complex is conserved between plants and mammals and is related to the 26S proteasome regulatory complex. *Curr Biol*, **8**, 919-922.
- Wertz, I.E., O'Rourke, K.M., Zhang, Z., Dornan, D., Arnott, D., Deshaies, R.J. und Dixit, V.M. (2004) Human De-etiololed-1 regulates c-Jun by assembling a CUL4A ubiquitin ligase. *Science*, **303**, 1371-1374.
- Wilson, M.P., Sun, Y., Cao, L. und Majerus, P.W. (2001) Inositol 1,3,4-trisphosphate 5/6-kinase is a protein kinase that phosphorylates the transcription factors c-Jun and ATF-2. *J Biol Chem*, **276**, 40998-41004.
- Winston, J.T., Strack, P., Beer-Romero, P., Chu, C.Y., Elledge, S.J. und Harper, J.W. (1999) The SCFbeta-TRCP-ubiquitin ligase complex associates specifically

- with phosphorylated destruction motifs in I κ B and β -catenin and stimulates I κ B ubiquitination in vitro. *Genes Dev*, **13**, 270-283.
- Wirkner, U., Voss, H., Lichter, P. und Pyerin, W. (1994) Human protein kinase CK2 genes. *Cell Mol Biol Res*, **40**, 489-499.
- Wolf, D.A., Zhou, C. und Wee, S. (2003) The COP9 signalosome: an assembly and maintenance platform for cullin ubiquitin ligases? *Nat Cell Biol*, **5**, 1029-1033.
- Wu, K., Chen, A. und Pan, Z.Q. (2000) Conjugation of Nedd8 to CUL1 enhances the ability of the ROC1-CUL1 complex to promote ubiquitin polymerization. *J Biol Chem*, **275**, 32317-32324.
- Wu, K., Yamoah, K., Dolios, G., Gan-Erdene, T., Tan, P., Chen, A., Lee, C.G., Wei, N., Wilkinson, K.D., Wang, R. und Pan, Z.Q. (2003) DEN1 is a dual function protease capable of processing the C terminus of Nedd8 and deconjugating hyper-neddylated CUL1. *J Biol Chem*, **278**, 28882-28891.
- Wu, X., Bayle, J.H., Olson, D. und Levine, A.J. (1993) The p53-mdm-2 autoregulatory feedback loop. *Genes Dev*, **7**, 1126-1132.
- Yaffe, M.B. und Elia, A.E. (2001) Phosphoserine/threonine-binding domains. *Curr Opin Cell Biol*, **13**, 131-138.
- Yang-Feng, T.L., Naiman, T., Kopatz, I., Eli, D., Dafni, N. und Canaani, D. (1994) Assignment of the human casein kinase II α' subunit gene (CSNK2A1) to chromosome 16p13.2-p13.3. *Genomics*, **19**, 173.
- Yaron, A., Gonen, H., Alkalay, I., Hatzubai, A., Jung, S., Beyth, S., Mercurio, F., Manning, A.M., Ciechanover, A. und Ben-Neriah, Y. (1997) Inhibition of NF- κ B cellular function via specific targeting of the I κ B-ubiquitin ligase. *Embo J*, **16**, 6486-6494.
- Yaron, A., Hatzubai, A., Davis, M., Lavon, I., Amit, S., Manning, A.M., Andersen, J.S., Mann, M., Mercurio, F. und Ben-Neriah, Y. (1998) Identification of the receptor component of the I κ B-ubiquitin ligase. *Nature*, **396**, 590-594.
- Yuan, J., Bae, D., Cantrell, D., Nel, A.E. und Rozengurt, E. (2002) Protein kinase D is a downstream target of protein kinase C θ . *Biochem Biophys Res Commun*, **291**, 444-452.
- Zandomeni, R.O. (1989) Kinetics of inhibition by 5,6-dichloro-1- β -D-ribofuranosylbenzimidazole on calf thymus casein kinase II. *Biochem J*, **262**, 469-473.
- Zhang, Y. und Xiong, Y. (1999) Mutations in human ARF exon 2 disrupt its nucleolar localization and impair its ability to block nuclear export of MDM2 and p53. *Mol Cell*, **3**, 579-591.
- Zhang, Y., Xiong, Y. und Yarbrough, W.G. (1998) ARF promotes MDM2 degradation and stabilizes p53: ARF-INK4a locus deletion impairs both the Rb and p53 tumor suppression pathways. *Cell*, **92**, 725-734.
- Zhong, W., Feng, H., Santiago, F.E. und Kipreos, E.T. (2003) CUL-4 ubiquitin ligase maintains genome stability by restraining DNA-replication licensing. *Nature*, **423**, 885-889.
- Zhou, C., Wee, S., Rhee, E., Naumann, M., Dubiel, W. und Wolf, D.A. (2003) Fission yeast COP9/signalosome suppresses cullin activity through recruitment of the deubiquitylating enzyme Ubp12p. *Mol Cell*, **11**, 927-938.