

Literaturverzeichnis

- [1] Smith, H.W. (1943): The evolution of the kidney.
- [2] Briggs, J. P.; Kriz, W und Schnermann, J. (2001): Overview of Renal Function and Structure, Greenberg, A, Primer on Kidney Diseases , 3. Auflage, Seite 3-20, National Kidney Foundation.
- [3] Hierholzer, K und Fromm, M (1995): Funktion der Nieren, Schmidt, RF und Thews, G, Physiologie des Menschen , 26. Auflage, Seite 719-776, Springer-Verlag, Berlin, Heidelberg, New York.
- [4] THURAU, K. und Schnermann, J. (1965): [THE SODIUM CONCENTRATION IN THE MACULA DENSA CELLS AS A REGULATING FACTOR FOR GLOMERULAR FILTRATION (MICROPUNCTURE EXPERIMENTS)], Klin.Wochenschr. (Band 43:410-3.), Seite 410-413.
- [5] Schnermann, J.; Traynor, T.; Yang, T.; Arend, L.; Huang, Y. G.; Smart, A. und Briggs, J. P. (1998): Tubuloglomerular feedback: new concepts and developments, Kidney Int.Suppl (Band 67:S40-5.), Seite S40-S45.
- [6] Thomson, S.; Bao, D.; Deng, A. und Vallon, V. (2000): Adenosine formed by 5'-nucleotidase mediates tubuloglomerular feedback, J Clin.Invest (Band 106), Nr. 2, Seite 289-298.
- [7] Schnermann, J. und Levine, D. Z. (2003): Paracrine factors in tubuloglomerular feedback: adenosine, ATP, and nitric oxide, Annu.Rev.Physiol (Band 65:501-29. Epub;%2002 May 01.), Seite 501-529.
- [8] Bell, P. D.; Lapointe, J. Y. und Peti-Peterdi, J. (2003): Macula densa cell signaling, Annu.Rev.Physiol (Band 65:481-500. Epub;%2003 Jan 09.), Seite 481-500.
- [9] Vallon, V. (2003): Tubuloglomerular feedback and the control of glomerular filtration rate, News Physiol Sci. (Band 18:169-74.), Seite 169-174.
- [10] Nishiyama, A.; Majid, D. S.; Walker, M., III; Miyatake, A. und Navar, L. G. (2001): Renal interstitial atp responses to changes in arterial pressure during alterations in tubuloglomerular feedback activity, Hypertension (Band 37), Nr. 2 Part 2, Seite 753-759.
- [11] Thomson, S. C.; Bachmann, S.; Bostanjoglo, M.; Ecelbarger, C. A.; Peterson, O. W.; Schwartz, D.; Bao, D. und Blantz, R. C. (1999): Temporal adjustment of the juxtaglomerular apparatus during sustained inhibition of proximal reabsorption, J.Clin.Invest (Band 104), Nr. 8, Seite 1149-1158.
- [12] Taugner, R.; Buhrlé, C. P.; Hackenthal, E.; Mannek, E. und Nobiling, R. (1984): Morphology of the juxtaglomerular apparatus and secretory mechanisms, Contrib.Nephrol. (Band 43:76-101.), Seite 76-101.
- [13] Bosse, H. M.; Bohm, R.; Resch, S. und Bachmann, S. (1995): Parallel regulation of constitutive NO synthase and renin at JGA of rat kidney under various stimuli, Am.J Physiol (Band 269), Nr. 6 Pt 2, Seite F793-F805.
- [14] Gallinat, O; Edling, O und Unger, Th (1999): AT1-Rezeptorantagonisten; Angiotensin II, ISBN: 3-7985-1171-3.
- [15] Wolf, G.; Butzmann, U. und Wenzel, U. O. (2003): The renin-angiotensin system and progression of renal disease: from hemodynamics to cell biology, Nephron Physiol (Band 93), Nr. 1, Seite 3-13. URL: PM:12411725
- [16] Campbell, D. J. (1996): Endogenous angiotensin II levels and the mechanism of action of angiotensin-converting enzyme inhibitors and angiotensin receptor type 1 antagonists, Clin.Exp Pharmacol.Physiol Suppl (Band 3:S125-31.), Seite S125-S131.
- [17] Van Dongen, R. und Peart, W. S. (1974): Calcium dependence of the inhibitory effect of angiotensin on renin secretion in the isolated perfused kidney of the rat, Br.J Pharmacol. (Band 50), Nr. 1, Seite 125-129.
- [18] Ackermann, M.; Ritthaler, T.; Rieger, G.; Kurtz, A. und Kramer, B. K. (1995): Endothelin inhibits cAMP-induced renin release from isolated renal juxtaglomerular cells, J Cardiovasc.Pharmacol. (Band 26 Suppl 3:S135-7.), Seite S135-S137.
- [19] Ritthaler, T.; Scholz, H.; Ackermann, M.; Rieger, G.; Kurtz, A. und Kramer, B. K. (1995): Effects of endothelins on renin secretion from isolated mouse renal juxtaglomerular cells, Am.J Physiol (Band 268), Nr. 1 Pt 2, Seite F39-F45.

- [20] Lorenz, J. N.; Weihprecht, H.; He, X. R.; Skott, O.; Briggs, J. P. und Schnermann, J. (1993): Effects of adenosine and angiotensin on macula densa-stimulated renin secretion, Am.J Physiol (Band 265), Nr. 2 Pt 2, Seite F187-F194.
- [21] Schnermann, J.; Weihprecht, H. und Briggs, J. P. (1990): Inhibition of tubuloglomerular feedback during adenosine1 receptor blockade, Am.J Physiol (Band 258), Nr. 3 Pt 2, Seite F553-F561.
- [22] Fray, J. C. (1978): Stretch receptor control of renin release in perfused rat kidney: effect of high perfusate potassium, J Physiol (Band 282:207-17.), Seite 207-217.
- [23] Carey, R. M.; McGrath, H. E.; Pentz, E. S.; Gomez, R. A. und Barrett, P. Q. (1997): Biomechanical coupling in renin-releasing cells, J Clin.Invest (Band 100), Nr. 6, Seite 1566-1574.
- [24] Kurtz, A.; Della, Bruna R.; Pratz, J. und Cavero, I. (1988): Rat juxtaglomerular cells are endowed with DA-1 dopamine receptors mediating renin release, J Cardiovasc.Pharmacol. (Band 12), Nr. 6, Seite 658-663.
- [25] Vandongen, R. (1976): Suppression of renin secretion in the isolated rat kidney by cycloheximide, Eur.J Pharmacol. (Band 40), Nr. 1, Seite 179-181.
- [26] Kurtz, A.; Muff, R.; Born, W.; Lundberg, J. M.; Millberg, B. I.; Gnadinger, M. P.; Uehlinger, D. E.; Weidmann, P.; Hokfelt, T. und Fischer, J. A. (1988): Calcitonin gene-related peptide is a stimulator of renin secretion, J Clin.Invest (Band 82), Nr. 2, Seite 538-543.
- [27] Jensen, B. L.; Schmid, C. und Kurtz, A. (1996): Prostaglandins stimulate renin secretion and renin mRNA in mouse renal juxtaglomerular cells, Am.J Physiol (Band 271), Nr. 3 Pt 2, Seite F659-F669.
- [28] Fang, F. C. (2004): Antimicrobial reactive oxygen and nitrogen species: concepts and controversies, Nat.Rev.Microbiol. (Band 2), Nr. 10, Seite 820-832.
- [29] Bachmann, S. und Mundel, P. (1994): Nitric oxide in the kidney: synthesis, localization, and function, Am.J Kidney Dis. (Band 24), Nr. 1, Seite 112-129. URL: PM:7517625
- [30] Plato, C. F.; Shesely, E. G. und Garvin, J. L. (2000): eNOS mediates L-arginine-induced inhibition of thick ascending limb chloride flux, Hypertension (Band 35), Nr. 1 Pt 2, Seite 319-323.
- [31] Wagner, C. und Kurtz, A. (1998): Regulation of renal renin release, Curr.Opin.Nephrol.Hypertens. (Band 7), Nr. 4, Seite 437-441.
- [32] Kurtz, A.; Gotz, K. H.; Hamann, M.; Kieninger, M. und Wagner, C. (1998): Stimulation of renin secretion by NO donors is related to the cAMP pathway, Am.J Physiol (Band 274), Nr. 4 Pt 2, Seite F709-F717.
- [33] Baker, M. E. (1995): Endocrine activity of plant-derived compounds: an evolutionary perspective, Proc Soc Exp Biol.Med. (Band 208), Nr. 1, Seite 131-138.
- [34] Botha, A.; Paul, I.; Roux, C.; Kock, J. L.; Coetzee, D. J.; Strauss, T. und Maree, C. (1999): An isolation procedure for arachidonic acid producing Mortierella species, Antonie Van Leeuwenhoek (Band 75), Nr. 3, Seite 253-256.
- [35] Kujubu, D. A.; Fletcher, B. S.; Varnum, B. C.; Lim, R. W. und Herschman, H. R. (1991): TIS10, a phorbol ester tumor promoter-inducible mRNA from Swiss 3T3 cells, encodes a novel prostaglandin synthase/cyclooxygenase homologue, J Biol.Chem. (Band 266), Nr. 20, Seite 12866-12872.
- [36] O'Banion, M. K.; Sadowski, H. B.; Winn, V. und Young, D. A. (1991): A serum- and glucocorticoid-regulated 4-kilobase mRNA encodes a cyclooxygenase-related protein, J Biol.Chem. (Band 266), Nr. 34, Seite 23261-23267.
- [37] Davies, N. M.; Good, R. L.; Roupe, K. A. und Yanez, J. A. (2004): Cyclooxygenase-3: axiom, dogma, anomaly, enigma or splice error?--Not as easy as 1, 2, 3, J.Pharm.Pharm.Sci. (Band 7), Nr. 2, Seite 217-226.
- [38] DeWitt, D. L. (1991): Prostaglandin endoperoxide synthase: regulation of enzyme expression, Biochim.Biophys.Acta (Band 1083), Nr. 2, Seite 121-134.
- [39] Yang, T.; Huang, Y.; Heasley, L. E.; Berl, T.; Schnermann, J. B. und Briggs, J. P. (2000): MAPK mediation of hypertonicity-stimulated cyclooxygenase-2 expression in renal medullary collecting duct cells, J Biol.Chem. (Band 275), Nr. 30, Seite 23281-23286.
- [40] Campean, V.; Theilig, F.; Paliege, A.; Breyer, M. und Bachmann, S. (2003): Key enzymes for renal prostaglandin synthesis: site-specific expression in rodent kidney (rat, mouse), Am.J Physiol Renal Physiol (Band 285), Nr. 1, Seite F19-F32. URL: PM:12657565
- [41] DeWitt, D. L. und Smith, W. L. (1988): Primary structure of prostaglandin G/H synthase from

- sheep vesicular gland determined from the complementary DNA sequence, Proc.Natl.Acad.Sci.U.S.A (Band 85), Nr. 5, Seite 1412-1416.
- [42] O'Neill, G. P. und Ford-Hutchinson, A. W. (1993): Expression of mRNA for cyclooxygenase-1 and cyclooxygenase-2 in human tissues, FEBS Lett. (Band 330), Nr. 2, Seite 156-160.
- [43] McKenna, J. A.; Zhang, M. Z.; Wang, J. L.; Cheng, H. und Harris, R. C. (1998): Constitutive expression of cyclooxygenase-2 in rat vas deferens, Am.J Physiol (Band 275), Nr. 1 Pt 2, Seite R227-R233.
- [44] Crofford, L. J. (1997): COX-1 and COX-2 tissue expression: implications and predictions, J.Rheumatol.Suppl (Band 49:15-9.), Seite 15-19.
- [45] Yang, T.; Park, J. M.; Arend, L.; Huang, Y.; Topaloglu, R.; Pasumarthi, A.; Praetorius, H.; Spring, K.; Briggs, J. P. und Schnermann, J. (2000): Low chloride stimulation of prostaglandin E2 release and cyclooxygenase-2 expression in a mouse macula densa cell line, J Biol.Chem. (Band 275), Nr. 48, Seite 37922-37929.
- [46] Peti-Peterdi, J.; Komlosi, P.; Fuson, A. L.; Guan, Y.; Schneider, A.; Qi, Z.; Redha, R.; Rosivall, L.; Breyer, M. D. und Bell, P. D. (2003): Luminal NaCl delivery regulates basolateral PGE2 release from macula densa cells, J.Clin.Invest (Band 112), Nr. 1, Seite 76-82.
- [47] Ichihara, A.; Imig, J. D. und Navar, L. G. (1999): Cyclooxygenase-2 modulates afferent arteriolar responses to increases in pressure, Hypertension (Band 34), Nr. 4 Pt 2, Seite 843-847.
- [48] Ichihara, A.; Imig, J. D.; Inscho, E. W. und Navar, L. G. (1998): Cyclooxygenase-2 participates in tubular flow-dependent afferent arteriolar tone: interaction with neuronal NOS, Am.J Physiol (Band 275), Nr. 4 Pt 2, Seite F605-F612.
- [49] Roig, F.; Llinas, M. T.; Lopez, R. und Salazar, F. J. (2002): Role of cyclooxygenase-2 in the prolonged regulation of renal function, Hypertension (Band 40), Nr. 5, Seite 721-728. URL: PM:12411468
- [50] Purdy, K. E. und Arendshorst, W. J. (1999): Prostaglandins buffer ANG II-mediated increases in cytosolic calcium in preglomerular VSMC, Am.J Physiol (Band 277), Nr. 6 Pt 2, Seite F850-F858.
- [51] Gerber, J. G.; Nies, A. S. und Olsen, R. D. (1981): Control of canine renin release: macula densa requires prostaglandin synthesis, J Physiol (Band 319:419-29.), Seite 419-429.
- [52] Baylis, C. (1987): Effects of administered thromboxanes on the intact, normal rat kidney, Ren Physiol (Band 10), Nr. 2, Seite 110-121.
- [53] Linas, S. L. (1984): Role of prostaglandins in renin secretion in the isolated kidney, Am.J Physiol (Band 246), Nr. 6 Pt 2, Seite F811-F818.
- [54] Harris, R. C.; Zhang, M. Z. und Cheng, H. F. (2004): Cyclooxygenase-2 and the renal renin-angiotensin system, Acta Physiol Scand. (Band 181), Nr. 4, Seite 543-547.
- [55] Yang, T.; Singh, I.; Pham, H.; Sun, D.; Smart, A.; Schnermann, J. B. und Briggs, J. P. (1998): Regulation of cyclooxygenase expression in the kidney by dietary salt intake, Am.J Physiol (Band 274), Nr. 3 Pt 2, Seite F481-F489.
- [56] Singh, I.; Grams, M.; Wang, W. H.; Yang, T.; Killen, P.; Smart, A.; Schnermann, J. und Briggs, J. P. (1996): Coordinate regulation of renal expression of nitric oxide synthase, renin, and angiotensinogen mRNA by dietary salt, Am.J Physiol (Band 270), Nr. 6 Pt 2, Seite F1027-F1037. URL: PM:8764322
- [57] Schricker, K.; Potzl, B.; Hamann, M. und Kurtz, A. (1996): Coordinate changes of renin and brain-type nitric-oxide-synthase (b-NOS) mRNA levels in rat kidneys, Pflugers Arch. (Band 432), Nr. 3, Seite 394-400.
- [58] Schricker, K.; Potzl, B.; Hamann, M. und Kurtz, A. (1996): Coordinate changes of renin and brain-type nitric-oxide-synthase (b-NOS) mRNA levels in rat kidneys, Pflugers Arch. (Band 432), Nr. 3, Seite 394-400.
- [59] Harris, R. C.; McKenna, J. A.; Akai, Y.; Jacobson, H. R.; Dubois, R. N. und Breyer, M. D. (1994): Cyclooxygenase-2 is associated with the macula densa of rat kidney and increases with salt restriction, J.Clin.Invest (Band 94), Nr. 6, Seite 2504-2510. URL: PM:7989609
- [60] Schricker, K.; Potzl, B.; Hamann, M. und Kurtz, A. (1996): Coordinate changes of renin and brain-type nitric-oxide-synthase (b-NOS) mRNA levels in rat kidneys, Pflugers Arch. (Band 432), Nr. 3, Seite 394-400.
- [61] Schricker, K.; Potzl, B.; Hamann, M. und Kurtz, A. (1996): Coordinate changes of renin and brain-type nitric-oxide-synthase (b-NOS) mRNA levels in rat kidneys, Pflugers Arch. (Band 432), Nr. 3, Seite 394-400.

- [62] Kammerl, M. C.; Richthammer, W.; Kurtz, A. und Kramer, B. K. (2002): Angiotensin II feedback is a regulator of renocortical renin, COX-2, and nNOS expression, Am.J.Physiol Regul.Integr.Comp Physiol (Band 282), Nr. 6, Seite R1613-R1617.
- [63] Schricker, K.; Hegyi, I.; Hamann, M.; Kaissling, B. und Kurtz, A. (1995): Tonic stimulation of renin gene expression by nitric oxide is counteracted by tonic inhibition through angiotensin II, Proc.Natl.Acad.Sci.U.S.A (Band 92), Nr. 17, Seite 8006-8010.
- [64] Schricker, K.; Hegyi, I.; Hamann, M.; Kaissling, B. und Kurtz, A. (1995): Tonic stimulation of renin gene expression by nitric oxide is counteracted by tonic inhibition through angiotensin II, Proc.Natl.Acad.Sci.U.S.A (Band 92), Nr. 17, Seite 8006-8010.
- [65] Hartner, A.; Goppelt-Struebe, M. und Hilgers, K. F. (1998): Coordinate expression of cyclooxygenase-2 and renin in the rat kidney in renovascular hypertension, Hypertension (Band 31), Nr. 1 Pt 2, Seite 201-205.
- [66] Mann, B.; Hartner, A.; Jensen, B. L.; Hilgers, K. F.; Hocherl, K.; Kramer, B. K. und Kurtz, A. (2001): Acute upregulation of COX-2 by renal artery stenosis, Am.J.Physiol Renal Physiol (Band 280), Nr. 1, Seite F119-F125.
- [67] Schricker, K.; Potzl, B.; Hamann, M. und Kurtz, A. (1996): Coordinate changes of renin and brain-type nitric-oxide-synthase (b-NOS) mRNA levels in rat kidneys, Pflugers Arch. (Band 432), Nr. 3, Seite 394-400.
- [68] Hartner, A.; Goppelt-Struebe, M. und Hilgers, K. F. (1998): Coordinate expression of cyclooxygenase-2 and renin in the rat kidney in renovascular hypertension, Hypertension (Band 31), Nr. 1 Pt 2, Seite 201-205.
- [69] Schricker, K.; Potzl, B.; Hamann, M. und Kurtz, A. (1996): Coordinate changes of renin and brain-type nitric-oxide-synthase (b-NOS) mRNA levels in rat kidneys, Pflugers Arch. (Band 432), Nr. 3, Seite 394-400.
- [70] Hartner, A.; Goppelt-Struebe, M. und Hilgers, K. F. (1998): Coordinate expression of cyclooxygenase-2 and renin in the rat kidney in renovascular hypertension, Hypertension (Band 31), Nr. 1 Pt 2, Seite 201-205.
- [71] Schricker, K.; Potzl, B.; Hamann, M. und Kurtz, A. (1996): Coordinate changes of renin and brain-type nitric-oxide-synthase (b-NOS) mRNA levels in rat kidneys, Pflugers Arch. (Band 432), Nr. 3, Seite 394-400.
- [72] Hartner, A.; Goppelt-Struebe, M. und Hilgers, K. F. (1998): Coordinate expression of cyclooxygenase-2 and renin in the rat kidney in renovascular hypertension, Hypertension (Band 31), Nr. 1 Pt 2, Seite 201-205.
- [73] Schricker, K.; Potzl, B.; Hamann, M. und Kurtz, A. (1996): Coordinate changes of renin and brain-type nitric-oxide-synthase (b-NOS) mRNA levels in rat kidneys, Pflugers Arch. (Band 432), Nr. 3, Seite 394-400.
- [74] Mann, B.; Hartner, A.; Jensen, B. L.; Kammerl, M.; Kramer, B. K. und Kurtz, A. (2001): Furosemide stimulates macula densa cyclooxygenase-2 expression in rats, Kidney Int. (Band 59), Nr. 1, Seite 62-68.
- [75] Schricker, K.; Potzl, B.; Hamann, M. und Kurtz, A. (1996): Coordinate changes of renin and brain-type nitric-oxide-synthase (b-NOS) mRNA levels in rat kidneys, Pflugers Arch. (Band 432), Nr. 3, Seite 394-400.
- [76] Schricker, K.; Potzl, B.; Hamann, M. und Kurtz, A. (1996): Coordinate changes of renin and brain-type nitric-oxide-synthase (b-NOS) mRNA levels in rat kidneys, Pflugers Arch. (Band 432), Nr. 3, Seite 394-400.
- [77] Wang, J. L.; Cheng, H. F.; Zhang, M. Z.; McKenna, J. A. und Harris, R. C. (1998): Selective increase of cyclooxygenase-2 expression in a model of renal ablation, Am.J.Physiol (Band 275), Nr. 4 Pt 2, Seite F613-F622.
- [78] Goncalves, A. R.; Fujihara, C. K.; Mattar, A. L.; Malheiros, D. M.; Noronha, Ide L.; de Nucci, G. und Zatz, R. (2004): Renal expression of COX-2, ANG II, and AT1 receptor in remnant kidney: strong renoprotection by therapy with losartan and a nonsteroidal anti-inflammatory, Am.J Physiol Renal Physiol (Band 286), Nr. 5, Seite F945-F954.
- [79] Szabo, A. J.; Wagner, L.; Erdely, A.; Lau, K. und Baylis, C. (2003): Renal neuronal nitric oxide synthase protein expression as a marker of renal injury, Kidney Int. (Band 64), Nr. 5, Seite 1765-1771.
- [80] Griffin, K. A.; Picken, M. M.; Churchill, M.; Churchill, P. und Bidani, A. K. (2000): Functional

- and structural correlates of glomerulosclerosis after renal mass reduction in the rat, J.Am.Soc.Nephrol. (Band 11), Nr. 3, Seite 497-506.
- [81] Weichert, W.; Paliege, A.; Provoost, A. P. und Bachmann, S. (2001): Upregulation of juxtaglomerular NOS1 and COX-2 precedes glomerulosclerosis in fawn-hooded hypertensive rats, Am.J.Physiol Renal Physiol (Band 280), Nr. 4, Seite F706-F714.
- [82] Welch, W. J.; Tojo, A.; Lee, J. U.; Kang, D. G.; Schnackenberg, C. G. und Wilcox, C. S. (1999): Nitric oxide synthase in the JGA of the SHR: expression and role in tubuloglomerular feedback, Am.J.Physiol (Band 277), Nr. 1 Pt 2, Seite F130-F138. URL: PM:10409306
- [83] Bagby, S. P.; McDonald, W. J. und Mass, R. D. (1979): Serial renin-angiotensin studies in spontaneously hypertensive and Wistar-Kyoto normotensive rats. Transition from normal- to high-renin status during the established phase of spontaneous hypertension, Hypertension (Band 1), Nr. 4, Seite 347-354.
- [84] Goodwin, D. C.; Landino, L. M. und Marnett, L. J. (1999): Effects of nitric oxide and nitric oxide-derived species on prostaglandin endoperoxide synthase and prostaglandin biosynthesis, FASEB J (Band 13), Nr. 10, Seite 1121-1136. URL: PM:10385604
- [85] Cheng, H. F.; Wang, J. L.; Zhang, M. Z.; McKenna, J. A. und Harris, R. C. (2000): Nitric oxide regulates renal cortical cyclooxygenase-2 expression, Am.J.Physiol Renal Physiol (Band 279), Nr. 1, Seite F122-F129.
- [86] Harris, R. C.; Cheng, H.; Wang, J.; Zhang, M. und McKenna, J. A. (2000): Interactions of the renin-angiotensin system and neuronal nitric oxide synthase in regulation of cyclooxygenase-2 in the macula densa, Acta Physiol Scand. (Band 168), Nr. 1, Seite 47-51. URL: PM:10691779
- [87] Theilig, F.; Campean, V.; Paliege, A.; Breyer, M.; Briggs, J. P.; Schnermann, J. und Bachmann, S. (2002): Epithelial COX-2 expression is not regulated by nitric oxide in rodent renal cortex, Hypertension (Band 39), Nr. 4, Seite 848-853.
- [88] Beierwaltes, W. H. (2002): Cyclooxygenase-2 products compensate for inhibition of nitric oxide regulation of renal perfusion, Am.J.Physiol Renal Physiol (Band 283), Nr. 1, Seite F68-F72.
- [89] Castrop, H.; Kammerl, M.; Mann, B.; Jensen, B. L.; Kramer, B. K. und Kurtz, A. (2000): Cyclooxygenase 2 and neuronal nitric oxide synthase expression in the renal cortex are not interdependent in states of salt deficiency, Pflugers Arch (Band 441), Nr. 2-3, Seite 235-240.
- [90] Vallon, V.; Traynor, T.; Barajas, L.; Huang, Y. G.; Briggs, J. P. und Schnermann, J. (2001): Feedback control of glomerular vascular tone in neuronal nitric oxide synthase knockout mice, J Am.Soc Nephrol. (Band 12), Nr. 8, Seite 1599-1606.
- [91] Choate, J. K.; Danson, E. J.; Morris, J. F. und Paterson, D. J. (2001): Peripheral vagal control of heart rate is impaired in neuronal NOS knockout mice, Am.J Physiol Heart Circ.Physiol (Band 281), Nr. 6, Seite H2310-H2317.
- [92] Barouch, L. A.; Harrison, R. W.; Skaf, M. W.; Rosas, G. O.; Cappola, T. P.; Kobeissi, Z. A.; Hobai, I. A.; Lemmon, C. A.; Burnett, A. L.; O'Rourke, B.; Rodriguez, E. R.; Huang, P. L.; Lima, J. A.; Berkowitz, D. E. und Hare, J. M. (2002): Nitric oxide regulates the heart by spatial confinement of nitric oxide synthase isoforms, Nature (Band 416), Nr. 6878, Seite 337-339.
- [93] Castrop, H.; Schweda, F.; Mizel, D.; Huang, Y.; Briggs, J.; Kurtz, A. und Schnermann, J. (2004): Permissive role of nitric oxide in macula densa control of renin secretion, Am.J Physiol Renal Physiol (Band 286), Nr. 5, Seite F848-F857. URL: PM:15075180
- [94] Ren, Y. L.; Garvin, J. L.; Ito, S. und Carretero, O. A. (2001): Role of neuronal nitric oxide synthase in the macula densa, Kidney Int. (Band 60), Nr. 5, Seite 1676-1683.
- [95] Wang, H.; Carretero, O. A. und Garvin, J. L. (2002): Nitric oxide produced by THAL nitric oxide synthase inhibits TGF, Hypertension (Band 39), Nr. 2 Pt 2, Seite 662-666.
- [96] Cheng, H. F.; Wang, J. L.; Zhang, M. Z.; Wang, S. W.; McKenna, J. A. und Harris, R. C. (2001): Genetic deletion of COX-2 prevents increased renin expression in response to ACE inhibition, Am.J.Physiol Renal Physiol (Band 280), Nr. 3, Seite F449-F456.
- [97] Yang, T.; Endo, Y.; Huang, Y. G.; Smart, A.; Briggs, J. P. und Schnermann, J. (2000): Renin expression in COX-2-knockout mice on normal or low-salt diets, Am.J Physiol Renal Physiol (Band 279), Nr. 5, Seite F819-F825.
- [98] Morham, S. G.; Langenbach, R.; Loftin, C. D.; Tiano, H. F.; Vouloumanos, N.; Jennette, J. C.; Mahler, J. F.; Kluckman, K. D.; Ledford, A.; Lee, C. A. und Smithies, O. (1995): Prostaglandin synthase 2 gene disruption causes severe renal pathology in the mouse, Cell (Band 83), Nr. 3, Seite 473-482.

- [99] Dinchuk, J. E.; Car, B. D.; Focht, R. J.; Johnston, J. J.; Jaffee, B. D.; Covington, M. B.; Contel, N. R.; Eng, V. M.; Collins, R. J.; Czerniak, P. M. und . (1995): Renal abnormalities and an altered inflammatory response in mice lacking cyclooxygenase II, *Nature* (Band 378), Nr. 6555, Seite 406-409.
- [100] Norwood, V. F.; Morham, S. G. und Smithies, O. (2000): Postnatal development and progression of renal dysplasia in cyclooxygenase-2 null mice, *Kidney Int.* (Band 58), Nr. 6, Seite 2291-2300.
- [101] Nath, K. A.; Chmielewski, D. H. und Hostetter, T. H. (1987): Regulatory role of prostanoids in glomerular microcirculation of remnant nephrons, *Am.J.Physiol* (Band 252), Nr. 5 Pt 2, Seite F829-F837.
- [102] Yoshida, Y.; Kawamura, T.; Ikoma, M.; Fogo, A. und Ichikawa, I. (1989): Effects of antihypertensive drugs on glomerular morphology, *Kidney Int.* (Band 36), Nr. 4, Seite 626-635.
- [103] Wang, J. L.; Cheng, H. F.; Shappell, S. und Harris, R. C. (2000): A selective cyclooxygenase-2 inhibitor decreases proteinuria and retards progressive renal injury in rats, *Kidney Int.* (Band 57), Nr. 6, Seite 2334-2342.
- [104] Zoja, C.; Perico, N.; Corna, D.; Benigni, A.; Gabanelli, M.; Morigi, M.; Bertani, T. und Remuzzi, G. (1990): Thromboxane synthesis inhibition increases renal prostacyclin and prevents renal disease progression in rats with remnant kidney, *J.Am.Soc.Nephrol.* (Band 1), Nr. 5, Seite 799-807.
- [105] Fogo, A. und Ichikawa, I. (1991): Evidence for a pathogenic linkage between glomerular hypertrophy and sclerosis, *Am.J.Kidney Dis.* (Band 17), Nr. 6, Seite 666-669.
- [106] Bachmann, S. und Ramasubbu, K. (1997): Immunohistochemical colocalization of the alpha-subunit of neutrophil NADPH oxidase and ecto-5'-nucleotidase in kidney and liver, *Kidney Int.* (Band 51), Nr. 2, Seite 479-482. URL: PM:9027726
- [107] Chabashvili, T.; Tojo, A.; Onozato, M. L.; Kitayakara, C.; Quinn, M. T.; Fujita, T.; Welch, W. J. und Wilcox, C. S. (2002): Expression and cellular localization of classic NADPH oxidase subunits in the spontaneously hypertensive rat kidney, *Hypertension* (Band 39), Nr. 2, Seite 269-274. URL: PM:11847196
- [108] Kalinina, N.; Agrotis, A.; Tararak, E.; Antropova, Y.; Kanellakis, P.; Ilyinskaya, O.; Quinn, M. T.; Smirnov, V. und Bobik, A. (2002): Cytochrome b558-dependent NAD(P)H oxidase-phox units in smooth muscle and macrophages of atherosclerotic lesions, *Arterioscler.Thromb.Vasc.Biol.* (Band 22), Nr. 12, Seite 2037-2043.
- [109] Griendling, K. K. und Ushio-Fukai, M. (1998): Redox control of vascular smooth muscle proliferation, *J Lab Clin.Med* (Band 132), Nr. 1, Seite 9-15.
- [110] Sanders, K. A.; Sundar, K. M.; He, L.; Dinger, B.; Fidone, S. und Hoidal, J. R. (2002): Role of components of the phagocytic NADPH oxidase in oxygen sensing, *J Appl.Physiol* (Band 93), Nr. 4, Seite 1357-1364.
- [111] Fukuyama, M.; Rokutan, K.; Sano, T.; Miyake, H.; Shimada, M. und Tashiro, S. (2005): Overexpression of a novel superoxide-producing enzyme, NADPH oxidase 1, in adenoma and well differentiated adenocarcinoma of the human colon, *Cancer Lett.* (Band 221), Nr. 1, Seite 97-104.
- [112] Gwinner, W. und Grone, H. J. (2000): Role of reactive oxygen species in glomerulonephritis, *Nephrol.Dial.Transplant.* (Band 15), Nr. 8, Seite 1127-1132.
- [113] Griendling, K. K.; Sorescu, D. und Ushio-Fukai, M. (2000): NAD(P)H oxidase: role in cardiovascular biology and disease, *Circ.Res.* (Band 86), Nr. 5, Seite 494-501.
- [114] Rajagopalan, S.; Kurz, S.; Munzel, T.; Tarpey, M.; Freeman, B. A.; Griendling, K. K. und Harrison, D. G. (1996): Angiotensin II-mediated hypertension in the rat increases vascular superoxide production via membrane NADH/NADPH oxidase activation. Contribution to alterations of vasomotor tone, *J Clin.Invest* (Band 97), Nr. 8, Seite 1916-1923.
- [115] Romero, F.; Herrera, J.; Nava, M. und Rodriguez-Iturbe, B. (1999): Oxidative stress in renal transplantation with uneventful postoperative course, *Transplant.Proc.* (Band 31), Nr. 6, Seite 2315-2316.
- [116] Vaziri, N. D.; Dicus, M.; Ho, N. D.; Boroujerdi-Rad, L. und Sindhu, R. K. (2003): Oxidative stress and dysregulation of superoxide dismutase and NADPH oxidase in renal insufficiency, *Kidney Int.* (Band 63), Nr. 1, Seite 179-185.
- [117] Zalba, G.; Beaumont, F. J.; San Jose, G.; Fortuno, A.; Fortuno, M. A.; Etayo, J. C. und Diez, J. (2000): Vascular NADH/NADPH oxidase is involved in enhanced superoxide production in

- spontaneously hypertensive rats, Hypertension (Band 35), Nr. 5, Seite 1055-1061.
- [118] Morrow, J. D. und Roberts, L. J. (1997): The isoprostanes: unique bioactive products of lipid peroxidation, Prog.Lipid Res. (Band 36), Nr. 1, Seite 1-21. URL: PM:9373618
- [119] Marnett, L. J.; Rowlinson, S. W.; Goodwin, D. C.; Kalgutkar, A. S. und Lanzo, C. A. (1999): Arachidonic acid oxygenation by COX-1 and COX-2. Mechanisms of catalysis and inhibition, J.Biol.Chem. (Band 274), Nr. 33, Seite 22903-22906. URL: PM:10438452
- [120] Beckman, J. S. und Koppenol, W. H. (1996): Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and ugly, Am.J Physiol (Band 271), Nr. 5 Pt 1, Seite C1424-C1437. URL: PM:8944624
- [121] Welch, W. J.; Tojo, A. und Wilcox, C. S. (2000): Roles of NO and oxygen radicals in tubuloglomerular feedback in SHR, Am.J Physiol Renal Physiol (Band 278), Nr. 5, Seite F769-F776. URL: PM:10807588
- [122] Vaziri, N. D.; Ni, Z. und Oveis, F. (1998): Upregulation of renal and vascular nitric oxide synthase in young spontaneously hypertensive rats, Hypertension (Band 31), Nr. 6, Seite 1248-1254. URL: PM:9622137
- [123] Schnackenberg, C. G. und Wilcox, C. S. (1999): Two-week administration of tempol attenuates both hypertension and renal excretion of 8-Iso prostaglandin f2alpha, Hypertension (Band 33), Nr. 1 Pt 2, Seite 424-428. URL: PM:9931141
- [124] Rodriguez-Iturbe, B.; Zhan, C. D.; Quiroz, Y.; Sindhu, R. K. und Vaziri, N. D. (2003): Antioxidant-rich diet relieves hypertension and reduces renal immune infiltration in spontaneously hypertensive rats, Hypertension (Band 41), Nr. 2, Seite 341-346. URL: PM:12574105
- [125] Vaziri, N. D.; Ni, Z.; Oveis, F. und Trnavsky-Hobbs, D. L. (2000): Effect of antioxidant therapy on blood pressure and NO synthase expression in hypertensive rats, Hypertension (Band 36), Nr. 6, Seite 957-964. URL: PM:11116107
- [126] Nava, M.; Quiroz, Y.; Vaziri, N. und Rodriguez-Iturbe, B. (2003): Melatonin reduces renal interstitial inflammation and improves hypertension in spontaneously hypertensive rats, Am.J.Physiol Renal Physiol (Band 284), Nr. 3, Seite F447-F454.
- [127] Welch, W. J.; Tojo, A. und Wilcox, C. S. (2000): Roles of NO and oxygen radicals in tubuloglomerular feedback in SHR, Am.J.Physiol Renal Physiol (Band 278), Nr. 5, Seite F769-F776. URL: PM:10807588
- [128] El Bekay, R.; Alvarez, M.; Monteseirin, J.; Alba, G.; Chacon, P.; Vega, A.; Martin-Nieto, J.; Jimenez, J.; Pintado, E.; Bedoya, F. J. und Sobrino, F. (2003): Oxidative stress is a critical mediator of the angiotensin II signal in human neutrophils: involvement of mitogen-activated protein kinase, calcineurin, and the transcription factor NF-kappaB, Blood (Band 102), Nr. 2, Seite 662-671.
- [129] Schmiedeberg, O (1883): Über die wirksamen bestandtheile der wurzel von *Apocynum cannabinum* L., Arch.Exp.Path.Pharm (Band 16), Seite 161-164.
- [130] Finnemore, H (1908): The constituents of Canadian hemp; Part I: Apocynin., J.Chem.Soc. (Band 93), Seite 1513-1520.
- [131] Otto, Th (1891): Über die Synthese des Acetovanillons aus Guajacol und Eisessig., Berichte (Band 24), Seite 2869-2870.
- [132] Hale, EM (1873): EDWIN M. HALE'S NEUE AMERIKANISCHE HEILMITTEL, Leipzig.
- [133] Basu, K; Dasgupta, B; Bhattacharya, S. K und Debnath, P. K (1971): Chemistry and pharmacology of apocynin, isolated from *Picrorhiza kurroa*, Royle ex Benth.Curr.Sci (Band 40), Nr. 22, Seite 603-604.
- [134] Welch, W. J.; Tojo, A. und Wilcox, C. S. (2000): Roles of NO and oxygen radicals in tubuloglomerular feedback in SHR, Am.J.Physiol Renal Physiol (Band 278), Nr. 5, Seite F769-F776. URL: PM:10807588
- [135] Meneton, P.; Ichikawa, I.; Inagami, T. und Schnermann, J. (2000): Renal physiology of the mouse, Am.J.Physiol Renal Physiol (Band 278), Nr. 3, Seite F339-F351.
- [136] Xue, B.; Pamidimukkala, J. und Hay, M. (2005): Sex differences in the development of angiotensin II-induced hypertension in conscious mice, Am.J.Physiol Heart Circ.Physiol (Band 288), Nr. 5, Seite H2177-H2184.
- [137] Huang, P. L.; Dawson, T. M.; Bredt, D. S.; Snyder, S. H. und Fishman, M. C. (1993): Targeted disruption of the neuronal nitric oxide synthase gene, Cell (Band 75), Nr. 7, Seite 1273-1286.

- [138] Beswick, R. A.; Zhang, H.; Marable, D.; Catravas, J. D.; Hill, W. D. und Webb, R. C. (2001): Long-term antioxidant administration attenuates mineralocorticoid hypertension and renal inflammatory response, *Hypertension* (Band 37), Nr. 2 Part 2, Seite 781-786. URL: PM:11230373
- [139] Rachmilewitz, D.; Okon, E. und Karmeli, F. (1997): Sulphydryl blocker induced small intestinal inflammation in rats: a new model mimicking Crohn's disease, *Gut* (Band 41), Nr. 3, Seite 358-365.
- [140] Cotter, M. A. und Cameron, N. E. (2003): Effect of the NAD(P)H oxidase inhibitor, apocynin, on peripheral nerve perfusion and function in diabetic rats, *Life Sci.* (Band 73), Nr. 14, Seite 1813-1824.
- [141] Park, Y. M.; Park, M. Y.; Suh, Y. L. und Park, J. B. (2004): NAD(P)H oxidase inhibitor prevents blood pressure elevation and cardiovascular hypertrophy in aldosterone-infused rats, *Biochem.Biophys.Res.Commun.* (Band 313), Nr. 3, Seite 812-817.
- [142] Kawai, J.; Ando, K.; Tojo, A.; Shimosawa, T.; Takahashi, K.; Onozato, M. L.; Yamasaki, M.; Ogita, T.; Nakaoka, T. und Fujita, T. (2004): Endogenous adrenomedullin protects against vascular response to injury in mice, *Circulation* (Band 109), Nr. 9, Seite 1147-1153.
- [143] Krege, J. H.; Hodgin, J. B.; Hagaman, J. R. und Smithies, O. (1995): A noninvasive computerized tail-cuff system for measuring blood pressure in mice, *Hypertension* (Band 25), Nr. 5, Seite 1111-1115.
- [144] Lykkegaard und Poulsen (1976): *Anal.Biochem* (Band 75), Seite 250-259.
- [145] Barton, M.; Vos, I.; Shaw, S.; Boer, P.; D'Uscio, L. V.; Grone, H. J.; Rabelink, T. J.; Lattmann, T.; Moreau, P. und Luscher, T. F. (2000): Dysfunctional renal nitric oxide synthase as a determinant of salt-sensitive hypertension: mechanisms of renal artery endothelial dysfunction and role of endothelin for vascular hypertrophy and Glomerulosclerosis, *J.Am.Soc.Nephrol.* (Band 11), Nr. 5, Seite 835-845.
- [146] Barton, M.; Vos, I.; Shaw, S.; Boer, P.; D'Uscio, L. V.; Grone, H. J.; Rabelink, T. J.; Lattmann, T.; Moreau, P. und Luscher, T. F. (2000): Dysfunctional renal nitric oxide synthase as a determinant of salt-sensitive hypertension: mechanisms of renal artery endothelial dysfunction and role of endothelin for vascular hypertrophy and Glomerulosclerosis, *J.Am.Soc.Nephrol.* (Band 11), Nr. 5, Seite 835-845.
- [147] Cheng, H. F.; Wang, S. W.; Zhang, M. Z.; McKenna, J. A.; Breyer, R. und Harris, R. C. (2002): Prostaglandins that increase renin production in response to ACE inhibition are not derived from cyclooxygenase-1, *Am.J.Physiol Regul.Integr.Comp Physiol* (Band 283), Nr. 3, Seite R638-R646.
- [148] Holmer, S. R.; Kaissling, B.; Putnik, K.; Pfeifer, M.; Kramer, B. K.; Riegger, G. A. und Kurtz, A. (1997): Beta-adrenergic stimulation of renin expression in vivo, *J Hypertens.* (Band 15), Nr. 12 Pt 1, Seite 1471-1479.
- [149] Krusell, L. R.; Christensen, C. K. und Pedersen, O. L. (1992): Alpha-adrenoceptor blockade in patients with mild to moderate hypertension: long-term renal effects of doxazosin, *J Cardiovasc.Pharmacol.* (Band 20), Nr. 3, Seite 440-444.
- [150] de Leeuw, P. W. und Birkenhager, W. H. (1988): Alpha-adrenoceptors and the kidney, *J Hypertens.Suppl* (Band 6), Nr. 2, Seite S21-S24.
- [151] de Leeuw, P. W.; van Es, P. N.; De Bos, R. und Birkenhager, W. H. (1986): Acute renal effects of doxazosin in man, *Br.J Clin.Pharmacol.* (Band 21 Suppl 1:41S-43S.), Seite 41S-43S.
- [152] Bauer, J. H. (1985): Adrenergic blocking agents and the kidney, *J Clin.Hypertens.* (Band 1), Nr. 3, Seite 199-221.
- [153] Goldberg, M. und Gehr, M. (1985): Effects of alpha-2 agonists on renal function in hypertensive humans, *J Cardiovasc.Pharmacol.* (Band 7 Suppl 8:S34-7.), Seite S34-S37.
- [154] SoRelle, R. (2000): National Heart, Lung, and Blood Institute halts part of antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT), *Circulation* (Band 101), Nr. 12, Seite E9025.
- [155] Cheng, H. F.; Wang, J. L.; Zhang, M. Z.; McKenna, J. A. und Harris, R. C. (2000): Role of p38 in the regulation of renal cortical cyclooxygenase-2 expression by extracellular chloride, *J.Clin.Invest* (Band 106), Nr. 5, Seite 681-688.
- [156] Ridley, S. H.; Dean, J. L.; Sarsfield, S. J.; Brook, M.; Clark, A. R. und Saklatvala, J. (1998): A p38 MAP kinase inhibitor regulates stability of interleukin-1-induced cyclooxygenase-2 mRNA, *FEBS Lett.* (Band 439), Nr. 1-2, Seite 75-80.

- [157] Hiller, G. und Sundler, R. (1999): Activation of arachidonate release and cytosolic phospholipase A2 via extracellular signal-regulated kinase and p38 mitogen-activated protein kinase in macrophages stimulated by bacteria or zymosan, *Cell Signal.* (Band 11), Nr. 12, Seite 863-869. URL: PM:10659994
- [158] Parfenova, H.; Haffner, J. und Leffler, C. W. (1999): Phosphorylation-dependent stimulation of prostanoid synthesis by nigericin in cerebral endothelial cells, *Am.J.Physiol* (Band 277), Nr. 4 Pt 1, Seite C728-C738.
- [159] Fowler, B. C.; Chang, Y. S.; Laamarti, A.; Higdon, M.; Lapointe, J. Y. und Bell, P. D. (1995): Evidence for apical sodium proton exchange in macula densa cells, *Kidney Int.* (Band 47), Nr. 3, Seite 746-751.
- [160] Theilig, F.; Bostanjoglo, M.; Pavenstadt, H.; Grupp, C.; Holland, G.; Slosarek, I.; Gressner, A. M.; Russwurm, M.; Koesling, D. und Bachmann, S. (2001): Cellular distribution and function of soluble guanylyl cyclase in rat kidney and liver, *J Am.Soc Nephrol.* (Band 12), Nr. 11, Seite 2209-2220.
- [161] Paliege, A.; Mizel, D.; Medina, C.; Pasumarthy, A.; Huang, Y. G.; Bachmann, S.; Briggs, J. P.; Schnermann, J. B. und Yang, T. (2004): Inhibition of nNOS expression in the macula densa by COX-2-derived prostaglandin E(2), *Am.J.Physiol Renal Physiol* (Band 287), Nr. 1, Seite F152-F159.
- [162] Cheng, H. F.; Wang, J. L.; Zhang, M. Z.; McKenna, J. A. und Harris, R. C. (2000): Nitric oxide regulates renal cortical cyclooxygenase-2 expression, *Am.J.Physiol Renal Physiol* (Band 279), Nr. 1, Seite F122-F129.
- [163] Southan, G. J. und Szabo, C. (1996): Selective pharmacological inhibition of distinct nitric oxide synthase isoforms, *Biochem Pharmacol.* (Band 51), Nr. 4, Seite 383-394.
- [164] Moore, P. K. und Bland-Ward, P. A. (1996): 7-nitroindazole: an inhibitor of nitric oxide synthase, *Methods Enzymol.* (Band 268:393-8.), Seite 393-398.
- [165] Vaziri, N. D. und Wang, X. Q. (1999): cGMP-mediated negative-feedback regulation of endothelial nitric oxide synthase expression by nitric oxide, *Hypertension* (Band 34), Nr. 6, Seite 1237-1241.
- [166] Bachmann, S.; Bosse, H. M. und Mundel, P. (1995): Topography of nitric oxide synthesis by localizing constitutive NO synthases in mammalian kidney, *Am.J Physiol* (Band 268), Nr. 5 Pt 2, Seite F885-F898. URL: PM:7539586
- [167] Kakoki, M.; Zou, A. P. und Mattson, D. L. (2001): The influence of nitric oxide synthase 1 on blood flow and interstitial nitric oxide in the kidney, *Am.J.Physiol Regul.Integr.Comp Physiol* (Band 281), Nr. 1, Seite R91-R97.
- [168] Wagner, C.; Godecke, A.; Ford, M.; Schnermann, J.; Schrader, J. und Kurtz, A. (2000): Regulation of renin gene expression in kidneys of eNOS- and nNOS-deficient mice, *Pflugers Arch.* (Band 439), Nr. 5, Seite 567-572. URL: PM:10764216
- [169] Persson, P. B.; Baumann, J. E.; Ehmke, H.; Hackenthal, E.; Kirchheim, H. R. und Nafz, B. (1993): Endothelium-derived NO stimulates pressure-dependent renin release in conscious dogs, *Am.J.Physiol* (Band 264), Nr. 6 Pt 2, Seite F943-F947.
- [170] Navarro, J.; Sanchez, A.; Saiz, J.; Ruilope, L. M.; Garcia-Estan, J.; Romero, J. C.; Moncada, S. und Lahera, V. (1994): Hormonal, renal, and metabolic alterations during hypertension induced by chronic inhibition of NO in rats, *Am.J.Physiol* (Band 267), Nr. 6 Pt 2, Seite R1516-R1521.
- [171] Tharaux, P. L.; Dussaule, J. C.; Pauti, M. D.; Vassitch, Y.; Ardaillou, R. und Chatziantoniou, C. (1997): Activation of renin synthesis is dependent on intact nitric oxide production, *Kidney Int.* (Band 51), Nr. 6, Seite 1780-1787.
- [172] Persson, P. B.; Baumann, J. E.; Ehmke, H.; Hackenthal, E.; Kirchheim, H. R. und Nafz, B. (1993): Endothelium-derived NO stimulates pressure-dependent renin release in conscious dogs, *Am.J.Physiol* (Band 264), Nr. 6 Pt 2, Seite F943-F947.
- [173] Deng, X.; Welch, W. J. und Wilcox, C. S. (1994): Renal vasoconstriction during inhibition of NO synthase: effects of dietary salt, *Kidney Int.* (Band 46), Nr. 3, Seite 639-646.
- [174] Schricker, K. und Kurtz, A. (1996): Blockade of nitric oxide formation inhibits the stimulation of the renin system by a low salt intake, *Pflugers Arch.* (Band 432), Nr. 2, Seite 187-191.
- [175] Ribeiro, M. O.; Antunes, E.; de Nucci, G.; Lovisolo, S. M. und Zatz, R. (1992): Chronic inhibition of nitric oxide synthesis. A new model of arterial hypertension, *Hypertension* (Band 20), Nr. 3, Seite 298-303.

- [176] Zanchi, A.; Schaad, N. C.; Osterheld, M. C.; Grouzmann, E.; Nussberger, J.; Brunner, H. R. und Waeber, B. (1995): Effects of chronic NO synthase inhibition in rats on renin-angiotensin system and sympathetic nervous system, Am.J.Physiol (Band 268), Nr. 6 Pt 2, Seite H2267-H2273.
- [177] Yamada, S. S.; Sasaki, A. L.; Fujihara, C. K.; Malheiros, D. M.; de Nucci, G. und Zatz, R. (1996): Effect of salt intake and inhibitor dose on arterial hypertension and renal injury induced by chronic nitric oxide blockade, Hypertension (Band 27), Nr. 5, Seite 1165-1172.
- [178] Hartlroad, J. Y.; Beharry, K. D.; Hausman, N.; Stavitsky, Y.; Asrat, T. und Modanlou, H. D. (2005): Effect of maternal administration of selective cyclooxygenase (COX)-2 inhibitors on renal size, growth factors, proteinases, and COX-2 secretion in the fetal rabbit, Biol.Neonate (Band 87), Nr. 4, Seite 246-253.
- [179] Guron, G. und Friberg, P. (2000): An intact renin-angiotensin system is a prerequisite for normal renal development, J.Hypertens. (Band 18), Nr. 2, Seite 123-137. URL: PM:10694179
- [180] Norwood, V. F.; Morham, S. G. und Smithies, O. (2000): Postnatal development and progression of renal dysplasia in cyclooxygenase-2 null mice, Kidney Int. (Band 58), Nr. 6, Seite 2291-2300.
- [181] Reese, J.; Paria, B. C.; Brown, N.; Zhao, X.; Morrow, J. D. und Dey, S. K. (2000): Coordinated regulation of fetal and maternal prostaglandins directs successful birth and postnatal adaptation in the mouse, Proc.Natl.Acad.Sci.U.S.A (Band 97), Nr. 17, Seite 9759-9764.
- [182] Komers, R.; Lindsley, J. N.; Oyama, T. T. und Anderson, S. (2004): Effects of long-term inhibition of neuronal nitric oxide synthase (NOS1) in uninephrectomized diabetic rats, Nitric.Oxide. (Band 11), Nr. 2, Seite 147-155.
- [183] Cosenzi, A.; Bernobich, E.; Bonavita, M.; Gris, F.; Odoni, G. und Bellini, G. (2002): Role of nitric oxide in the early renal changes induced by high fructose diet in rats, Kidney Blood Press Res. (Band 25), Nr. 6, Seite 363-369.
- [184] Schnermann, J. (1998): Juxtaglomerular cell complex in the regulation of renal salt excretion, Am.J Physiol (Band 274), Nr. 2 Pt 2, Seite R263-R279.
- [185] Sen, S.; Smeby, R. R. und Bumpus, F. M. (1972): Renin in rats with spontaneous hypertension, Circ.Res. (Band 31), Nr. 6, Seite 876-880.
- [186] Harbrecht, B. G.; Kim, Y. M.; Wirant, E. A.; Simmons, R. L. und Billiar, T. R. (1997): Timing of prostaglandin exposure is critical for the inhibition of LPS- or IFN-gamma-induced macrophage NO synthesis by PGE2, J Leukoc.Biol. (Band 61), Nr. 6, Seite 712-720.
- [187] Welch, W. J.; Tojo, A. und Wilcox, C. S. (2000): Roles of NO and oxygen radicals in tubuloglomerular feedback in SHR, Am.J.Physiol Renal Physiol (Band 278), Nr. 5, Seite F769-F776. URL: PM:10807588
- [188] Peti-Peterdi, J.; Komlosi, P.; Fuson, A. L.; Guan, Y.; Schneider, A.; Qi, Z.; Redha, R.; Rosivall, L.; Breyer, M. D. und Bell, P. D. (2003): Luminal NaCl delivery regulates basolateral PGE2 release from macula densa cells, J.Clin.Invest (Band 112), Nr. 1, Seite 76-82.
- [189] Traynor, T. R.; Smart, A.; Briggs, J. P. und Schnermann, J. (1999): Inhibition of macula densa-stimulated renin secretion by pharmacological blockade of cyclooxygenase-2, Am.J Physiol (Band 277), Nr. 5 Pt 2, Seite F706-F710.
- [190] Harding, P.; Sigmon, D. H.; Alfie, M. E.; Huang, P. L.; Fishman, M. C.; Beierwaltes, W. H. und Carretero, O. A. (1997): Cyclooxygenase-2 mediates increased renal renin content induced by low-sodium diet, Hypertension (Band 29), Nr. 1 Pt 2, Seite 297-302.
- [191] Kammerl, M. C.; Nusing, R. M.; Schweda, F.; Endemann, D.; Stubanus, M.; Kees, F.; Lackner, K. J.; Fischereder, M. und Kramer, B. K. (2001): Low sodium and furosemide-induced stimulation of the renin system in man is mediated by cyclooxygenase 2, Clin.Pharmacol.Ther. (Band 70), Nr. 5, Seite 468-474.
- [192] Ichikawa, I. und Fogo, A. (1996): Focal segmental glomerulosclerosis, Pediatr.Nephrol. (Band 10), Nr. 3, Seite 374-391.
- [193] D'Agati, V. (1994): The many masks of focal segmental glomerulosclerosis, Kidney Int. (Band 46), Nr. 4, Seite 1223-1241.
- [194] Cameron, J. S. (2003): Focal segmental glomerulosclerosis in adults, Nephrol.Dial.Transplant. (Band 18 Suppl 6:vi45-51.), Seite vi45-vi51.
- [195] Waldherr, R. und Gretz, N. (1988): Natural course of the development of histological lesions after 5/6 nephrectomy, Contrib.Nephrol. (Band 60:64-72.), Seite 64-72.

- [196] Welch, W. J.; Wilcox, C. S. und Thomson, S. C. (1999): Nitric oxide and tubuloglomerular feedback, *Semin.Nephrol.* (Band 19), Nr. 3, Seite 251-262.
- [197] Adler, S. und Huang, H. (2004): Oxidant stress in kidneys of spontaneously hypertensive rats involves both oxidase overexpression and loss of extracellular superoxide dismutase, *Am.J Physiol Renal Physiol* (Band 287), Nr. 5, Seite F907-F913.
- [198] Ulker, S.; McKeown, P. P. und Bayraktutan, U. (2003): Vitamins reverse endothelial dysfunction through regulation of eNOS and NAD(P)H oxidase activities, *Hypertension* (Band 41), Nr. 3, Seite 534-539.
- [199] Sen, S.; Smeby, R. R. und Bumpus, F. M. (1972): Renin in rats with spontaneous hypertension, *Circ.Res.* (Band 31), Nr. 6, Seite 876-880.
- [200] Morton, J. J.; Beattie, E. C.; Griffin, S. A.; MacPherson, F.; Lyall, F. und Russo, D. (1990): Vascular hypertrophy, renin and blood pressure in the young spontaneously hypertensive rat, *Clin.Sci.(Lond)* (Band 79), Nr. 5, Seite 523-530.
- [201] Welch, W. J.; Tojo, A. und Wilcox, C. S. (2000): Roles of NO and oxygen radicals in tubuloglomerular feedback in SHR, *Am.J.Physiol Renal Physiol* (Band 278), Nr. 5, Seite F769-F776. URL: PM:10807588
- [202] Xu, H.; Fink, G. D. und Galligan, J. J. (2002): Nitric oxide-independent effects of tempol on sympathetic nerve activity and blood pressure in DOCA-salt rats, *Am.J Physiol Heart Circ.Physiol* (Band 283), Nr. 3, Seite H885-H892. URL: PM:12181115
- [203] Zanzinger, J. und Czachurski, J. (2000): Chronic oxidative stress in the RVLM modulates sympathetic control of circulation in pigs, *Pflugers Arch.* (Band 439), Nr. 4, Seite 489-494. URL: PM:10678747
- [204] de Richelieu, L. T.; Sorensen, C. M.; Holstein-Rathlou, N. H. und Salomonsson, M. (2005): NO-independent mechanism mediates tempol-induced renal vasodilation in SHR, *Am.J Physiol Renal Physiol.* (Band 289), Nr. 6, Seite F1227-F1234.
- [205] Beswick, R. A.; Zhang, H.; Marable, D.; Catravas, J. D.; Hill, W. D. und Webb, R. C. (2001): Long-term antioxidant administration attenuates mineralocorticoid hypertension and renal inflammatory response, *Hypertension* (Band 37), Nr. 2 Part 2, Seite 781-786. URL: PM:11230373
- [206] Beswick, R. A.; Zhang, H.; Marable, D.; Catravas, J. D.; Hill, W. D. und Webb, R. C. (2001): Long-term antioxidant administration attenuates mineralocorticoid hypertension and renal inflammatory response, *Hypertension* (Band 37), Nr. 2 Part 2, Seite 781-786. URL: PM:11230373
- [207] Fortuno, A.; Olivan, S.; Beloqui, O.; San Jose, G.; Moreno, M. U.; Diez, J. und Zalba, G. (2004): Association of increased phagocytic NADPH oxidase-dependent superoxide production with diminished nitric oxide generation in essential hypertension, *J Hypertens.* (Band 22), Nr. 11, Seite 2169-2175.
- [208] Rodriguez-Iturbe, B.; Quiroz, Y.; Kim, C. H. und Vaziri, N. D. (2005): Hypertension induced by aortic coarctation above the renal arteries is associated with immune cell infiltration of the kidneys, *Am.J Hypertens.* (Band 18), Nr. 11, Seite 1449-1456.
- [209] Joffe, H. V. und Adler, G. K. (2005): Effect of aldosterone and mineralocorticoid receptor blockade on vascular inflammation, *Heart Fail.Rev.* (Band 10), Nr. 1, Seite 31-37.
- [210] Mervaala, E. M.; Muller, D. N.; Park, J. K.; Schmidt, F.; Lohn, M.; Breu, V.; Dragun, D.; Ganter, D.; Haller, H. und Luft, F. C. (1999): Monocyte infiltration and adhesion molecules in a rat model of high human renin hypertension, *Hypertension* (Band 33), Nr. 1 Pt 2, Seite 389-395.
- [211] Theuer, J.; Dechend, R.; Muller, D. N.; Park, J. K.; Fiebeler, A.; Barta, P.; Ganter, D.; Haller, H.; Dietz, R. und Luft, F. C. (2002): Angiotensin II induced inflammation in the kidney and in the heart of double transgenic rats, *BMC.Cardiovasc.Disord.* (Band 2:3. Epub;%2002 Jan 18.), Seite 3.
- [212] Abumiya, T.; Masuda, J.; Kawai, J.; Suzuki, T. und Ogata, J. (1996): Heterogeneity in the appearance and distribution of macrophage subsets and their possible involvement in hypertensive vascular lesions in rats, *Lab Invest* (Band 75), Nr. 2, Seite 125-136.
- [213] Muller, D. N.; Shagdarsuren, E.; Park, J. K.; Dechend, R.; Mervaala, E.; Hampich, F.; Fiebeler, A.; Ju, X.; Finckenberg, P.; Theuer, J.; Viedt, C.; Kreuzer, J.; Heidecke, H.; Haller, H.; Zenke, M. und Luft, F. C. (2002): Immunosuppressive treatment protects against angiotensin II-induced renal damage, *Am.J.Pathol.* (Band 161), Nr. 5, Seite 1679-1693.
- [214] Mervaala, E.; Finckenberg, P.; Lapatto, R.; Muller, D. N.; Park, J. K.; Dechend, R.; Ganter, D.; Vapaatalo, H. und Luft, F. C. (2003): Lipoic acid supplementation prevents angiotensin II-induced renal injury, *Kidney Int.* (Band 64), Nr. 2, Seite 501-508.

- [215] Muller, D. N.; Dechend, R.; Mervaala, E. M.; Park, J. K.; Schmidt, F.; Fiebeler, A.; Theuer, J.; Breu, V.; Ganten, D.; Haller, H. und Luft, F. C. (2000): NF-kappaB inhibition ameliorates angiotensin II-induced inflammatory damage in rats, *Hypertension* (Band 35), Nr. 1 Pt 2, Seite 193-201.
- [216] Feld, L. G.; Van Liew, J. B.; Galaske, R. G. und Boylan, J. W. (1977): Selectivity of renal injury and proteinuria in the spontaneously hypertensive rat, *Kidney Int.* (Band 12), Nr. 5, Seite 332-343.
- [217] Gil-Longo, J.; Fernandez-Grandal, D.; Alvarez, M.; Sieira, M. und Orallo, F. (1996): Study of in vivo and in vitro resting vasodilator nitric oxide tone in normotensive and genetically hypertensive rats, *Eur.J.Pharmacol.* (Band 310), Nr. 2-3, Seite 175-183.
- [218] Vaziri, N. D.; Ni, Z.; Oveis, F. und Trnavsky-Hobbs, D. L. (2000): Effect of antioxidant therapy on blood pressure and NO synthase expression in hypertensive rats, *Hypertension* (Band 36), Nr. 6, Seite 957-964.
- [219] Cowley, A. W., Jr.; Mori, T.; Mattson, D. und Zou, A. P. (2003): Role of renal NO production in the regulation of medullary blood flow, *Am.J.Physiol Regul.Integr.Comp Physiol* (Band 284), Nr. 6, Seite R1355-R1369.
- [220] Cowley, A. W., Jr.; Mori, T.; Mattson, D. und Zou, A. P. (2003): Role of renal NO production in the regulation of medullary blood flow, *Am.J.Physiol Regul.Integr.Comp Physiol* (Band 284), Nr. 6, Seite R1355-R1369.
- [221] Sullivan, JC; Williams, JM; Pollock, DM und Pollock, JS (2002): Effect of Apocynin on NOS Activity and Expression in Mesenteric Arteries in DOCA-salt Hypertension, *FASEB J.* (Band 16), Nr. 4, Seite 96-96.
- [222] Asaba, K.; Tojo, A.; Onozato, M. L.; Goto, A.; Quinn, M. T.; Fujita, T. und Wilcox, C. S. (2005): Effects of NADPH oxidase inhibitor in diabetic nephropathy, *Kidney Int.* (Band 67), Nr. 5, Seite 1890-1898.
- [223] Kobayashi, N.; Honda, T.; Yoshida, K.; Nakano, S.; Ohno, T.; Tsubokou, Y. und Matsuoka, H. (2005): Critical role of bradykinin-eNOS and oxidative stress-LOX-1 pathway in cardiovascular remodeling under chronic angiotensin-converting enzyme inhibition, *Atherosclerosis*. (Band .).
- [224] Limas, C. J. und Limas, C. (1977): Prostaglandin metabolism in the kidneys of spontaneously hypertensive rats, *Am.J Physiol* (Band 233), Nr. 1, Seite H87-H92. URL: PM:18024
- [225] Hocherl, K.; Endemann, D.; Kammerl, M. C.; Grobecker, H. F. und Kurtz, A. (2002): Cyclooxygenase-2 inhibition increases blood pressure in rats, *Br.J.Pharmacol.* (Band 136), Nr. 8, Seite 1117-1126.
- [226] Lee, S. A.; Park, J. K.; Kang, E. K.; Bae, H. R.; Bae, K. W. und Park, H. T. (2000): Calmodulin-dependent activation of p38 and p42/44 mitogen-activated protein kinases contributes to c-fos expression by calcium in PC12 cells: modulation by nitric oxide, *Brain Res.Mol.Brain Res.* (Band 75), Nr. 1, Seite 16-24.
- [227] Takeda, Y.; Miyamori, I. und Takeda, R. (1995): Significance of 19-noraldosterone, a new mineralocorticoid, in clinical and experimental hypertension, *Steroids*. (Band 60), Nr. 1, Seite 137-142.
- [228] Hocherl, K.; Kammerl, M.; Kees, F.; Kramer, B. K.; Grobecker, H. F. und Kurtz, A. (2002): Role of renal nerves in stimulation of renin, COX-2, and nNOS in rat renal cortex during salt deficiency, *Am.J Physiol Renal Physiol.* (Band 282), Nr. 3, Seite F478-F484.
- [229] Schricker, K.; Holmer, S.; Kramer, B. K.; Riegger, G. und Kurtz, A. (1994): Control of renin gene expression in 2 kidney-1 clip rats, *Kidney Int.* (Band 46), Nr. 6, Seite 1539-1541.
- [230] Cheng, H. F.; Wang, J. L.; Zhang, M. Z.; Miyazaki, Y.; Ichikawa, I.; McKenna, J. A. und Harris, R. C. (1999): Angiotensin II attenuates renal cortical cyclooxygenase-2 expression, *J.Clin.Invest* (Band 103), Nr. 7, Seite 953-961.
- [231] Harris, R. C.; Cheng, H.; Wang, J.; Zhang, M. und McKenna, J. A. (2000): Interactions of the renin-angiotensin system and neuronal nitric oxide synthase in regulation of cyclooxygenase-2 in the macula densa, *Acta Physiol Scand.* (Band 168), Nr. 1, Seite 47-51.
- [232] Sen, S.; Smeby, R. R. und Bumpus, F. M. (1972): Renin in rats with spontaneous hypertension, *Circ.Res.* (Band 31), Nr. 6, Seite 876-880.
- [233] Morton, J. J.; Beattie, E. C.; Griffin, S. A.; MacPherson, F.; Lyall, F. und Russo, D. (1990): Vascular hypertrophy, renin and blood pressure in the young spontaneously hypertensive rat, *Clin.Sci.(Lond)* (Band 79), Nr. 5, Seite 523-530.

- [234] Sen, S.; Smeby, R. R. und Bumpus, F. M. (1972): Renin in rats with spontaneous hypertension, Circ.Res. (Band 31), Nr. 6, Seite 876-880.
- [235] Kurtz, A. und Wagner, C. (1998): Role of nitric oxide in the control of renin secretion, Am.J Physiol (Band 275), Nr. 6 Pt 2, Seite F849-F862. URL: PM:9843901
- [236] Meneton, P.; Ichikawa, I.; Inagami, T. und Schnermann, J. (2000): Renal physiology of the mouse, Am.J.Physiol Renal Physiol (Band 278), Nr. 3, Seite F339-F351.
- [237] OKAMOTO, K. (1972): Spontaneous Hypertension, Igaku Shoin Ltd, Tokyo.
- [238] Welch, W. J.; Tojo, A. und Wilcox, C. S. (2000): Roles of NO and oxygen radicals in tubuloglomerular feedback in SHR, Am.J.Physiol Renal Physiol (Band 278), Nr. 5, Seite F769-F776. URL: PM:10807588