

## 6 List of References

- Aasland R., Akslen L.A., Varhaug J.E., Lillehaug J.R. (1990).  
Co-expression of the genes encoding transforming growth factor-alpha and its receptor in papillary carcinoma of the thyroid.  
*Int. J. Cancer* 46: 382- 387.
- Akslen L. A. and Varhaug J. E. (1995)  
Oncoproteins and tumor progression in papillary thyroid carcinoma.  
*Cancer* 76: 1643- 1654.
- Allgeier A., Offermans S., Van Sande J., Spicher K., Schultz G., Dumont J.E. (1994).  
The human thyrotropin receptor activates G-proteins G<sub>s</sub> and G<sub>q/11</sub>.  
*J. Biol. Chem.* 269: 13733- 13735.
- Ambesi-Impiombato F.S., Parks L.A.M., Coon H.G. (1980)  
Culture of hormone-dependent functional epithelial cells from rat thyroids.  
*Proc. Natl. Acad. Sci. USA* 77: 3455- 3459.
- Amorino G.P., Hamilton V.M., Valerie K., Dent P., Lammering G., Schmidt-Ullrich R.K. (2002).  
Epidermal growth factor receptor dependence of radiation- induced transcription factor activation in human breast carcinoma cells.  
*Mol. Biol. Cell* 13: 2233- 2244.
- Armstrong R., Wen W., Meinkoth J., Taylor S., Montminy M. (1995).  
A refractory phase in cyclic AMP- responsive transcription requires down regulation of protein kinase A.  
*Mol. Cell. Biol.* 15: 1826- 1832.
- Atkinson S. and Kendall-Taylor P. (1987).  
Effect of thyrotropin on epidermal growth factor receptors in monolayer cultures of porcine thyroid cells.  
*J. Endocrinol.* 114: 179- 184.
- Becks G.P., Logan A., Phillips I.D., Wang J.F., Smith C., DeSousa D., Hill D.J. (1994).  
Increase of basic FGF and FGF receptor mRNA during rat hyperplasia: temporal changes and cell distribution.  
*J. Endocrinol.* 142: 325- 338.
- Belfiore A., Pandini G., Vella V., Frasca F., Pezzino V., Vigneri R. (1999).  
Insulin/IGF-I hybrid receptors play a major role in IGF-I signaling in thyroid cancer.  
26<sup>th</sup> Annual Meeting of the European Thyroid Association, Milan 1999.

- Berg V., Vassart G., Christophe D. (1996).  
Identification of a thyroid-specific and cAMP-responsive enhancer in the upstream sequences of the human thyroglobulin promoter.  
*Biochim. Biophys. Acta* 1307: 35- 38.
- Bertherat J., chanson P., Montminy m. (1995).  
The Cyclic Adenosine 3',5'- Monophosphate-Responsive Factor CREB is Constitutively activated in Human Somatotroph Adenomas.  
*Mol. Endocrinol.* 9: 777- 783.
- Bifulco M., Perillo B., Saji M., Laezza C., Tedesco I., Kohn L.D., Aloj S.M. (1995).  
Regulation of the 3-hydroxy-3-methylglutaryl coenzyme A reductase gene expression in FRTL-5 cells. I. Identification and characterization of a cyclic AMP-responsive element in the rat reductase promoter.  
*J. Biol. Chem.* 270: 15231- 15236.
- Blendy J.A., Kaestner K.H., Schmid W., Gass P., Schütz G. (1996).  
Targeting of the CREB gene leads to up-regulation of a novel CREB mRNA isoform.  
*EMBO J.* 15: 1098- 1106.
- Brenner-Gati L., Berg K.A., Gershengorn M.C. (1989).  
Insulin-like growth factor-I potentiates thyrotropin stimulation of adenylyl cyclase in FRTL-5 cells.  
*Endocrinology* 125: 1315- 1320.
- Brindle P., Nakajima T., Montminy M. (1995).  
Multiple protein kinase A-regulated events are required for transcriptional induction by cAMP.  
*Proc. Natl. Acad. Sci. USA* 92: 10521- 10525.
- Broecker M., Hammer J., Derwahl M. (1998).  
Excessive activation of tyrosine kinases leads to inhibition of proliferation in a thyroid carcinoma cell line.  
*Life Sciences* 63: 2373- 2386.
- Broecker M., Mayr G.W., Derwahl M. (1997).  
Suppression of Thyrotropin Receptor-G Protein- Phospholipase C Coupling by Activation of Protein Kinase C in Thyroid Carcinoma Cells.  
*Endocrinology* 138: 3787- 3796.
- Bubenhofer R. and Hedinger C. (1977).  
Thyroid neoplasms before and after the prophylactic supplementation of table salt with iodine.  
*Schweiz. Med. Wochenschr.* 107: 733- 741.

- Burikhanov R., Coulonval K., Pirson I., Lamy F., Dumont J.E., Roger P.P. (1996).  
Thyrotropin via cyclic AMP induces insulin receptor expression and insulin co-stimulation of growth and amplifies insulin and insulin-like growth factor signaling pathways in dog thyroid epithelial cells.  
*J. Biol. Chem.* 271: 29400- 29406.
- Cheng G., Lewis A.E., Meinkoth J.L. (2003).  
Ras stimulates aberrant cell cycle progression and apoptosis in rat thyroid cells.  
*Mol. Endocrinol.* *17*: 450-459.
- Cheung N.W., Lou J.C., Boyages S.C. (1996).  
GH does not increase thyroid size in the absence of thyrotropin: a study in adults with hypopituitarism.  
*J. Clin. Endocrinol. Metab.* 81: 1179- 1183.
- Chijiwa T., Mishima A., Hagiwara M., Sano M., Hayashi K., Inoue T., Naito K., Toshioka T., Hidaka H. (1990).  
Inhibition of Forskolin-induced Neurite Outgrowth and Protein phosphorylation by a newly Synthesized Selective Inhibitor of Cyclic AMP-dependent Protein Kinase, *N*-[2-(*p*-Bromocinnamylamino)ethyl]-5-isoquinolinesulphonamide (H-89), of PC12D Pheochromocytoma Cells.  
*J. Biol. Chem.* 265: 5267- 5272.
- Ciardiello F., and Tortora G. (1998).  
Interactions between the epidermal growth factor receptor and type I protein kinase A: biological significance and therapeutic implications.  
*Clin. Cancer Res.* 4: 821- 828.
- Clark R. (1997).  
The somatogenic hormones and insulin-like growth factor-1: stimulators of lymphopoiesis and immune function.  
*Endocr. Rev.* 18: 157- 179.
- Contessa J.N., Reardon D.B., Todd D., Dent P., Mikkelsen R.B., Valerie K., Bowers G.D., Schmidt-Ullrich R.K. (1999).  
The inducible expression of dominant- negative epidermal growth factor receptor-CD533 results in radiosensitization of human mammary carcinoma cells.  
*Clin. Cancer Res.* 5: 405- 411.
- Deak M., Clifton A.D., Lucocq J.M., Alessi D.R. (1998).  
Mitogen- and stress-activated protein kinase-1 (MSK1) is directly activated by MAPK and SAPK2/p38, and may mediate activation of CREB.  
*EMBO J.* 17: 4426- 4441.
- De Cesare D., Jaquot S., Hanauer A., Sassone-Corsi P. (1998).  
Rsk-2 activity is necessary for epidermal growth factor-induced phosphorylation of CREB protein and transcription of c-fos gene.  
*PNAS* 95: 12202- 12207.

- De Groot R.P., den Hertog J., Vandenheede J.R., Goris J., Sassone-Corsi P. (1993).  
Multiple and cooperative phosphorylation events regulate the CREM activator  
function.  
EMBO J. 12: 3903- 3911..
- Dennings M.F., Dlugosz A.A., Threadgill D.W., Magnuson T., Yuspa S.H. (1996).  
Activation of the Epidermal Growth Factor Receptor Signal Transduction Pathway  
Stimulates Tyrosine Phosphorylation of Protein Kinase C  $\delta$ .  
J. Biol. Chem. 271: 5325- 5331.
- Dent P., Haser W., Haystead T.A., Vincent L.A., Roberts T.M., Sturgill T.W. (1992).  
Activation of mitogen-activated protein kinase kinase by v-Raf in NIH 3T3 cells and  
*in vitro*.  
Science 257: 1404-1407.
- Dent P., Reardan D.B., Park J.S., Bowers G., Logsdon C., Valerie K., Schmidt-Ullrich R. (1999).  
Radiation- induced release of transforming growth factor alpha activates the  
epidermal growth factor receptor and mitogen- activated protein kinase pathway in  
carcinoma cells, leading to increased proliferation and protection from radiation-  
induced cell death.  
Mol. Biol. Cell. 10: 2493- 2506.
- Derwahl M., Broecker M., Kraiem Z. (1999).  
Thyrotropin may not be the dominant growth factor in benign and malignant  
thyroid tumors.  
J. Endocrinol. Metab. 84: 829- 833.
- Derwahl M., Hamacher C., Russo D., Broecker M., Manole D., Schatz H., Kopp P., Filetti S.  
(1996).  
Constitutive activation of the Gs alpha protein- adenylate cyclase pathway may not  
be sufficient to generate toxic thyroid adenomas.  
J. Clin. Endocrinol. Metab.: 81: 1898- 1904.
- Derwahl M., Kuemmel M., Goretzki P., Schatz H., Broecker M. (1993).  
Expression of the human TSH receptor in a human thyroid carcinoma cell line that  
lacks an endogenous TSH receptor: growth inhibition by cAMP.  
Biochem. Biophys. Res. Commun. 191: 1131- 1138.
- Derwahl M., Manole D., Sobke A., Broecker M. (1998).  
Pathogenesis of toxic thyroid adenomas and nodules: relevance of activating  
mutations in the TSH-receptor and Gs-alpha gene, the possible role of iodine  
deficiency and secondary and TSH-independent molecular mechanisms.  
Exp. Clin. Endocrinol. Diabetes Suppl. 4: S6- S9.
- Derwahl M. and Studer H. (1998).  
Pathogenesis and treatment of multinodular goiter.  
In: Fagin J. A. (Ed.): Thyroid cancer. Kluwer, Boston, 155-186.

Desdouets C., Matesic G., Molina C.A., Foulkes N.S., Sassone-Corsi P., Brechot C., Sobczak-Thepot J. (1995).

Cell cycle regulation of cyclin A gene expression by the cyclic AMP-responsive transcription factors CREB and CREM.

Mol. Cell. Biol. 15: 3301- 3309.

Di Carlo A., Mariano A., Pisrano A., Parmeggiani U., Beguinot L., Macchia V. (1990).

Epidermal growth factor receptor and thyrotropin response in human thyroid tissues.

J. Endocrinol. Invest. 13: 293- 299.

Dumont J.E., Maenhaut C., Pirson I., Baptist M., Roger P.P. (1991).

Growth factors controlling the thyroid gland.

Clin. Endocrinol. Metab. 5: 727- 754.

Dumont J. E., Lamy F., Roger P., Maenhaut C. (1992),

Physiological and pathological regulation of thyroid cell proliferation and differentiation by thyrotropin and other factors.

Physiol. Rev. 72: 667- 697.

Dutil E. M., Toker A., Newton A. C. (1998).

Regulation of conventional protein kinase C isozymes by phosphoinositide-dependent kinase 1 (PDK-1).

Curr. Biol. 8: 1366- 1375.

Eggo M.C., Bachrach L.K., Burrow G.N. (1990).

Interaction of TSH, insulin and insulin-like growth factors in regulating thyroid growth and function.

Growth Factors 2: 99- 109.

Eggo M.C., Bachrach L.K., Fayet G., Burrow G.N. (1984).

The effects of growth factors and serum on DNA synthesis and differentiation in thyroid cells in culture.

Mol. Cell. Endocrinol. 38: 141- 150.

Eggo M.C., King W.J., Black E.G., Sheppard M.C. (1996).

Functional human thyroid cells and their insulin-like growth factor-binding proteins: regulation by thyrotropin, cyclic 3', 5' adenosine monophosphate, and growth factors.

J. Clin. Endocrinol. Metab. 81: 3056-3062.

Eggo M.C., Pratt C., Becks G.P., Burrow G.N. (1989).

Regulation of growth and differentiation in follicle cells.

Adv. Exp. Med. Biol. 261: 341- 356.

- Endo T., Shimura H., Saito T., Onaya T. (1990).  
Cloning of malignantly transformed rat thyroid (FRTL) cells with thyrotropin receptors and their growth inhibition by 3', 5'-cyclic adenosine monophosphate. *Endocrinology* 126: 1492- 1497.
- Feliars D. and Pavlovic-Hournac M. (1994).  
Species differences of the thyroid protein kinase C heterogeneity. *Thyroid* 4: 459- 465.
- Frauman A.G., Moses A.C. (1990).  
Oncogenes and growth factors in thyroid carcinogenesis. *Endocrinol. Metab. Clin. North Am.* 19: 479-493.
- Gärtner R., Bechtner G., Stübner D., Greil W. (1990).  
Growth regulation of porcine thyroid follicles in vitro by growth factors. *Horm. Metab. Res. Suppl.* 23: 225- 230.
- Gärtner R., Greil W., Demhardt R., Horn K. (1985).  
Involvement of cyclic AMP, iodine and metabolites of arachidonic acid in the regulation of cell proliferation of isolated porcine thyroid follicles. *Mol. Cell. Endocrinol.* 42: 145- 155.
- Girardet C., Walker H.W., Habener J.F. (1996).  
An alternatively spliced polycistronic mRNA encoding cyclic adenosine 3', 5'-monophosphate (cAMP)-responsive transcription factor CREB (cAMP response element-binding protein) in human testis extinguishes expression of an internally translated inhibitor CREB isoform. *Mol. Endocrinology* 10: 879- 891.
- Gonzalez G.A., Yamamoto K.K., Fischer W.H., Karr D., Menzel P., Biggs III W., Vale W.W., Montminy M.R. (1989).  
A cluster of phosphorylation sites on the cyclic AMP-regulated factor CREB predicted by its sequence. *Nature* 337: 749- 752.
- Goretzki P.E., Frilling A., Simon D., Roehrer D. (1990).  
Growth regulation of normal thyroids and thyroid tumors in man. *Recent Results Cancer Res.* 118: 48- 63.
- Gudi T., Casteel D.E., Vinson C., Boss G.R., Pilz R.B. (2000).  
NO activation of fos promoter elements requires nuclear translocation of G-kinase I and CREB phosphorylation but is independent of MAP kinase activation. *Oncogene* 19: 6324- 6333.
- Hagiwara M., Brindle P., Harootunian A., Armstrong R., Rivier J., Vale W., Tsien R., Montminy M.R. (1993).  
Coupling of hormonal stimulation and transcription via the cyclic AMP- responsive factor CREB is rate limited by nuclear entry of protein kinase A. *Mol. Cell. Biol.* 13: 4852- 4859.

- Hai T. and Curran T. (1991).  
Cross-family dimerization of transcription factors Fos/Jun and ATF/CREB alters DNA binding specificity.  
Proc. Natl. Acad. Sci. USA 88: 3720- 3724.
- Hamacher C., Studer H., Zbaeren J., Schatz H., Derwahl M. (1995)  
Expression of functional stimulatory guanine nucleotide binding protein in nonfunctioning thyroid adenomas is not correlated to adenylate cyclase and growth of these tumors.  
J. Clin. Endocrinol. Metab.: 80: 1724- 1732.
- Hatada T., Sakarone Y., Kusunoki M., Kobayashi A., Utsunomiya J. (1992).  
Protein kinase C activity in human thyroid carcinoma and adenoma.  
Cancer 70: 2918- 2922.
- Haugen D. R. F., Akslen L. A., Varhaug J. E., Lillehaug J. R. (1993).  
Demonstration of a TGF $\alpha$ -EGF-receptor autocrine loop and c-myc protein over-expression in papillary thyroid carcinomas.  
Int. J. Cancer 55: 37- 43.
- Heldin N.-E., Cvejic D., Smeds S., Westermark B. (1991).  
Coexpression of functionally active receptors for thyrotropin and platelet-derived growth factor in human thyroid carcinoma cells.  
Endocrinology 129: 2187- 2193.
- Heldin N.-E., Gustavsson G., Claesson-Welsh L. et al. (1988).  
Aberrant expression of receptors for platelet-derived growth factor in an anaplastic thyroid carcinoma cell line.  
Proc. Natl. Acad. Sci. USA 85: 9302- 9306.
- Heldin N.-E. and Westermark B. (1988).  
Epidermal growth factor, but not thyrotropin stimulates the expression of c-fos and c-myc messenger ribonucleic acid in porcine thyroid follicle cells in primary culture.  
Endocrinology 122: 1042
- Heldin N.-E. and Westermark B. (1991).  
The molecular biology of the human anaplastic thyroid carcinoma cell.  
Thyroidology 3: 127- 131.
- Hoeffler J.P., Meyer T.E., Yun Y., Jameson J.L., Habener J.F. (1988).  
Cyclic AMP-responsive DNA-binding protein: structure based on a cloned placental cDNA.  
Science 242: 1430- 1433.
- Hoelting T., Siperstein A.E., Clark O.H., Duh Q.-Y. (1994).  
Epidermal growth factor I messenger ribonucleic acid expression in porcine thyroid follicles is regulated by thyrotropin and iodine.  
Eur. J. Endocrinol. 132: 605- 610.

- Hofbauer L.C., Rafferzeder M., Janssen E.E., Gärtner R. (1995).  
Insulin like growth factor I messenger ribonucleic acid expression in porcine thyroid follicles is regulated by thyrotropin and iodine.  
Eur. J. Endocrinol. 132: 605- 610.
- Hummler E., Cole T.J., Blendy J.A., Ganss R., Aguzzi A., Schmid W., Beermann F., Schütz G. (1994).  
Targeted mutation of the CREB gene: compensation within the CREB/ATF family of transcription factors.  
Proc. Natl. Acad. Sci. USA 91: 5647- 5651.
- Ikuyama S., Niller H.H., Shimura H., Akamizu T., Kohn L.D. (1992).  
Characterization of the 5'-flanking region of the rat thyrotropin receptor gene.  
Mol. Endocrinol. 6: 793- 804.
- Isler M. (1974).  
Loss of mitotic response of the thyroid gland to TSH in hypophysectomized rat and its partial restoration by anterior and posterior pituitary hormones.  
Anat. Rec. 180: 369- 376.
- Jolin T., Morreale de Escobar G., Escobar del Rey F. (1970).  
Differential effects in the rat of thyroidectomy, propylthiouracil and other goitrogens on plasma insulin and thyroid weight.  
Endocrinology 87: 99- 110.
- Jolin T., Tarin M.J., Garcia M.D. (1974).  
Effects of adrenalectomy and corticosterone on thyroid weight of goitrogen-treated rats. Role of adrenal-corticoids in the insulin increase of goitre weight.  
Acta Endocrinol. 75: 734- 747.
- Kanamori A., Abe Y., Yajima Y., Manabe Y., Ho K. (1989).  
Epidermal growth factor receptors in plasma membranes of normal and diseased human thyroid glands.  
J. Clin. Endocrinol. Metab. 68: 899- 903.
- Kikuchi A. and Williams L.T. (1996).  
Regulation of Interaction of *rasp21* with RalGDS and Raf-1 by Cyclic AMP-dependent Protein Kinase.  
J. Biol. Chem. 271: 588- 594.
- Kobayashi K., Shaver J.K., Liang W., Siperstein A.E., Duh Q. Y., Clark O.H. (1993).  
Increased phospholipase C activity in neoplastic thyroid membrane.  
Thyroid 3: 25- 29.
- Kolch W., Heidecker G., Kochs G., Hummel R., Vahidi H., Mischak H., Finkenzeller G., Marme D., Rapp U.R. (1993).  
Protein kinase C alpha activates RAF-1 by direct phosphorylation.  
Nature 364: 249- 252.



Konturek P.C., Dembinski A., Warzecha Z., Ihlm A., Ceranowicz P., Konturek S.J., Stachura J., Hahn E.G. (1998).

Comparison of epidermal growth factor and transforming growth factor-beta 1 expression in hormone-induced acute pancreatitis in rats.

*Digestion* 59: 110- 119.

Korc M., Friess H., Yamanaka Y., Kobrin M.S., Buchler M., Beger H.G. (1994).

Chronic pancreatitis is associated with increased concentrations of epidermal growth factor receptor, transforming growth factor  $\alpha$ , and phospholipase C $\gamma$ .

*Gut* 35: 1465- 1473.

Kraiem Z., Sadeh O., Yosef M., Aharon A. (1995).

Mutual Antagonistic Interactions between the Thyrotropin (Adenosine 3',5'-Monophosphate) and Protein Kinase C/Epidermal Growth Factor (Tyrosine Kinase) Pathways in Cell Proliferation and Differentiation of Cultured Human Thyroid Follicles.

*Endocrinology* 136: 585- 590.

Kyriakis J.M., App H., Zhang X.F., Banerjee P., Brautigan D.L., Rapp K.R., Arruch J. (1992).

Raf-1 activates MAP kinase- kinase.

*Nature* 358: 417- 421.

Lalli E. and Sassone-Corsi P. (1995).

Thyroid-stimulating hormone (TSH)-directed induction of the CREM gene in the thyroid gland participates in the long-term desensitization of the TSH-receptor.

*Biochemistry* 92: 9633- 9637.

Laoide B.M., Foulkes N.S., Schlotter F., Sassone-Corsi P. (1993).

The functional versatility of CREM is determined by its modular structure.

*EMBO J.* 12: 1179- 1191.

Laugwitz K.-L., Allgeier A., Offermanns S., Spicher K., Van Sande J., Dumont J.E., Schultz G. (1995).

The human thyrotropin receptor: a heptahelical receptor capable of stimulating members of all four G protein families.

*Proc. Natl. Acad. Sci. USA* 93: 116- 120.

Lee C.Q., Yun Y., Hoeffler J.P., Habener J.F. (1990).

Cyclic-AMP-responsive transcriptional activation of CREB-327 involves interdependent phosphorylated subdomains.

*EMBO J.* 9: 4455- 4465.

Lemoine N.R., Hughes C.M., Brown C.L., Wynford-Thomas D. (1991).

Abnormalities of the EGF receptor system in human thyroid neoplasia.

*Int. J. Cancer* 49: 558- 561.

- Lemoine N.R., Mayall E.S., Wyllie F.S., Farr C.J., Hughes D., Padua R.A., Thurston V., Williams E.D., Wynford-Thomas D. (1988).  
Activated ras oncogenes in human thyroid cancers.  
*Cancer Res.* 48: 4459- 4463.
- Lemoine N.R., Mayall E.S., Wyllie F.S., Williams E.D., Goyhs M., Stringer B., Wynford-Thomas D. (1989).  
High frequency of ras oncogene activation in all stages of human thyroid tumorigenesis.  
*Oncogene* 4: 159- 164.
- Li G., Hidaka H., Wollheim C.B. (1992)  
Inhibition of Voltage-Gated Ca<sup>2+</sup> Channels and Insulin Secretion in HIT Cells by the Ca<sup>2+</sup>/Calmodulin-Dependent Protein Kinase II Inhibitor KN-62: Comparison with Antagonists of Calmodulin and L-Type Ca<sup>2+</sup> Channels.  
*Mol. Pharmacology* 42: 489- 498.
- Li W., Mischak H., Yu J.-C., Wang L.-M., Mushinski J.F., Heidarman M.A., Pierce J.H. (1994).  
Tyrosine phosphorylation of protein kinase C- $\delta$  in response to its activation.  
*J. Biol. Chem.* 269: 2349-2352.
- Lin R., Beauparlant P., Markis C., Meloche S., Hiscott J. (1996).  
Phosphorylation of I $\kappa$ B $\alpha$  in the C-terminal PEST domain by casein kinase II affects intrinsic protein stability.  
*Mol. Cell. Biol.* 16: 1401- 1409.
- Logan A., Black E.G., Gonzalez A.M., Buscaglia M., Sheppard M.C. (1992).  
Basic FGF: an autocrine mitogen of rat thyroid follicular cells.  
*Endocrinology* 130: 2363- 2372.
- Maciel R.M.B., Moses A.C., Villone G., Tramontano D., Ingbar S.H. (1988).  
Demonstration of the production and physiological role of insulin-like growth factor II in rat thyroid follicular cells in culture.  
*J. Clin. Invest.* 82: 1546- 1553.
- Mäkinen T., Pekonen F., Franssila K., Lamberg B.A. (1988).  
Receptors for epidermal growth factor and thyrotropin in thyroid carcinoma.  
*Acta Endocrinol.* 117: 45- 50.
- Martiny-Baron G., Kazanietz M.G., Mischak H., Blumberg P.M., Kochs G., Hug H., Marme D., Schächtele C. (1993).  
Selective Inhibition of Protein Kinase C Isozymes by the Indolocarbazole Gö 6976.  
*J. Biol. Chem.* 268: 9194- 9197.
- Matowe W.C., Gupta S., Ginsberg J. (1996).  
Regulation of protein kinase C isoforms in FRTL-5 thyroid cells by TSH and phorbol ester.  
*Thyroid* 6: 53- 58..

- Mei F.C., Qiao J., Tsygankova O.M., Meinkoth J.L., Quilliam L.A., Cheng X. (2002).  
Differential Signaling of Cyclic AMP.  
*J. Biol. Chem.* 277: 11497- 11504.
- Miller M.J., Rioux L., Prendergast G.V., Cannon S., White M.A., Meinkoth J.L. (1998).  
Differential Effects of Protein kinase A on Ras Effector Pathways.  
*Mol. Cell. Biol.* 18: 3718- 3726.
- Minuto F., Barreca A., Del Monte P., Cariola G., Torre G.L., Giordano G. (1989).  
Immunoreactive insulin-like growth factor I (IGF-I) and IGF-I-binding protein  
content in human thyroid tissue.  
*J. Clin. Endocrinol. Metab.* 68: 621- 626.
- Miyamoto M., Sugawa H., Kuma K., Mori T., Imura H. (1989).  
Analysis of epidermal growth factor (EGF) receptor and effect of EGF on the  
growth of cultured Graves' and non-neoplastic human thyroid cells.  
*J. Endocrinol. Invest.* 12: 111- 117.
- Montminy M.R. and Bilezikjian L.M. (1987).  
Binding of a nuclear protein to the cyclic-AMP-responsive response element of the  
somatostatin gene.  
*Nature* 328: 175- 178.
- Nakajima T., Uchida C., Anderson S.F., Parvin J.D., Montminy M. (1997).  
Analysis of a cAMP-response activator reveals a two component mechanism for  
transcriptional induction via signal dependent factors.  
*Genes Dev.* 11: 738- 747.
- Namba H., Rubin S.A., Fagin J.A. (1990).  
Point mutations of ras oncogenes are an early event in thyroid tumorigenesis.  
*Mol. Endocrinol.* 4: 1474- 1479.
- Ness G. O., Haugen D. R. F., Varhaug J. E., Akslen L. A., Lillehaug J. R., (1996).  
Cytoplasmic localization of EGF receptor in papillary thyroid carcinomas:  
association with the 150-kDa receptor form.  
*Int. J. Cancer* 65: 161- 167.
- Ollis C.A., Hill D.J., Munro D.S. (1989).  
A role for insulin-like growth factor-I in the regulation of human thyroid cell  
growth by thyrotropin.  
*J. Clin. Endocrinol. Metab.* 123: 495- 500.
- Pandini G., Frasca F., Mineo R., Sciacca L., Vigneri R., Belfiore A. (2002).  
Insulin/insulin-like growth factor I hybrid receptors have different biological  
characteristics depending on the insulin receptor isoform involved.  
*J. Biol. Chem.* 277: 39684- 39695.
- Parma J., Duprez L., Van Sande J. (1993).  
Somatic mutations in the thyrotropin receptor.  
*Science* 246: 1620- 1622.

- Patel V.A., Hill D.J., Eggo M.C., Sheppard M.C., Becks G.P., Logan A. (1996).  
Changes in the immunohistochemical localisation of FGF-2, TGF $\beta$ 1 and thrombospondin-1 are associated with the early angiogenic events in the hyperplastic rat thyroid.  
*J. Endocrinol.* 148: 183- 199.
- Peter M.J., Gerber M., Studer H., Smeds S. (1985).  
Pathogenesis of heterogeneity in human multinodular goiter.  
*J. Clin. Invest.* 76: 1992- 2002.
- Pfannenstiel P., Hotze L.-A., Saller B. (1997).  
Schilddrüsenerkrankungen-Diagnose und Therapie.  
BMV Berliner Medizinische Verlagsanstalt Berlin
- Radice P., Sozzi G., Miozzo M., De Benedetti V., Cariani T., Bongarzone I., Spurr N. K., Pierotti M. A., Della Porta G. (1991).  
The human tropomyosin gene involved in the generation of the TRK oncogene maps to chromosome 1q31.  
*Oncogene* 6: 2145- 2148.
- Reardon D.B., Contessa J.N., Mikkelsen R.B., Valerie K., Amir C., Dent P., Schmidt-Ullrich R.K. (1999).  
Dominant negative EGFR- CD533 and inhibition of MAPK modify JNK1 activation and enhance radiation toxicity of human mammary carcinoma cells.  
*Oncogene* 18: 4756- 4766.
- Rechsteiner M. and Rogers S.W. (1996).  
PEST sequences and regulation by proteolysis.  
*TIBS* 21: 267-171.
- Reiners C., Wegscheider K., Schicha H., Theissen P., Vaupel R., Wrbitzky R., Schumm-Draeger P.-M. (2004).  
Prevalence of thyroid disorders in the working population of Germany: ultrasonography screening in 96,278 unselected employees.  
*Thyroid* 14: 926-932.
- Roger P.P., Hotimsky A., Moreau C., Dumont J.E. (1982).  
Stimulation by thyrotropin, cholera toxin and dibutyryl cyclic AMP of the multiplication of differentiated thyroid cells in vitro.  
*Mol. Cell. Endocrinol.* 26: 165- 176.
- Roger P.P. and Dumont J.E. (1982).  
Epidermal growth factor controls the proliferation and the expression of differentiation in canine thyroid cells in primary culture.  
*FEBS Lett.* 144: 209- 212.

- Roger P.P., Servais P., Dumont J.E. (1987).  
Regulation of dog thyroid epithelial cell cycle by forskolin, an adenylate cyclase activator.  
Exp. Cell Res. 172: 282- 292.
- Roger P.P., Reuse S., Servais P., Van Heuverswyn B., Dumont J.E. (1986).  
Stimulation of cell proliferation and inhibition of differentiation expression by tumor-promoting phorbol esters in dog thyroid cells in primary culture.  
Cancer Res. 46: 898- 906.
- Roger P.P., Taton M., Van Sande J., Dumont J.E. (1988).  
Mitogenic effects of thyrotropin and adenosine 3',5'-monophosphate in differentiated normal human thyroid cells in vitro.  
J. Clin. Endocrinol. Metab. 66: 1158- 1165.
- Saji M., Akamizu T., Sanchez M., Obici S., Avvedimento E., Gottesman M.E., Kohn L.D. (1992).  
Regulation of thyrotropin receptor gene expression in rat FRTL-5 thyroid cells.  
Endocrinology 130: 520- 533.
- Saller B., Stapfer G., Bein B., Hoermann R., Spelsberg F., Mann K. (1993).  
Increased binding capacity of receptors for the epidermal growth factor in benign thyroid nodules and thyroid malignancies.  
Clin. Investig. 71: 898- 902.
- Santisteban P., Acebron A., Polycarpou-Schwarz M., Di Lauro R. (1992).  
Insulin and insulin-like growth factor I regulate a thyroid-specific nuclear protein that binds to the thyroglobulin promoter.  
Mol. Endocrinol. 6: 1310- 1317.
- Santisteban P., Kohn L.D., Di Lauro R. (1987).  
Thyroglobulin gene expression is regulated by insulin and insulin-like growth factor I, as well as thyrotropin, in FRTL5 thyroid cells.  
J. Biol. Chem. 262: 4048-
- Schmidt-Ullrich R.K., Contessa J.N., Dent P., Mikkelsen R.B., Valerie K., Reardon D.B., Bowers G., Lin P.S. (1999).  
Molecular mechanisms of radiation- induced accelerated Upopulation.  
Radiat. Oncol. Investig. 7: 321- 330.
- Shi Y.F., Zou M.J., Schmidt H., Juhasz F.m Stensky V., Robb D., Farid N.R. (1991).  
High rates of ras codon 61 mutation in thyroid tumors in an iodide-deficient area.  
Cancer Res. 51: 2690- 2693.
- Shimura Y., Shimura H., Ohmori M., Ikuyama S., Kohn L.D. (1994).  
Identification of a novel insulin-responsive element in the rat thyrotropin receptor promoter.  
J. Biol. Chem. 269: 31908- 31914.

- Shaver J. K., Tezelman S., Siperstein A. E., Duh Q. Y., Clark O. H. (1993).  
Thyroid-stimulating hormone activates phospholipase C in normal and neoplastic thyroid tissue.  
Surgery 114: 1064- 1069.
- Studer H. and Derwahl M. (1995).  
Mechanisms of Nonneoplastic Endocrine Hyperplasia- A Changing Concept:  
A Review Focused on the Thyroid Gland.  
Endocr. Reviews 16: 411- 426.
- Studer H., Gerber H., Zbaeren J., Peter H.J. (1992).  
Histomorphological and immunohistochemical evidence that human nodular goiters grow by episodic replication of multiple clusters of thyroid follicular cells.  
J. Clin. Endocrinol. Metab.: 75: 1151- 1158.
- Studer H., Peter H. J., Gerber H. (1989).  
Natural heterogeneity of thyroid cells: The basis for understanding thyroid function and nodular goiter growth.  
Endocr. Rev. 10: 125- 135.
- Suarez Nunez J.M. (1970).  
Effect of removal of submaxillary glands on the thyroid gland.  
J. Dent. Res. 49: 454.
- Takasu N., Charrier B., Mauchamp J., Lissitzky S. (1977).  
Positive and negative regulation by thyrotropin of thyroid cyclic AMP response to thyrotropin in porcine thyroid cells.  
FEBS Lett. 84: 191- 194.
- Taylor S.S., Buechler J. A., Yonemoto W. (1990).  
cAMP-dependent protein kinase: framework for a diverse family of regulatory enzymes.  
Ann. Rev. Biochem. 59: 971- 1005.
- Thomas M.J., Umayahara Y., Shu H., Centrella M., Rotwein P., McCarthy T.L. (1996).  
Identification of the cAMP response element that controls transcriptional activation of the insulin-like growth factor-I gene by prostaglandin E<sub>2</sub> in osteoblasts.  
J. Biol. Chem. 271: 21835- 21841.
- Thompson S.D., Franklyn J.A., Watkinson J.C., Verhaeg J.M., Sheppard M.C., Eggo M.C. (1998).  
Fibroblast growth factors 1 and 2 and fibroblast growth factor receptor 1 are elevated in thyroid hyperplasia.  
J. Clin. Endocrinol. Metab.: 83: 1336- 1341.
- Thorburn G.D., Waters M.J., Young I.R., Buntine D., Hopkins P.S. (1981).  
Epidermal growth factor: a critical factor in fetal maturation.  
In: The fetus and independent life, pp 172- 198, Pitman, London.

- Tode B., Serio M., Rotella C.M., Galli G., Franceschelli F., Tanini A., Toccafondi R. (1989).  
 Insulin-like growth factor-I: autocrine secretion by human thyroid follicular cells in primary culture.  
*J. Clin. Endocrinol. Metab.* 69: 91- 96.
- Tomaszewska R., Dembenski A., Warzecha Z., Ceranowicz P., Konturek S.J., Stachura J. (2002).  
 The influence of epidermal growth factor on the course of ischemia-reperfusion induced pancreatitis in rats.  
*J. Physiol. Pharmacol.* 53: 183- 198.
- Tseng Y.C., Burman K.D., Schaudies R.P., Ahmann A.J., D'Avis J., Geelhoed G.W., Wartofsky L. (1989).  
 Effects of epidermal growth factor on thyroglobulin and adenosine 3',5'-monophosphate production by cultured human thyrocytes.  
*J. Clin. Endocrinol. Metab.* 69: 771- 775.
- Ueda Y., Hirai S., Osada S., Suzuki A., Mizuno K., Ohno S. (1996)  
 Protein kinase C $\delta$  activates the MEK-ERK pathway in a manner independent of Ras and dependent on Raf.  
*J. Biol. Chem.* 271: 23512- 23519.
- Uyttensprot N., Costagliola S., Dumont J.E., Miot F. (1999).  
 Requirement for cAMP-responsive element (CRE) binding protein/CRE modulator transcription factors in thyrotropin-induced proliferation of dog thyroid cells in primary culture.  
*Eur. J. Biochem.* 259: 370- 378.
- Uyttensprot N. and Miot F. (1997).  
 Dog CREM transcription factors: cloning, tissue distribution, and identification of new isoforms.  
*Biochem. Biophys. Res. Commun.* 237: 74- 78.
- Van Aelst L., Barr M., Marcus S., Polverino A., Wigler M. (1993).  
 Complex formation between RAS and RAF and other protein kinases.  
*PNAS* 90: 6213- 6217.
- Vannelli G.B., Barni T., Modigliani U., Paulin I., Serio M., Maggi M., Fiorelli G., Balboni G.C. (1990).  
 Insulin-like growth factor receptor I in nonfunctioning thyroid nodules.  
*J. Clin. Endocrinol. Metab.* 71: 1175- 1182.
- Vella V., Sciacca L., Pandini G., Mineo R., Sanatrito S., Vigneri R., Belfiore A. ( )  
 The IGF system in thyroid cancer: new concepts.  
*Mol. Path.* 54: 121- 124.

- Vinson C.R., Sigler P.B., McKnight S.L. (1989).  
Scissor-grip model for DNA recognition by a family of leucine zipper proteins.  
*Science* 246: 911- 916.
- Völzke H., Alte D., Kohlmann T., Lüdemann J., Nauck M., John U., Meng W. (2005).  
Reference intervals of serum thyroid function tests in a previously iodine-deficient area.  
*Thyroid* 15: 279-285.
- Wang X.D., Kiang J.G., Smallridge R.C. (1995).  
Identification of protein kinase C and its multiple isoforms in FRTL-5 thyroid cells.  
*Thyroid* 5: 137- 40.
- Warne P.H., Viciane P.R., Downward J. (1993).  
Direct interaction of Ras and the amino-terminal region of Raf-1 in vitro.  
*Nature* 364: 352- 355.
- Waters M.J., Tweedale R.C., Whip T.A., Shaw G., Manley S.W., Bourke J.R. (1987).  
Dedifferentiation of cultured thyroid cells by epidermal growth factor: some insights into the mechanisms.  
*Mol. Cell. Endocrinol.* 49: 109- 117.
- Westermarck B., Karlsson F.A., Walindor O. (1979).  
Thyrotropin is not a growth factor for human thyroid cells in culture.  
*Proc. Natl. Acad. Sci. USA* 76: 2022- 2026.
- Westermarck K., Karlsson F.A., Westermarck B. (1985).  
Thyrotropin modulates EGF receptor function in porcine thyroid follicle cells.  
*Mol. Cell. Endocrinol.* 40: 17- 23.
- Westermarck K., Lundquist M., Wallin G., Dahlmann T., Hacker G.W., Heldin N.-E., Grimelins L. (1996).  
EGF- receptors in human normal and pathological thyroid tissue.  
*Histopathology* 28: 221- 227.
- Westermarck K., Westermarck B., Karlsson F.A., Ericson L.E. (1986).  
Location of epidermal growth factor receptors on porcine thyroid follicle cells and receptor regulation by thyrotropin.  
*Endocrinology* 118: 1040- 1046.
- Williams D.W., Wynford-Thomas D., Williams E.D. (1987).  
Control of human thyroid follicular cell proliferation in suspension and monolayer culture.  
*Mol. Cell. Endocrinol.* 61: 139- 145.
- Williams D.W., Williams E.D., Wynford-Thomas D. (1989).  
Evidence for autocrine production of IGF-1 in human thyroid adenomas.  
*Mol. Cell. Endocrinology* 61: 139- 143.



- Wollman S.H. and Breitman T.R. (1970)  
Changes in DANN and weight of thyroid glands during hyperplasia and involution.  
*Endocrinology* 86: 322- 327.
- Woloshin P.I., Walton K.M., Rehfuss R.P., Goodman R.H., Cone R.D. (1992).  
3',5'-cyclic adenosine monophosphate-regulated enhancer binding (CREB) activity  
is required for normal growth and differentiated phenotype in the FRTL5 thyroid  
follicular cell line.  
*Mol. Endocrinol.* 6: 1725- 1733.
- Wu J., Dent P., Jelinek T., Wolfman A., Weber M., Sturgill T. (1993).  
Inhibition of the EGF-activating MAP kinase signalling pathway by adenosine-  
3',5'- monophosphate.  
*Science* 262: 1065- 1069.
- Wynford-Thomas D., Smith P., Williams E.D. (1987).  
Proliferative response to cyclic AMP elevation of thyroid epithelium in suspension  
culture.  
*Mol. Cell. Endocrinol.* 51: 163- 166.
- Wynford-Thomas D., Stringer B.M.J., Williams E.D. (1982).  
Circadian rhythm of mitotic activity in normal and hyperplastic rat thyroid.  
*Acta Endocrinol.* 100: 46- 50.
- Xing J., Ginty D.D., Greenberg M.E. (1996).  
Coupling of the RAS-MAPK pathway to gene activation by RSK2, a growth factor-  
regulated CREB kinase.  
*Science* 273: 959- 963.
- Xing J., Kornhauser J.M., Xia Z., Thiele E.A., Greenberg M.E. (1998).  
Nerve growth factor activates extracellular signal-regulated kinase and p38  
mitogen-activated protein kinase pathways to stimulate CREB serine 133  
phosphorylation.  
*Mol. Cell. Biol.* 18: 1946- 1955.
- Yamamoto K.K., Gonzalez G.A., Briggs W.H. III, Montminy M.R. (1988).  
Phosphorylation-induced binding and transcriptional efficacy of nuclear factor  
CREB.  
*Nature* 334: 494- 498.
- Yehia G., Schlotter F., Razavi R., Alessandrini A., Molina C.A. (2001).  
Mitogen-activated protein kinase phosphorylates and targets inducible cAMP early  
repressor to ubiquitin-mediated destruction.  
*J. Biol. Chem.* 276: 35272- 35279.
- Yoshizumi M., Hsieh C.-M., Zhou F., Tsai J.-C., Patterson C., Perrella M.A.,  
Lee M.-E. (1995).  
The ATF site mediates downregulation of the cyclin A gene during contact  
inhibition in vascular endothelial cell.  
*Mol. Cell. Biol.* 15: 3266- 3272.

Zielke A., Tezelman S., Jossart G. H., Wong M., Siperstein A. E., Duh Q.-Y., Clark O. H. (1998).

Establishment of a differentiated thyroid cancer cell line of Hürthle cell origin. *Thyroid* 8: 475- 483.

## Erklärung

„Ich, Astrid C. S. Sobke, erkläre, daß ich die vorgelegte Dissertationsschrift mit dem Thema : „Growth and differentiation control in the thyroid: all ways leading to CREB “ selbst verfasst und keine anderen als die angegebenen Quellen und Hilfsmittel benutzt, ohne die (unzulässige) Hilfe Dritter und auch in Teilen keine Kopien anderer Arbeiten dargestellt habe.“

Datum

Unterschrift

Mein Lebenslauf wird aus Datenschutzgründen in der elektronischen Version meiner Arbeit nicht mit veröffentlicht.