

## 8 Literaturverzeichnis

- Ahrens, P. B., Solursh, M., & Reiter, R. S. (1977). Stage-related capacity for limb chondrogenesis in cell culture. *Dev Biol*, 60(1), 69-82.
- Albrecht, A. N., Schwabe, G. C., Stricker, S., Boddrich, A., Wanker, E. E., & Mundlos, S. (2002). The synpolydactyly homolog (spdh) mutation in the mouse -- a defect in patterning and growth of limb cartilage elements. *Mech Dev*, 112(1-2), 53-67.
- Archer, C. W., Dowthwaite, G. P., & Francis-West, P. (2003). Development of synovial joints. *Birth Defects Res C Embryo Today*, 69(2), 144-155.
- Ashique, A. M., Fu, K., & Richman, J. M. (2002). Signalling via type IA and type IB bone morphogenetic protein receptors (BMPR) regulates intramembranous bone formation, chondrogenesis and feather formation in the chicken embryo. *Int J Dev Biol*, 46(2), 243-253.
- Baur, S. T., Mai, J. J., & Dymecki, S. M. (2000). Combinatorial signaling through BMP receptor IB and GDF5: shaping of the distal mouse limb and the genetics of distal limb diversity. *Development*, 127(3), 605-619.
- Bell, J. (1951). In *Treasury of Human Inheritance* (Vol. 5, pp. 1-31.). London: Cambridge University Press.
- Botting, J. (2002). The History of Thalidomide. *Drug News Perspect*, 15(9), 604-611.
- Buxton, P., Edwards, C., Archer, C. W., & Francis-West, P. (2001). Growth/differentiation factor-5 (GDF-5) and skeletal development. *J Bone Joint Surg Am*, 83-A Suppl 1(Pt 1), S23-30.
- Chang, C., & Hemmati-Brivanlou, A. (1999). Xenopus GDF6, a new antagonist of noggin and a partner of BMPs. *Development*, 126(15), 3347-3357.
- Chang, S. C., Hoang, B., Thomas, J. T., Vukicevic, S., Luyten, F. P., Ryba, N. J., Kozak, C. A., Reddi, A. H., & Moos, M., Jr. (1994). Cartilage-derived morphogenetic proteins. New members of the transforming growth factor-beta superfamily predominantly expressed in long bones during human embryonic development. *J Biol Chem*, 269(45), 28227-28234.
- Chen, Y., & Zhao, X. (1998). Shaping limbs by apoptosis. *J Exp Zool*, 282(6), 691-702.
- Cushing, H. (1916). Hereditary ankylosis of proximal phalangeal joints (symphalangism). *Genetics*, 1:, 90-106.

- DeLise, A. M., Stringa, E., Woodward, W. A., Mello, M. A., & Tuan, R. S. (2000). Embryonic limb mesenchyme micromass culture as an in vitro model for chondrogenesis and cartilage maturation. *Methods Mol Biol*, 137, 359-375.
- Demirhan, O., Türkmen, S., Schwabe, G., Soyupak, E., Akgün, D., Tastemir, D., Karahan, D., Mundlos, S., & Lehmann, K. (2004). A homozygous BMPR1B Mutation Causes a New Subtype of Acromesomelic Chondrodysplasia with Genital Anomalies. *J Med Genet*, 42, 314-317.
- Dore, J. J., Jr., Yao, D., Edens, M., Garamszegi, N., Sholl, E. L., & Leof, E. B. (2001). Mechanisms of transforming growth factor-beta receptor endocytosis and intracellular sorting differ between fibroblasts and epithelial cells. *Mol Biol Cell*, 12(3), 675-684.
- Everman, D. B., Bartels, C. F., Yang, Y., Yanamandra, N., Goodman, F. R., Mendoza-Londono, J. R., Savarirayan, R., White, S. M., Graham, J. M., Jr., Gale, R. P., Svarch, E., Newman, W. G., Kleckers, A. R., Francomano, C. A., Govindaiah, V., Singh, L., Morrison, S., Thomas, J. T., & Warman, M. L. (2002). The mutational spectrum of brachydactyly type C. *Am J Med Genet*, 112(3), 291-296.
- Faiyaz-Ul-Haque, M., Ahmad, W., Zaidi, S. H., Haque, S., Teebi, A. S., Ahmad, M., Cohn, D. H., & Tsui, L. C. (2002). Mutation in the cartilage-derived morphogenetic protein-1 (CDMP1) gene in a kindred affected with fibular hypoplasia and complex brachydactyly (DuPan syndrome). *Clin Genet*, 61(6), 454-458.
- Francis-West, P. H., Abdelfattah, A., Chen, P., Allen, C., Parish, J., Ladher, R., Allen, S., MacPherson, S., Luyten, F. P., & Archer, C. W. (1999). Mechanisms of GDF-5 action during skeletal development. *Development*, 126(6), 1305-1315.
- Freire-Maia, N., Maia, N. A., & Pacheco, C. N. (1980). Mohr-Wriedt (A2) brachydactyly: analysis of a large Brazilian kindred. *Hum Hered*, 30(4), 225-231.
- Friedlaender, G. E., Perry, C. R., Cole, J. D., Cook, S. D., Cierny, G., Muschler, G. F., Zych, G. A., Calhoun, J. H., LaForte, A. J., & Yin, S. (2001). Osteogenic protein-1 (bone morphogenetic protein-7) in the treatment of tibial nonunions. *J Bone Joint Surg Am*, 83-A Suppl 1(Pt 2), S151-158.

- Fuhrmann, W., & Steffens, C. (1966). [Dominant hereditary bilateral dysplasia and synostosis of the elbow joint, with symmetrical brachymesophalangy and brachymetacarpal as well as synostoses in the finger, carpal and tarsal region]. *Humangenetik*, 3(1), 64-75.
- Gao, B., & He, L. (2004). Answering a century old riddle: brachydactyly type A1. *Cell Res*, 14(3), 179-187.
- Garamszegi, N., Dore, J. J., Jr., Penheiter, S. G., Edens, M., Yao, D., & Leof, E. B. (2001). Transforming growth factor beta receptor signaling and endocytosis are linked through a COOH terminal activation motif in the type I receptor. *Mol Biol Cell*, 12(9), 2881-2893.
- Gilbert, S. F. (2003). *Developmental Biology* (seventh ed.).
- Gilbert, S. F. (2003). Opening Darwin's black box: teaching evolution through developmental genetics. *Nat Rev Genet*, 4(9), 735-741.
- Gilboa, L., Nohe, A., Geissendorfer, T., Sebald, W., Henis, Y. I., & Knaus, P. (2000). Bone morphogenetic protein receptor complexes on the surface of live cells: a new oligomerization mode for serine/threonine kinase receptors. *Mol Biol Cell*, 11(3), 1023-1035.
- Gong, Y., Krakow, D., Marcelino, J., Wilkin, D., Chitayat, D., Babul-Hirji, R., Hudgins, L., Cremers, C. W., Cremers, F. P., Brunner, H. G., Reinker, K., Rimoin, D. L., Cohn, D. H., Goodman, F. R., Reardon, W., Patton, M., Francomano, C. A., & Warman, M. L. (1999). Heterozygous mutations in the gene encoding noggin affect human joint morphogenesis. *Nat Genet*, 21(3), 302-304.
- Goodstadt, L., & Ponting, C. P. (2001). CHROMA: consensus-based colouring of multiple alignments for publication. *Bioinformatics*, 17(9), 845-846.
- Groppe, J., Greenwald, J., Wiater, E., Rodriguez-Leon, J., Economides, A. N., Kwiatkowski, W., Affolter, M., Vale, W. W., Belmonte, J. C., & Choe, S. (2002). Structural basis of BMP signalling inhibition by the cystine knot protein Noggin. *Nature*, 420(6916), 636-642.
- Guex, N., & Peitsch, M. C. (1997). SWISS-MODEL and the Swiss-PdbViewer: an environment for comparative protein modeling. *Electrophoresis*, 18(15), 2714-2723.

- Hartmann, C., & Tabin, C. J. (2001). Wnt-14 plays a pivotal role in inducing synovial joint formation in the developing appendicular skeleton. *Cell*, 104(3), 341-351.
- Hassel, S., Schmitt, S., Hartung, A., Roth, M., Nohe, A., Petersen, N., Ehrlich, M., Henis, Y. I., Sebald, W., & Knaus, P. (2003). Initiation of Smad-dependent and Smad-independent signaling via distinct BMP-receptor complexes. *J Bone Joint Surg Am*, 85-A Suppl 3, 44-51.
- Hinck, A. P., Archer, S. J., Qian, S. W., Roberts, A. B., Sporn, M. B., Weatherbee, J. A., Tsang, M. L., Lucas, R., Zhang, B. L., Wenker, J., & Torchia, D. A. (1996). Transforming growth factor beta 1: three-dimensional structure in solution and comparison with the X-ray structure of transforming growth factor beta 2. *Biochemistry*, 35(26), 8517-8534.
- Hotten, G., Neidhardt, H., Jacobowsky, B., & Pohl, J. (1994). Cloning and expression of recombinant human growth/differentiation factor 5. *Biochem Biophys Res Commun*, 204(2), 646-652.
- Hughes, S. H., Greenhouse, J. J., Petropoulos, C. J., & Sutrave, P. (1987). Adaptor plasmids simplify the insertion of foreign DNA into helper-independent retroviral vectors. *J Virol*, 61(10), 3004-3012.
- Hurle, J. M., & Colvee, E. (1982). Surface changes in the embryonic interdigital epithelium during the formation of the free digits: a comparative study in the chick and duck foot. *J Embryol Exp Morphol*, 69, 251-263.
- Huse, M., Muir, T. W., Xu, L., Chen, Y. G., Kuriyan, J., & Massague, J. (2001). The TGF beta receptor activation process: an inhibitor- to substrate-binding switch. *Mol Cell*, 8(3), 671-682.
- Johnson, D., Kan, S. H., Oldridge, M., Trembath, R. C., Roche, P., Esnouf, R. M., Giele, H., & Wilkie, A. O. (2003). Missense mutations in the homeodomain of HOXD13 are associated with brachydactyly types D and E. *Am J Hum Genet*, 72(4), 984-997.
- Johnson, R. L., & Tabin, C. J. (1997). Molecular models for vertebrate limb development. *Cell*, 90(6), 979-990.
- Keller, S., Nickel, J., Zhang, J. L., Sebald, W., & Mueller, T. D. (2004). Molecular recognition of BMP-2 and BMP receptor IA. *Nat Struct Mol Biol*, 11(5), 481-488.

- Kirsch, T., Nickel, J., & Sebald, W. (2000). BMP-2 antagonists emerge from alterations in the low-affinity binding epitope for receptor BMPR-II. *Embo J*, 19(13), 3314-3324.
- Kirsch, T., Sebald, W., & Dreyer, M. K. (2000). Crystal structure of the BMP-2-BRIA ectodomain complex. *Nat Struct Biol*, 7(6), 492-496.
- Kjaer, K. W., Eiberg, H., Hansen, L., van der Hagen, C. B., Rosendahl, K., Tommerup, N., & Mundlos, S. (2005). A mutation in the receptor binding site of GDF5 causes Mohr-Wriedt brachydactyly type A2. *J Med Genet*.
- Kornak, U., & Mundlos, S. (2003). Genetic disorders of the skeleton: a developmental approach. *Am J Hum Genet*, 73(3), 447-474.
- Krakow, D., Reinker, K., Powell, B., Cantor, R., Priore, M. A., Garber, A., Lachman, R. S., Rimoin, D. L., & Cohn, D. H. (1998). Localization of a multiple synostoses-syndrome disease gene to chromosome 17q21-22. *Am J Hum Genet*, 63(1), 120-124.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227(5259), 680-685.
- Lanske, B., Karaplis, A. C., Lee, K., Luz, A., Vortkamp, A., Pirro, A., Karperien, M., Defize, L. H., Ho, C., Mulligan, R. C., Abou-Samra, A. B., Juppner, H., Segre, G. V., & Kronenberg, H. M. (1996). PTH/PTHrP receptor in early development and Indian hedgehog-regulated bone growth. *Science*, 273(5275), 663-666.
- Lehmann, K., Seemann, P., Stricker, S., Sammar, M., Meyer, B., Suring, K., Majewski, F., Tinschert, S., Grzeschik, K. H., Muller, D., Knaus, P., Nurnberg, P., & Mundlos, S. (2003). Mutations in bone morphogenetic protein receptor 1B cause brachydactyly type A2. *Proc Natl Acad Sci U S A*, 100(21), 12277-12282.
- Logan, M. (2003). Finger or toe: the molecular basis of limb identity. *Development*, 130(26), 6401-6410.
- Logan, M., & Tabin, C. (1998). Targeted gene misexpression in chick limb buds using avian replication-competent retroviruses. *Methods*, 14(4), 407-420.
- Lutz, M., & Knaus, P. (2002). Integration of the TGF-beta pathway into the cellular signalling network. *Cell Signal*, 14(12), 977-988.
- Luyten, F. P. (1997). Cartilage-derived morphogenetic protein-1. *Int J Biochem Cell Biol*, 29(11), 1241-1244.

- Massague, J., & Chen, Y. G. (2000). Controlling TGF-beta signaling. *Genes Dev*, 14(6), 627-644.
- Merino, R., Macias, D., Ganan, Y., Economides, A. N., Wang, X., Wu, Q., Stahl, N., Sampath, K. T., Varona, P., & Hurle, J. M. (1999). Expression and function of Gdf-5 during digit skeletogenesis in the embryonic chick leg bud. *Dev Biol*, 206(1), 33-45.
- Minina, E., Wenzel, H. M., Kreschel, C., Karp, S., Gaffield, W., McMahon, A. P., & Vortkamp, A. (2001). BMP and Ihh/PTHrP signaling interact to coordinate chondrocyte proliferation and differentiation. *Development*, 128(22), 4523-4534.
- Mishina, Y., Suzuki, A., Ueno, N., & Behringer, R. R. (1995). Bmpr encodes a type I bone morphogenetic protein receptor that is essential for gastrulation during mouse embryogenesis. *Genes Dev*, 9(24), 3027-3037.
- Mohr, O. L., & Wried, C. (1919). *A new type of hereditary brachphalangy in man.*: Institution of Washington.
- Morgan, B. A., & Fekete, D. M. (1996). Manipulating gene expression with replication-competent retroviruses. *Methods Cell Biol*, 51, 185-218.
- Morgan, E. A., Nguyen, S. B., Scott, V., & Stadler, H. S. (2003). Loss of Bmp7 and Fgf8 signaling in Hoxa13-mutant mice causes hypospadias. *Development*, 130(14), 3095-3109.
- Mujagic, H., Chabner, B. A., & Mujagic, Z. (2002). Mechanisms of action and potential therapeutic uses of thalidomide. *Croat Med J*, 43(3), 274-285.
- Mulsant, P., Lecerf, F., Fabre, S., Schibler, L., Monget, P., Lanneluc, I., Pisset, C., Riquet, J., Monniaux, D., Callebaut, I., Cribiu, E., Thimonier, J., Teyssier, J., Bodin, L., Cognie, Y., Chitour, N., & Elsen, J. M. (2001). Mutation in bone morphogenetic protein receptor-IB is associated with increased ovulation rate in Booroola Merino ewes. *Proc Natl Acad Sci U S A*, 98(9), 5104-5109.
- Nakamura, K., Shirai, T., Morishita, S., Uchida, S., Saeki-Miura, K., & Makishima, F. (1999). p38 mitogen-activated protein kinase functionally contributes to chondrogenesis induced by growth/differentiation factor-5 in ATDC5 cells. *Exp Cell Res*, 250(2), 351-363.

- Naski, M. C., Colvin, J. S., Coffin, J. D., & Ornitz, D. M. (1998). Repression of hedgehog signaling and BMP4 expression in growth plate cartilage by fibroblast growth factor receptor 3. *Development*, 125(24), 4977-4988.
- Nickel, J., Kotzsch, A., Sebald, W., & Mueller, T. D. (2005). A single residue of GDF-5 defines binding specificity to BMP receptor IB. *J Mol Biol*, 349(5), 933-947.
- Nishitoh, H., Ichijo, H., Kimura, M., Matsumoto, T., Makishima, F., Yamaguchi, A., Yamashita, H., Enomoto, S., & Miyazono, K. (1996). Identification of type I and type II serine/threonine kinase receptors for growth/differentiation factor-5. *J Biol Chem*, 271(35), 21345-21352.
- Niswander, L. (2003). Pattern formation: old models out on a limb. *Nat Rev Genet*, 4(2), 133-143.
- Niswander, L., Jeffrey, S., Martin, G. R., & Tickle, C. (1994). A positive feedback loop coordinates growth and patterning in the vertebrate limb. *Nature*, 371(6498), 609-612.
- Nohe, A., Hassel, S., Ehrlich, M., Neubauer, F., Sebald, W., Henis, Y. I., & Knaus, P. (2002). The mode of bone morphogenetic protein (BMP) receptor oligomerization determines different BMP-2 signaling pathways. *J Biol Chem*, 277(7), 5330-5338.
- Nohe, A., Keating, E., Knaus, P., & Petersen, N. O. (2004). Signal transduction of bone morphogenetic protein receptors. *Cell Signal*, 16(3), 291-299.
- Oldridge, M., Fortuna, A. M., Maringa, M., Propping, P., Mansour, S., Pollit, C., DeChiara, T. M., Kimble, R. B., Valenzuela, D. M., Yancopoulos, G. D., & Wilkie, A. O. (2000). Dominant mutations in ROR2, encoding an orphan receptor tyrosine kinase, cause brachydactyly type B. *Nat Genet*, 24(3), 275-278.
- Parr, B. A., & McMahon, A. P. (1995). Dorsalizing signal Wnt-7a required for normal polarity of D-V and A-P axes of mouse limb. *Nature*, 374(6520), 350-353.
- Peitsch, M. C., Wells, T. N., Stampf, D. R., & Sussman, J. L. (1995). The Swiss-3DImage collection and PDB-Browser on the World-Wide Web. *Trends Biochem Sci*, 20(2), 82-84.

- Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C., & Ferrin, T. E. (2004). UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem*, 25(13), 1605-1612.
- Pizette, S., Abate-Shen, C., & Niswander, L. (2001). BMP controls proximodistal outgrowth, via induction of the apical ectodermal ridge, and dorsoventral patterning in the vertebrate limb. *Development*, 128(22), 4463-4474.
- Polinkovsky, A., Robin, N. H., Thomas, J. T., Irons, M., Lynn, A., Goodman, F. R., Reardon, W., Kant, S. G., Brunner, H. G., van der Burgt, I., Chitayat, D., McGaughran, J., Donnai, D., Luyten, F. P., & Warman, M. L. (1997). Mutations in CDMP1 cause autosomal dominant brachydactyly type C. *Nat Genet*, 17(1), 18-19.
- Riddle, R. D., Ensini, M., Nelson, C., Tsuchida, T., Jessell, T. M., & Tabin, C. (1995). Induction of the LIM homeobox gene Lmx1 by WNT7a establishes dorsoventral pattern in the vertebrate limb. *Cell*, 83(4), 631-640.
- Riddle, R. D., Johnson, R. L., Laufer, E., & Tabin, C. (1993). Sonic hedgehog mediates the polarizing activity of the ZPA. *Cell*, 75(7), 1401-1416.
- Rountree, R. B., Schoor, M., Chen, H., Marks, M. E., Harley, V., Mishina, Y., & Kingsley, D. M. (2004). BMP receptor signaling is required for postnatal maintenance of articular cartilage. *PLoS Biol*, 2(11), e355.
- Sambrook, J., Russell, D. W., & Sambrook, J. E. (2001). *Molecular Cloning: A laboratory manual*: Cold Spring Harbor Laboratory Press.
- Sammar, M., Stricker, S., Schwabe, G. C., Sieber, C., Hartung, A., Hanke, M., Oishi, I., Pohl, J., Minami, Y., Sebald, W., Mundlos, S., & Knaus, P. (2004). Modulation of GDF5/BRI-b signalling through interaction with the tyrosine kinase receptor Ror2. *Genes Cells*, 9(12), 1227-1238.
- Sanger, F., Nicklen, S., & Coulson, A. R. (1992). DNA sequencing with chain-terminating inhibitors. 1977. *Biotechnology*, 24, 104-108.
- Schreuder, H., Liesum, A., Pohl, J., Kruse, M., & Koyama, M. (2005). Crystal structure of recombinant human growth and differentiation factor 5: evidence for interaction of the type I and type II receptor-binding sites. *Biochem Biophys Res Commun*, 329(3), 1076-1086.

- Schwabe, G. C., & Mundlos, S. (2004). Genetics of congenital hand anomalies. *Handchir Mikrochir Plast Chir*, 36(2-3), 85-97.
- Schwabe, G. C., Tinschert, S., Buschow, C., Meinecke, P., Wolff, G., Gillessen-Kaesbach, G., Oldridge, M., Wilkie, A. O., Komec, R., & Mundlos, S. (2000). Distinct mutations in the receptor tyrosine kinase gene ROR2 cause brachydactyly type B. *Am J Hum Genet*, 67(4), 822-831.
- Schwabe, G. C., Turkmen, S., Leschik, G., Palanduz, S., Stover, B., Goecke, T. O., & Mundlos, S. (2004). Brachydactyly type C caused by a homozygous missense mutation in the prodomain of CDMP1. *Am J Med Genet*, 124A(4), 356-363.
- Schwede, T., Kopp, J., Guex, N., & Peitsch, M. C. (2003). SWISS-MODEL: An automated protein homology-modeling server. *Nucleic Acids Res*, 31(13), 3381-3385.
- Sebald, W., Nickel, J., Zhang, J. L., & Mueller, T. D. (2004). Molecular recognition in bone morphogenetic protein (BMP)/receptor interaction. *Biol Chem*, 385(8), 697-710.
- Seemann, P., Schwappacher, R., Kjaer, K. W., Krakow, D., Lehmann, K., Dawson, K., Stricker, S., Pohl, J., Ploger, F., Staub, E., Nickel, J., Sebald, W., Knaus, P., & Mundlos, S. (2005). Activating and deactivating mutations in the receptor interaction site of GDF5 cause symphalangism or brachydactyly type A2. *J Clin Invest*, 115(9), 2373-2381.
- Solursh, M., Ahrens, P. B., & Reiter, R. S. (1978). A tissue culture analysis of the steps in limb chondrogenesis. *In Vitro*, 14(1), 51-61.
- Storm, E. E., Huynh, T. V., Copeland, N. G., Jenkins, N. A., Kingsley, D. M., & Lee, S. J. (1994). Limb alterations in brachypodism mice due to mutations in a new member of the TGF beta-superfamily. *Nature*, 368(6472), 639-643.
- Storm, E. E., & Kingsley, D. M. (1996). Joint patterning defects caused by single and double mutations in members of the bone morphogenetic protein (BMP) family. *Development*, 122(12), 3969-3979.
- Storm, E. E., & Kingsley, D. M. (1999). GDF5 coordinates bone and joint formation during digit development. *Dev Biol*, 209(1), 11-27.
- Stricker, S., Fundele, R., Vortkamp, A., & Mundlos, S. (2002). Role of Runx genes in chondrocyte differentiation. *Dev Biol*, 245(1), 95-108.

- Sun, X., Mariani, F. V., & Martin, G. R. (2002). Functions of FGF signalling from the apical ectodermal ridge in limb development. *Nature*, 418(6897), 501-508.
- Szczaluba, K., Hilbert, K., Obersztyn, E., Zabel, B., Mazurczak, T., & Kozlowski, K. (2005). Du Pan syndrome phenotype caused by heterozygous pathogenic mutations in CDMP1 gene. *Am J Med Genet A*, 138(4), 379-383.
- Thomas, J. T., Kilpatrick, M. W., Lin, K., Erlacher, L., Lembessis, P., Costa, T., Tsipouras, P., & Luyten, F. P. (1997). Disruption of human limb morphogenesis by a dominant negative mutation in CDMP1. *Nat Genet*, 17(1), 58-64.
- Thomas, J. T., Lin, K., Nandedkar, M., Camargo, M., Cervenka, J., & Luyten, F. P. (1996). A human chondrodysplasia due to a mutation in a TGF-beta superfamily member. *Nat Genet*, 12(3), 315-317.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F., & Higgins, D. G. (1997). The CLUSTAL\_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res*, 25(24), 4876-4882.
- Tsumaki, N., Tanaka, K., Arikawa-Hirasawa, E., Nakase, T., Kimura, T., Thomas, J. T., Ochi, T., Luyten, F. P., & Yamada, Y. (1999). Role of CDMP-1 in skeletal morphogenesis: promotion of mesenchymal cell recruitment and chondrocyte differentiation. *J Cell Biol*, 144(1), 161-173.
- Urist, M. R. (1965). Bone: formation by autoinduction. *Science*, 150(698), 893-899.
- Vesalius, A. (1551). *Anatomia - Ein kurtzer Auszug der beschreibung aller glider menschlichs Leybs*: Reprint Verlag Leipzig.
- Vortkamp, A., Lee, K., Lanske, B., Segre, G. V., Kronenberg, H. M., & Tabin, C. J. (1996). Regulation of rate of cartilage differentiation by Indian hedgehog and PTH-related protein. *Science*, 273(5275), 613-622.
- Wan, M., & Cao, X. (2005). BMP signaling in skeletal development. *Biochem Biophys Res Commun*, 328(3), 651-657.
- Wieser, R., Wrana, J. L., & Massague, J. (1995). GS domain mutations that constitutively activate T beta R-I, the downstream signaling component in the TGF-beta receptor complex. *Embo J*, 14(10), 2199-2208.

- Wilson, T., Wu, X. Y., Juengel, J. L., Ross, I. K., Lumsden, J. M., Lord, E. A., Dodds, K. G., Walling, G. A., McEwan, J. C., O'Connell, A. R., McNatty, K. P., & Montgomery, G. W. (2001). Highly prolific Booroola sheep have a mutation in the intracellular kinase domain of bone morphogenetic protein IB receptor (ALK-6) that is expressed in both oocytes and granulosa cells. *Biol Reprod*, 64(4), 1225-1235.
- Wozney, J. M., Rosen, V., Celeste, A. J., Mitsock, L. M., Whitters, M. J., Kriz, R. W., Hewick, R. M., & Wang, E. A. (1988). Novel regulators of bone formation: molecular clones and activities. *Science*, 242(4885), 1528-1534.
- Zimmerman, L. B., De Jesus-Escobar, J. M., & Harland, R. M. (1996). The Spemann organizer signal noggin binds and inactivates bone morphogenetic protein 4. *Cell*, 86(4), 599-606.
- Zou, H., Wieser, R., Massague, J., & Niswander, L. (1997). Distinct roles of type I bone morphogenetic protein receptors in the formation and differentiation of cartilage. *Genes Dev*, 11(17), 2191-2203.