

7 REFERENCES

- Abu-Baker, S., Qi, X., Newstadt, J., and Lorian, G. A. (2005). Structural changes in a binary mixed phospholipid bilayer of DOPG and DOPS upon saposin C interaction at acidic pH utilizing 31P and 2H solid-state NMR spectroscopy. *Biochim Biophys Acta* 1717, 58-66.
- Adam-Klages, S., Adam, D., Wiegmann, K., Struve, S., Kolanus, W., Schneider-Mergener, J., and Kronke, M. (1996). FAN, a novel WD-repeat protein, couples the p55 TNF-receptor to neutral sphingomyelinase. *Cell* 86, 937-947.
- Adams, P. D., Grosse-Kunstleve, R. W., Hung, L. W., Ioerger, T. R., McCoy, A. J., Moriarty, N. W., Read, R. J., Sacchettini, J. C., Sauter, N. K., and Terwilliger, T. C. (2002). PHENIX: building new software for automated crystallographic structure determination. *Acta Crystallogr D Biol Crystallogr* 58, 1948-1954.
- Ahn, V. E., Faull, K. F., Whitelegge, J. P., Fluharty, A. L., and Prive, G. G. (2003). Crystal structure of saposin B reveals a dimeric shell for lipid binding. *Proc Natl Acad Sci U S A* 100, 38-43.
- Ahn, V. E., Leyko, P., Alattia, J. R., Chen, L., and Prive, G. G. (2006). Crystal structures of saposins A and C. *Protein Sci* 15, 1849-1857.
- Alattia, J. R., Shaw, J. E., Yip, C. M., and Prive, G. G. (2006). Direct visualization of saposin remodelling of lipid bilayers. *J Mol Biol* 362, 943-953.
- Alattia, J. R., Shaw, J. E., Yip, C. M., and Prive, G. G. (2007). Molecular imaging of membrane interfaces reveals mode of beta-glucosidase activation by saposin C. *Proc Natl Acad Sci U S A* 104, 17394-17399.
- Amidon, B., Brown, A., and Waite, M. (1996). Transacylase and phospholipases in the synthesis of bis(monoacylglycerophosphate). *Biochemistry* 35, 13995-14002.
- Anderson, D. H., Sawaya, M. R., Cascio, D., Ernst, W., Modlin, R., Krensky, A., and Eisenberg, D. (2003). Granulysin crystal structure and a structure-derived lytic mechanism. *J Mol Biol* 325, 355-365.
- Arroyo, M., de la Mata, I., Acebal, C., and Castillon, M. P. (2003). Biotechnological applications of penicillin acylases: state-of-the-art. *Appl Microbiol Biotechnol* 60, 507-514.
- Artymiuk, P. J. (1995). A sting in the (N-terminal) tail. *Nat Struct Biol* 2, 1035-1037.
- Arutyunyan, A. M., Rafikova, E. R., Drachev, V. A., and Dobrov, E. N. (2001). Appearance of "beta-like" circular dichroism spectra on protein aggregation that is not accompanied by transition to beta-structure. *Biochemistry (Mosc)* 66, 1378-1380.
- Aubert-Jousset, E., Sbarra, V., and Lombardo, D. (2004). Site-directed mutagenesis of the distal basic cluster of pancreatic bile salt-dependent lipase. *J Biol Chem* 279, 39697-39704.
- Bailey, S. (1994). The CCP4 Suite - Programs for Protein Crystallography. *Acta Crystallographica Section D-Biological Crystallography* 50, 760-763.
- Barton, G. J. (1993). ALSCRIPT: a tool to format multiple sequence alignments. *Protein Eng* 6, 37-40.
- Batta, A. K., Salen, G., and Shefer, S. (1984). Substrate specificity of cholylglycine hydrolase for the hydrolysis of bile acid conjugates. *J Biol Chem* 259, 15035-15039.
- Bernardo, K., Hurwitz, R., Zenk, T., Desnick, R. J., Ferlinz, K., Schuchman, E. H., and Sandhoff, K. (1995). Purification, characterization, and biosynthesis of human acid ceramidase. *J Biol Chem* 270, 11098-11102.
- Bornhauser, B. C., Olsson, P. A., and Lindholm, D. (2003). MSAP is a novel MIR-interacting protein that enhances neurite outgrowth and increases myosin regulatory light chain. *J Biol Chem* 278, 35412-35420.

- Brady, R. O., Kanfer, J. N., and Shapiro, D. (1965). Metabolism of Glucocerebrosides. II. Evidence of an Enzymatic Deficiency in Gaucher's Disease. *Biochem Biophys Res Commun* 18, 221-225.
- Brannigan, J. A., Dodson, G., Duggleby, H. J., Moody, P. C., Smith, J. L., Tomchick, D. R., and Murzin, A. G. (1995). A protein catalytic framework with an N-terminal nucleophile is capable of self-activation. *Nature* 378, 416-419.
- Bretscher, M. S. (1973). Membrane structure: some general principles. *Science* 181, 622-629.
- Brunger, A. T., Adams, P. D., Clore, G. M., DeLano, W. L., Gros, P., Grosse-Kunstleve, R. W., Jiang, J. S., Kuszewski, J., Nilges, M., Pannu, N. S., et al. (1998). Crystallography & NMR system: A new software suite for macromolecular structure determination. *Acta Crystallographica Section D-Biological Crystallography* 54, 905-921.
- Budisa, N., Steipe, B., Demange, P., Eckerskorn, C., Kellermann, J., and Huber, R. (1995). High-level biosynthetic substitution of methionine in proteins by its analogs 2-aminohexanoic acid, selenomethionine, telluromethionine and ethionine in *Escherichia coli*. *Eur J Biochem* 230, 788-796.
- Cevc, G., and Richardsen, H. (1999). Lipid vesicles and membrane fusion. *Adv Drug Deliv Rev* 38, 207-232.
- Christomanou, H., Aignesberger, A., and Linke, R. P. (1986). Immunochemical characterization of two activator proteins stimulating enzymic sphingomyelin degradation in vitro. Absence of one of them in a human Gaucher disease variant. *Biol Chem Hoppe Seyler* 367, 879-890.
- Ciaffoni, F., Salvioli, R., Tatti, M., Arancia, G., Crateri, P., and Vaccaro, A. M. (2001). Saposin D solubilizes anionic phospholipid-containing membranes. *J Biol Chem* 276, 31583-31589.
- Ciaffoni, F., Tatti, M., Salvioli, R., and Vaccaro, A. M. (2003). Interaction of saposin D with membranes: effect of anionic phospholipids and sphingolipids. *Biochem J* 373, 785-792.
- Coleman, J. P., and Hudson, L. L. (1995). Cloning and characterization of a conjugated bile acid hydrolase gene from *Clostridium perfringens*. *Appl Environ Microbiol* 61, 2514-2520.
- Cremesti, A. E., Goni, F. M., and Kolesnick, R. (2002). Role of sphingomyelinase and ceramide in modulating rafts: do biophysical properties determine biologic outcome? *FEBS Lett* 531, 47-53.
- de Alba, E., Weiler, S., and Tjandra, N. (2003). Solution structure of human saposin C: pH-dependent interaction with phospholipid vesicles. *Biochemistry* 42, 14729-14740.
- De Smet, I., De Boever, P., and Verstraete, W. (1998). Cholesterol lowering in pigs through enhanced bacterial bile salt hydrolase activity. *Br J Nutr* 79, 185-194.
- Desnick, R. J. (2004). Enzyme replacement and enhancement therapies for lysosomal diseases. *J Inherit Metab Dis* 27, 385-410.
- Desnick, R. J., and Schuchman, E. H. (2002). Enzyme replacement and enhancement therapies: lessons from lysosomal disorders. *Nat Rev Genet* 3, 954-966.
- Dietschy, J. M. (1974). Bile acids: their absorption from the gastrointestinal tract and role during fat absorption. *Verh Dtsch Ges Inn Med* 80, 399-407.
- Duttleby, H. J., Tolley, S. P., Hill, C. P., Dodson, E. J., Dodson, G., and Moody, P. C. (1995). Penicillin acylase has a single-amino-acid catalytic centre. *Nature* 373, 264-268.
- Egas, C., Lavoura, N., Resende, R., Brito, R. M., Pires, E., de Lima, M. C., and Faro, C. (2000). The saposin-like domain of the plant aspartic proteinase precursor is a potent inducer of vesicle leakage. *J Biol Chem* 275, 38190-38196.
- Elias, P. M. (1983). Epidermal lipids, barrier function, and desquamation. *J Invest Dermatol Suppl*, 44s-49s.
- Emsley, P., and Cowtan, K. (2004). Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr* 60, 2126-2132.

- Eyssen, H. (1973). Role of the gut microflora in metabolism of lipids and sterols. *Proc Nutr Soc* 32, 59-63.
- Farber, S., Cohen, J., and Uzman, L. L. (1957). Lipogranulomatosis; a new lipo-glycoprotein storage disease. *J Mt Sinai Hosp N Y* 24, 816-837.
- Fehrenbacher, N., and Jaattela, M. (2005). Lysosomes as targets for cancer therapy. *Cancer Res* 65, 2993-2995.
- Ferlinz, K., Kopal, G., Bernardo, K., Linke, T., Bar, J., Breiden, B., Neumann, U., Lang, F., Schuchman, E. H., and Sandhoff, K. (2001). Human acid ceramidase: processing, glycosylation, and lysosomal targeting. *J Biol Chem* 276, 35352-35360.
- Finnegan, C. M., Rawat, S. S., Puri, A., Wang, J. M., Ruscetti, F. W., and Blumenthal, R. (2004). Ceramide, a target for antiretroviral therapy. *Proc Natl Acad Sci U S A* 101, 15452-15457.
- Fischer, G., Demuth, H. U., and Barth, A. (1983). N,O-diacylhydroxylamines as enzyme-activated inhibitors for serine proteases. *Pharmazie* 38, 249-250.
- Flores, T. P., Moss, D. S., and Thornton, J. M. (1994). An algorithm for automatically generating protein topology cartoons. *Protein Eng* 7, 31-37.
- Furst, T., Bott, C., Stein, J., and Dressman, J. B. (2005). Enteric-coated cholylsarcosine microgranules for the treatment of short bowel syndrome. *J Pharm Pharmacol* 57, 53-60.
- Fürst, W., and Sandhoff, K. (1992). Activator proteins and topology of lysosomal sphingolipid catabolism. *Biochim Biophys Acta* 1126, 1-16.
- Fürst W., W. M., and K. Sandhoff (1988). The precursor of sulfatide activator protein is processed to three different proteins. *Biol Chem Hoppe Seyler* 369, 317-328.
- Gatt, S. (1963). Enzymic Hydrolysis and Synthesis of Ceramides. *J Biol Chem* 238, 3131-3133.
- Gaurav, K., Kundu, K., and Kundu, S. (2007). Microbial production of 7-aminocephalosporanic acid and new generation cephalosporins (cephalothin) by different processing strategies. *Artif Cells Blood Substit Immobil Biotechnol* 35, 345-358.
- Gonzalez, C., Langdon, G. M., Bruix, M., Galvez, A., Valdivia, E., Maqueda, M., and Rico, M. (2000). Bacteriocin AS-48, a microbial cyclic polypeptide structurally and functionally related to mammalian NK-lysin. *Proc Natl Acad Sci U S A* 97, 11221-11226.
- Goodsell, D. S., Morris, G. M., and Olson, A. J. (1996). Automated docking of flexible ligands: applications of AutoDock. *J Mol Recognit* 9, 1-5.
- Gopal-Srivastava, R., and Hylemon, P. B. (1988). Purification and characterization of bile salt hydrolase from *Clostridium perfringens*. *J Lipid Res* 29, 1079-1085.
- Groll, M., Ditzel, L., Lowe, J., Stock, D., Bochtler, M., Bartunik, H. D., and Huber, R. (1997). Structure of 20S proteasome from yeast at 2.4 Å resolution. *Nature* 386, 463-471.
- Gutsmann, T., Riekens, B., Bruhn, H., Wiese, A., Seydel, U., and Leippe, M. (2003). Interaction of amoebapores and NK-lysin with symmetric phospholipid and asymmetric lipopolysaccharide/phospholipid bilayers. *Biochemistry* 42, 9804-9812.
- Hakomori, S. (1981). Glycosphingolipids in cellular interaction, differentiation, and oncogenesis. *Annu Rev Biochem* 50, 733-764.
- Harzer, K., Paton, B. C., Christomanou, H., Chatelut, M., Levade, T., Hiraiwa, M., and O'Brien, J. S. (1997). Saposins (sap) A and C activate the degradation of galactosylceramide in living cells. *FEBS Lett* 417, 270-274.
- Hawkins, C. A., de Alba, E., and Tjandra, N. (2005). Solution structure of human saposin C in a detergent environment. *J Mol Biol* 346, 1381-1392.
- Hecht, O., Van Nuland, N. A., Schleinkofer, K., Dingley, A. J., Bruhn, H., Leippe, M., and Grotzinger, J. (2004). Solution structure of the pore-forming protein of *Entamoeba histolytica*. *J Biol Chem* 279, 17834-17841.
- Henseler, M., Klein, A., Glombitza, G. J., Suzuki, K., and Sandhoff, K. (1996). Expression of the three alternative forms of the sphingolipid activator protein precursor in baby hamster kidney cells and functional assays in a cell culture system. *J Biol Chem* 271, 8416-8423.

- Hiraiwa, M., O'Brien, J. S., Kishimoto, Y., Galzicka, M., Fluharty, A. L., Ginns, E. I., and Martin, B. M. (1993a). Isolation, characterization, and proteolysis of human prosaposin, the precursor of saposins (sphingolipid activator proteins). *Arch Biochem Biophys* 304, 110-116.
- Hiraiwa, M., Soeda, S., Martin, B. M., Fluharty, A. L., Hirabayashi, Y., O'Brien, J. S., and Kishimoto, Y. (1993b). The effect of carbohydrate removal on stability and activity of saposin B. *Arch Biochem Biophys* 303, 326-331.
- Ho, M.-W., and J. S. O'Brien (1971). Gaucher's disease: deficiency of "acid" beta-glucuronidase and reconstitution of enzyme activity in vitro. *Proc Natl Acad Sci USA* 68, 2810-2813.
- Hofmann, A. F., Sjovall, J., Kurz, G., Radominska, A., Schteingart, C. D., Tint, G. S., Vlahcevic, Z. R., and Setchell, K. D. (1992). A proposed nomenclature for bile acids. *J Lipid Res* 33, 599-604.
- Holm, L., and Sander, C. (1993). Protein structure comparison by alignment of distance matrices. *J Mol Biol* 233, 123-138.
- Holm, L., and Sander, C. (1998). Touring protein fold space with Dali/FSSP. *Nucleic Acids Res* 26, 316-319.
- Hooft, R. W., Vriend, G., Sander, C., and Abola, E. E. (1996). Errors in protein structures. *Nature* 381, 272.
- Huwiler, A., and Zangemeister-Wittke, U. (2007). Targeting the conversion of ceramide to sphingosine 1-phosphate as a novel strategy for cancer therapy. *Crit Rev Oncol Hematol* 63, 150-159.
- Isupov, M. N., Obmolova, G., Butterworth, S., Badet-Denisot, M. A., Badet, B., Polikarpov, I., Littlechild, J. A., and Teplyakov, A. (1996). Substrate binding is required for assembly of the active conformation of the catalytic site in Ntn amidotransferases: evidence from the 1.8 Å crystal structure of the glutaminase domain of glucosamine 6-phosphate synthase. *Structure* 4, 801-810.
- Jeyakumar, M., Dwek, R. A., Butters, T. D., and Platt, F. M. (2005). Storage solutions: treating lysosomal disorders of the brain. *Nat Rev Neurosci* 6, 713-725.
- John, M., Wendeler, M., Heller, M., Sandhoff, K., and Kessler, H. (2006). Characterization of human saposins by NMR spectroscopy. *Biochemistry* 45, 5206-5216.
- Jones, M. J., and Murray, A. W. (1995). Evidence that ceramide selectively inhibits protein kinase C-alpha translocation and modulates bradykinin activation of phospholipase D. *J Biol Chem* 270, 5007-5013.
- Kabsch, W., Kabsch, H., and Eisenberg, D. (1976). Packing in a new crystalline form of glutamine synthetase from Escherichia coli. *J Mol Biol* 100, 283-291.
- Kang, S. J., and Cresswell, P. (2004). Saposins facilitate CD1d-restricted presentation of an exogenous lipid antigen to T cells. *Nat Immunol* 5, 175-181.
- Kelley, L. A., MacCallum, R. M., and Sternberg, M. J. (2000). Enhanced genome annotation using structural profiles in the program 3D-PSSM. *J Mol Biol* 299, 499-520.
- Kervinen, J., Tobin, G. J., Costa, J., Waugh, D. S., Wlodawer, A., and Zdanov, A. (1999). Crystal structure of plant aspartic proteinase prophytepsin: inactivation and vacuolar targeting. *Embo J* 18, 3947-3955.
- Kim, G. B., Miyamoto, C. M., Meighen, E. A., and Lee, B. H. (2004). Cloning and characterization of the bile salt hydrolase genes (bsh) from *Bifidobacterium bifidum* strains. *Appl Environ Microbiol* 70, 5603-5612.
- Kim, J. H., Krahn, J. M., Tomchick, D. R., Smith, J. L., and Zalkin, H. (1996). Structure and function of the glutamine phosphoribosylpyrophosphate amidotransferase glutamine site and communication with the phosphoribosylpyrophosphate site. *J Biol Chem* 271, 15549-15557.

- Kim, J. K., Yang, I. S., Rhee, S., Dauter, Z., Lee, Y. S., Park, S. S., and Kim, K. H. (2003). Crystal structures of glutaryl 7-aminocephalosporanic acid acylase: insight into autoproteolytic activation. *Biochemistry* 42, 4084-4093.
- Kim, Y., Kim, S., Earnest, T. N., and Hol, W. G. (2002). Precursor structure of cephalosporin acylase. Insights into autoproteolytic activation in a new N-terminal hydrolase family. *J Biol Chem* 277, 2823-2829.
- Kim, Y., Yoon, K., Khang, Y., Turley, S., and Hol, W. G. (2000). The 2.0 Å crystal structure of cephalosporin acylase. *Structure Fold Des* 8, 1059-1068.
- Kirby, L. C., Klein, R. A., and Coleman, J. P. (1995). Continuous Spectrophotometric Assay of Conjugated Bile-Acid Hydrolase. *Lipids* 30, 863-867.
- Kishimoto, Y., Hiraiwa, M., and O'Brien, J. S. (1992). Saposins: structure, function, distribution, and molecular genetics. *J Lipid Res* 33, 1255-1267.
- Kolesnick, R. (2002). The therapeutic potential of modulating the ceramide/sphingomyelin pathway. *J Clin Invest* 110, 3-8.
- Kolter, T., and Sandhoff, K. (2006). Sphingolipid metabolism diseases. *Biochim Biophys Acta* 1758, 2057-2079.
- Kraulis, P. J. (1991). Molscript - a Program to Produce Both Detailed and Schematic Plots of Protein Structures. *Journal of Applied Crystallography* 24, 946-950.
- Kumar, R. S., Brannigan, J. A., Prabhune, A. A., Pundle, A. V., Dodson, G. G., Dodson, E. J., and Suresh, C. G. (2006). Structural and functional analysis of a conjugated bile salt hydrolase from *Bifidobacterium longum* reveals an evolutionary relationship with penicillin V acylase. *J Biol Chem* 281, 32516-32525.
- Kumar, R. S., Brannigan, J. A., Pundle, A., Prabhune, A., Dodson, G. G., and Suresh, C. G. (2004). Expression, purification, crystallization and preliminary X-ray diffraction analysis of conjugated bile salt hydrolase from *Bifidobacterium longum*. *Acta Crystallogr D Biol Crystallogr* 60, 1665-1667.
- Kurz, M., Brachvogel, V., Matter, H., Stengelin, S., Thuring, H., and Kramer, W. (2003). Insights into the bile acid transportation system: the human ileal lipid-binding protein-cholyltaurine complex and its comparison with homologous structures. *Proteins* 50, 312-328.
- Lansmann, S., Schuette, C. G., Bartelsen, O., Hoernschemeyer, J., Linke, T., Weisgerber, J., and Sandhoff, K. (2003). Human acid sphingomyelinase. *Eur J Biochem* 270, 1076-1088.
- Law, B., and Rossie, S. (1995). The dimeric and catalytic subunit forms of protein phosphatase 2A from rat brain are stimulated by C2-ceramide. *J Biol Chem* 270, 12808-12813.
- Li, C. M., Park, J. H., Simonaro, C. M., He, X., Gordon, R. E., Friedman, A. H., Ehleiter, D., Paris, F., Manova, K., Hepbildikler, S., et al. (2002). Insertional mutagenesis of the mouse acid ceramidase gene leads to early embryonic lethality in homozygotes and progressive lipid storage disease in heterozygotes. *Genomics* 79, 218-224.
- Li, S., Smith, J. L., and Zalkin, H. (1999). Mutational analysis of *Bacillus subtilis* glutamine phosphoribosylpyrophosphate amidotransferase propeptide processing. *J Bacteriol* 181, 1403-1408.
- Liepinsh, E., Andersson, M., Ruysschaert, J. M., and Otting, G. (1997). Saposin fold revealed by the NMR structure of NK-lysin. *Nat Struct Biol* 4, 793-795.
- Linke, T., Wilkening, G., Sadeghlar, F., Mozcall, H., Bernardo, K., Schuchman, E., and Sandhoff, K. (2001). Interfacial regulation of acid ceramidase activity. Stimulation of ceramide degradation by lysosomal lipids and sphingolipid activator proteins. *J Biol Chem* 276, 5760-5768.
- Liu, A., Wenzel, N., and Qi, X. (2005). Role of lysine residues in membrane anchoring of saposin C. *Arch Biochem Biophys* 443, 101-112.

- Lopez-Gallego, F., Betancor, L., Hidalgo, A., Mateo, C., Guisan, J. M., and Fernandez-Lafuente, R. (2004). Optimization of an industrial biocatalyst of glutaryl acylase: stabilization of the enzyme by multipoint covalent attachment onto new amino-epoxy Sepabeads. *J Biotechnol* 111, 219-227.
- Mahfoud, R., Garmy, N., Maresca, M., Yahia, N., Puigserver, A., and Fantini, J. (2002). Identification of a common sphingolipid-binding domain in Alzheimer, prion, and HIV-1 proteins. *J Biol Chem* 277, 11292-11296.
- Malinina, L., Malakhova, M. L., Teplov, A., Brown, R. E., and Patel, D. J. (2004). Structural basis for glycosphingolipid transfer specificity. *Nature* 430, 1048-1053.
- Marti-Renom, M. A., Stuart, A. C., Fiser, A., Sanchez, R., Melo, F., and Sali, A. (2000). Comparative protein structure modeling of genes and genomes. *Annu Rev Biophys Biomol Struct* 29, 291-325.
- Maxfield, F. R., and McGraw, T. E. (2004). Endocytic recycling. *Nat Rev Mol Cell Biol* 5, 121-132.
- McVey, C. E., Walsh, M. A., Dodson, G. G., Wilson, K. S., and Brannigan, J. A. (2001). Crystal structures of penicillin acylase enzyme-substrate complexes: structural insights into the catalytic mechanism. *J Mol Biol* 313, 139-150.
- Menard, R., Feng, R., Storer, A. C., Robinson, V. J., Smith, R. A., and Krantz, A. (1991). Nature of papain products resulting from inactivation by a peptidyl O-acyl hydroxamate. *FEBS Lett* 295, 27-30.
- Merritt, E. A., and Murphy, M. E. P. (1994). Raster3d Version-2.0 - a Program for Photorealistic Molecular Graphics. *Acta Crystallographica Section D-Biological Crystallography* 50, 869-873.
- Möbius, W., van Donselaar, E., Ohno-Iwashita, Y., Shimada, Y., Heijnen, H. F., Slot, J. W., and Geuze, H. J. (2003). Recycling compartments and the internal vesicles of multivesicular bodies harbor most of the cholesterol found in the endocytic pathway. *Traffic* 4, 222-231.
- Murshudov, G. N., Vagin, A. A., and Dodson, E. J. (1997). Refinement of macromolecular structures by the maximum- likelihood method. *Acta Crystallographica Section D-Biological Crystallography* 53, 240-255.
- Nair, P. P., Gordon, M., and Reback, J. (1967). The enzymatic cleavage of the carbon-nitrogen bond in 3-alpha, 7-alpha, 12-alpha-trihydroxy-5-beta-cholan-24-oylglycine. *J Biol Chem* 242, 7-11.
- Niere, M., Dettloff, M., Maier, T., Ziegler, M., and Wiesner, A. (2001). Insect immune activation by apolipophorin III is correlated with the lipid-binding properties of this protein. *Biochemistry* 40, 11502-11508.
- O'Brien, J. S., Carson, G. S., Seo, H. C., Hiraiwa, M., and Kishimoto, Y. (1994). Identification of prosaposin as a neurotrophic factor. *Proc Natl Acad Sci U S A* 91, 9593-9596.
- O'Brien, J. S., Carson, G. S., Seo, H. C., Hiraiwa, M., Weiler, S., Tomich, J. M., Barranger, J. A., Kahn, M., Azuma, N., and Kishimoto, Y. (1995). Identification of the neurotrophic factor sequence of prosaposin. *Faseb J* 9, 681-685.
- O'Brien, J. S., and Kishimoto, Y. (1991). Saposin proteins: structure, function, and role in human lysosomal storage disorders. *Faseb J* 5, 301-308.
- Oinonen, C., and Rouvinen, J. (2000). Structural comparison of Ntn-hydrolases. *Protein Sci* 9, 2329-2337.
- Oinonen, C., Tikkanen, R., Rouvinen, J., and Peltonen, L. (1995). Three-dimensional structure of human lysosomal aspartylglucosaminidase. *Nat Struct Biol* 2, 1102-1108.
- Okino, N., He, X., Gatt, S., Sandhoff, K., Ito, M., and Schuchman, E. H. (2003). The reverse activity of human acid ceramidase. *J Biol Chem* 278, 29948-29953.
- Otwinowski, Z., and Minor, W. (1997). Processing of X-ray diffraction data collected in oscillation mode. In *Macromolecular Crystallography*, Pt A, pp. 307-326.

- Park, J. H., and Schuchman, E. H. (2006). Acid ceramidase and human disease. *Biochim Biophys Acta* 1758, 2133-2138.
- Pei, J., and Grishin, N. V. (2003). Peptidase family U34 belongs to the superfamily of N-terminal nucleophile hydrolases. *Protein Sci* 12, 1131-1135.
- Peitsch, M. C. (1996). ProMod and Swiss-Model: Internet-based tools for automated comparative protein modelling. *Biochem Soc Trans* 24, 274-279.
- Perrakis, A., Morris, R., and Lamzin, V. S. (1999). Automated protein model building combined with iterative structure refinement. *Nature Structural Biology* 6, 458-463.
- Petrek, M., Otyepka, M., Banas, P., Kosinova, P., Koca, J., and Damborsky, J. (2006). CAVER: a new tool to explore routes from protein clefts, pockets and cavities. *BMC Bioinformatics* 7, 316.
- Pettus, B. J., Chalfant, C. E., and Hannun, Y. A. (2002). Ceramide in apoptosis: an overview and current perspectives. *Biochim Biophys Acta* 1585, 114-125.
- Piatigorsky, J., Norman, B., Dishaw, L. J., Kos, L., Horwitz, J., Steinbach, P. J., and Kozmik, Z. (2001). J3-crystallin of the jellyfish lens: similarity to saposins. *Proc Natl Acad Sci U S A* 98, 12362-12367.
- Platt, F. M., Jeyakumar, M., Andersson, U., Dwek, R. A., and Butters, T. D. (2005). New developments in treating glycosphingolipid storage diseases. *Adv Exp Med Biol* 564, 117-126.
- Powell, H. R. (1999). The Rossmann Fourier autoindexing algorithm in MOSFLM. *Acta Crystallogr D Biol Crystallogr* 55, 1690-1695.
- Prabhune, A. A., and Sivaraman, H. (1990). Evidence for involvement of arginyl residue at the catalytic site of penicillin acylase from Escherichia coli. *Biochem Biophys Res Commun* 173, 317-322.
- Qi, X., and Chu, Z. (2004). Fusogenic domain and lysines in saposin C. *Arch Biochem Biophys* 424, 210-218.
- Qi, X., and Grabowski, G. A. (1998). Acid beta-glucosidase: intrinsic fluorescence and conformational changes induced by phospholipids and saposin C. *Biochemistry* 37, 11544-11554.
- Qi, X., and Grabowski, G. A. (2001). Differential membrane interactions of saposins A and C: implications for the functional specificity. *J Biol Chem* 276, 27010-27017.
- Qi, X., Leonova, T., and Grabowski, G. A. (1994). Functional human saposins expressed in Escherichia coli. Evidence for binding and activation properties of saposin C with acid beta-glucosidase. *J Biol Chem* 269, 16746-16753.
- Rafi, M. A., de Gala, G., Zhang, X. L., and Wenger, D. A. (1993). Mutational analysis in a patient with a variant form of Gaucher disease caused by SAP-2 deficiency. *Somat Cell Mol Genet* 19, 1-7.
- Read, R. J. (2001). Pushing the boundaries of molecular replacement with maximum likelihood. *Acta Crystallogr D Biol Crystallogr* 57, 1373-1382.
- Saad, A. F., Meacham, W. D., Bai, A., Anelli, V., Elojeimy, S., Mahdy, A. E., Turner, L. S., Cheng, J., Bielawska, A., Bielawski, J., et al. (2007). The Functional Effects of Acid Ceramidase Overexpression in Prostate Cancer Progression and Resistance to Chemotherapy. *Cancer Biol Ther* 6.
- Saarela, J., Oinonen, C., Jalanko, A., Rouvinen, J., and Peltonen, L. (2004). Autoproteolytic activation of human aspartylglucosaminidase. *Biochem J* 378, 363-371.
- Sandhoff, K., and Kolter, T. (1996). Topology of glycosphingolipid degradation. *Trends Cell Biol* 6, 98-103.
- Schmassmann, A., Fehr, H. F., Locher, J., Lillienau, J., Schteingart, C. D., Rossi, S. S., and Hofmann, A. F. (1993). Cholylsarcosine, a new bile acid analogue: metabolism and effect on biliary secretion in humans. *Gastroenterology* 104, 1171-1181.
- Schnabel, D., Schroder, M., and Sandhoff, K. (1991). Mutation in the sphingolipid activator protein 2 in a patient with a variant of Gaucher disease. *FEBS Lett* 284, 57-59.

- Schuette, C. G., Pierstorff, B., Huettler, S., and Sandhoff, K. (2001). Sphingolipid activator proteins: proteins with complex functions in lipid degradation and skin biogenesis. *Glycobiology* 11, 81R-90R.
- Schultz-Heienbrok, R., Remmel, N., Klingenstein, R., Rossocha, M., Sandhoff, K., Saenger, W., and Maier, T. (2006). Crystallization and preliminary characterization of three different crystal forms of human saposin C heterologously expressed in *Pichia pastoris*. *Acta Crystallograph Sect F Struct Biol Cryst Commun* 62, 117-120.
- Schulze, H., Schepers, U., and Sandhoff, K. (2007). Overexpression and mass spectrometry analysis of mature human acid ceramidase. *Biol Chem* 388, 1333-1343.
- Schuttelkopf, A. W., and van Aalten, D. M. (2004). PRODRG: a tool for high-throughput crystallography of protein-ligand complexes. *Acta Crystallogr D Biol Crystallogr* 60, 1355-1363.
- Selzner, M., Bielawska, A., Morse, M. A., Rudiger, H. A., Sindram, D., Hannun, Y. A., and Clavien, P. A. (2001). Induction of apoptotic cell death and prevention of tumor growth by ceramide analogues in metastatic human colon cancer. *Cancer Res* 61, 1233-1240.
- Shewale JG, S. V. (1997). Penicillin V acylase: its potential in the production of 6-amino penicillanic acid. *Enzyme Microb Technol* 20, 402-410.
- Simoes, I., and Faro, C. (2004). Structure and function of plant aspartic proteinases. *Eur J Biochem* 271, 2067-2075.
- Smith, J. L., Zaluzec, E. J., Wery, J. P., Niu, L., Switzer, R. L., Zalkin, H., and Satow, Y. (1994). Structure of the allosteric regulatory enzyme of purine biosynthesis. *Science* 264, 1427-1433.
- Smith, R. A., Coles, P. J., Spencer, R. W., Copp, L. J., Jones, C. S., and Krantz, A. (1988). Peptidyl O-acyl hydroxamates: potent new inactivators of cathepsin B. *Biochem Biophys Res Commun* 155, 1201-1206.
- Sreerama, N., and Woody, R. W. (2000). Estimation of protein secondary structure from circular dichroism spectra: comparison of CONTIN, SELCON, and CDSSTR methods with an expanded reference set. *Anal Biochem* 287, 252-260.
- Staab, J. F., Ginkel, D. L., Rosenberg, G. B., and Munford, R. S. (1994). A saposin-like domain influences the intracellular localization, stability, and catalytic activity of human acyloxyacyl hydrolase. *J Biol Chem* 269, 23736-23742.
- Storoni, L. C., McCoy, A. J., and Read, R. J. (2004). Likelihood-enhanced fast rotation functions. *Acta Crystallogr D Biol Crystallogr* 60, 432-438.
- Sun, Y., Witte, D. P., Zamzow, M., Ran, H., Quinn, B., Matsuda, J., and Grabowski, G. A. (2007). Combined saposin C and D deficiencies in mice lead to a neuronopathic phenotype, glucosylceramide and alpha-hydroxy ceramide accumulation, and altered prosaposin trafficking. *Hum Mol Genet* 16, 957-971.
- Suresh, C. G., Pundle, A. V., SivaRaman, H., Rao, K. N., Brannigan, J. A., McVey, C. E., Verma, C. S., Dauter, Z., Dodson, E. J., and Dodson, G. G. (1999). Penicillin V acylase crystal structure reveals new Ntn-hydrolase family members. *Nat Struct Biol* 6, 414-416.
- Svergun D., B. C., Koch M.H.J. (1995). CRYSTOL-a Program to Evaluate X-ray Solution Scattering of Biological Macromolecules from Atomic Coordinates. *J Appl Cryst* 28, 768-773.
- Svergun, D. I. (1992). Determination of the Regularization Parameter in Indirect-Transform Methods Using Perceptual Criteria. *J Appl Cryst* 25, 495-503.
- Svergun, D. I., Barberato, C., and Koch, M. H. J. (1995). CRYSTOL - a program to evaluate X-ray solution scattering of biological macromolecules from atomic coordinates. *J Appl Crystallogr* 28, 768-773.
- Tanaka, H., Doesburg, K., Iwasaki, T., and Mierau, I. (1999). Screening of lactic acid bacteria for bile salt hydrolase activity. *J Dairy Sci* 82, 2530-2535.

- Tanaka, H., Hashiba, H., Kok, J., and Mierau, I. (2000). Bile salt hydrolase of *Bifidobacterium longum*-biochemical and genetic characterization. *Appl Environ Microbiol* 66, 2502-2512.
- Tettamanti, G. (2004). Ganglioside/glycosphingolipid turnover: new concepts. *Glycoconj J* 20, 301-317.
- Tikkanen, R., Riikonen, A., Oinonen, C., Rouvinen, R., and Peltonen, L. (1996). Functional analyses of active site residues of human lysosomal aspartylglucosaminidase: implications for catalytic mechanism and autocatalytic activation. *Embo J* 15, 2954-2960.
- Vaccaro, A. M., Tatti, M., Ciaffoni, F., Salvioli, R., Serafino, A., and Barca, A. (1994). Saposin C induces pH-dependent destabilization and fusion of phosphatidylserine-containing vesicles. *FEBS Lett* 349, 181-186.
- Vagin, A., and Teplyakov, A. (1997). MOLREP: an automated program for molecular replacement. *Journal of Applied Crystallography* 30, 1022-1025.
- Vaguine, A. A., Richelle, J., and Wodak, S. J. (1999). SFCHECK: a unified set of procedures for evaluating the quality of macromolecular structure-factor data and their agreement with the atomic model. *Acta Crystallogr D Biol Crystallogr* 55, 191-205.
- van Echten, G., and Sandhoff, K. (1993). Ganglioside metabolism. Enzymology, Topology, and regulation. *J Biol Chem* 268, 5341-5344.
- Van Langen LM, O. N., Guranda DT, Van Rantwijk F, Svedas VK, Sheldon RA (2000). Penicillin acylase-catalyzed resolution of amines in aqueous organic solvents. *Tetrahedron Assy* 11.
- Viard, M., Parolini, I., Rawat, S. S., Fecchi, K., Sargiacomo, M., Puri, A., and Blumenthal, R. (2004). The role of glycosphingolipids in HIV signaling, entry and pathogenesis. *Glycoconj J* 20, 213-222.
- Vielhaber, G., Hurwitz, R., and Sandhoff, K. (1996). Biosynthesis, processing, and targeting of sphingolipid activator protein (SAP) precursor in cultured human fibroblasts. Mannose 6-phosphate receptor-independent endocytosis of SAP precursor. *J Biol Chem* 271, 32438-32446.
- Vonrhein, C., Blanc, E., Roversi, P., and Bricogne, G. (2006). Automated Structure Solution With autoSHARP. *Methods Mol Biol* 364, 215-230.
- Wallace, A. C., Laskowski, R. A., and Thornton, J. M. (1995). LIGPLOT: a program to generate schematic diagrams of protein-ligand interactions. *Protein Eng* 8, 127-134.
- Wang, Y., Grabowski, G. A., and Qi, X. (2003). Phospholipid vesicle fusion induced by saposin C. *Arch Biochem Biophys* 415, 43-53.
- Watanabe, R., Funato, K., Venkataraman, K., Futerman, A. H., and Riezman, H. (2002). Sphingolipids are required for the stable membrane association of glycosylphosphatidylinositol-anchored proteins in yeast. *J Biol Chem* 277, 49538-49544.
- Whitmore, L., and Wallace, B. A. (2004). DICHROWEB, an online server for protein secondary structure analyses from circular dichroism spectroscopic data. *Nucleic Acids Res* 32, W668-673.
- Wiederstein, M., and Sippl, M. J. (2007). ProSA-web: interactive web service for the recognition of errors in three-dimensional structures of proteins. *Nucleic Acids Res* 35, W407-410.
- Wilkening, G., Linke, T., and Sandhoff, K. (1998). Lysosomal degradation on vesicular membrane surfaces. Enhanced glucosylceramide degradation by lysosomal anionic lipids and activators. *J Biol Chem* 273, 30271-30278.
- Winau, F., Schwierzeck, V., Hurwitz, R., Remmel, N., Sieling, P. A., Modlin, R. L., Porcelli, S. A., Brinkmann, V., Sugita, M., Sandhoff, K., et al. (2004). Saposin C is required for lipid presentation by human CD1b. *Nat Immunol* 5, 169-174.
- Wright, C. S., Mi, L. Z., Lee, S., and Rastinejad, F. (2005). Crystal structure analysis of phosphatidylcholine-GM2-activator product complexes: evidence for hydrolase activity. *Biochemistry* 44, 13510-13521.

- Yang, J., Liu, X., Bhalla, K., Kim, C. N., Ibrado, A. M., Cai, J., Peng, T. I., Jones, D. P., and Wang, X. (1997). Prevention of apoptosis by Bcl-2: release of cytochrome c from mitochondria blocked. *Science* 275, 1129-1132.
- Yuan, W., Qi, X., Tsang, P., Kang, S. J., Illarionov, P. A., Besra, G. S., Gumperz, J., and Cresswell, P. (2007). Saposin B is the dominant saposin that facilitates lipid binding to human CD1d molecules. *Proc Natl Acad Sci U S A* 104, 5551-5556.