

6 Referenzen

- Agre,P. and Kozono,D. (2003). Aquaporin water channels: molecular mechanisms for human diseases. *FEBS Lett.*, **555**, 72-78.
- Alto,N.M., Soderling,S.H., Hoshi,N., Langeberg,L.K., Fayos,R., Jennings,P.A., and Scott,J.D. (2003). Bioinformatic design of A-kinase anchoring protein-in silico: a potent and selective peptide antagonist of type II protein kinase A anchoring. *Proc. Natl. Acad. Sci. U. S. A*, **100**, 4445-4450.
- Asirvatham,A.L., Galligan,S.G., Schillace,R.V., Davey,M.P., Vasta,V., Beavo,J.A., and Carr,D.W. (2004). A-kinase anchoring proteins interact with phosphodiesterases in T lymphocyte cell lines. *J. Immunol.*, **173**, 4806-4814.
- Baillie,G.S., Sood,A., McPhee,I., Gall,I., Perry,S.J., Lefkowitz,R.J., and Houslay,M.D. (2003). beta-Arrestin-mediated PDE4 cAMP phosphodiesterase recruitment regulates beta-adrenoceptor switching from Gs to Gi. *Proc. Natl. Acad. Sci. U. S. A*, **100**, 940-945.
- Birnbaumer,M., Antaramian,A., Themmen,A.P., and Gilbert,S. (1992). Desensitization of the human V2 vasopressin receptor. Homologous effects in the absence of heterologous desensitization. *J. Biol. Chem.*, **267**, 11783-11788.
- Bregman,D.B., Bhattacharyya,N., and Rubin,C.S. (1989). High affinity binding protein for the regulatory subunit of cAMP-dependent protein kinase II-B. Cloning, characterization, and expression of cDNAs for rat brain P150. *J. Biol. Chem.*, **264**, 4648-4656.
- Brown,D. (2003). The ins and outs of aquaporin-2 trafficking. *Am. J. Physiol Renal Physiol*, **284**, F893-F901.
- Brown,R.L., August,S.L., Williams,C.J., and Moss,S.B. (2003). AKAP7gamma is a nuclear RI-binding AKAP. *Biochem. Biophys. Res. Commun.*, **306**, 394-401.
- Burger,P.M., Mehl,E., Cameron,P.L., Maycox,P.R., Baumert,M., Lottspeich,F., De Camilli,P., and Jahn,R. (1989). Synaptic vesicles immunoisolated from rat cerebral cortex contain high levels of glutamate. *Neuron*, **3**, 715-720.
- Burns-Hamuro,L.L., Ma,Y., Kammerer,S., Reineke,U., Self,C., Cook,C., Olson,G.L., Cantor,C.R., Braun,A., and Taylor,S.S. (2003). Designing isoform-specific peptide disruptors of protein kinase A localization. *Proc. Natl. Acad. Sci. U. S. A*, **100**, 4072-4077.
- Bustamante,M., Hasler,U., Kotova,O., Chibalin,A.V., Mordasini,D., Rousselot,M., Vandewalle,A., Martin,P.Y., and Feraille,E. (2005). Insulin potentiates AVP-induced AQP2 expression in cultured renal collecting duct principal cells. *Am. J. Physiol Renal Physiol*, **288**, F334-F344.

- Carlisle Michel,J.J., Dodge,K.L., Wong,W., Mayer,N.C., Langeberg,L.K., and Scott,J.D. (2004). PKA-phosphorylation of PDE4D3 facilitates recruitment of the mAKAP signalling complex. *Biochem. J.*, **381**, 587-592.
- Catterall,W.A. (2000). Structure and regulation of voltage-gated Ca²⁺ channels. *Annu. Rev. Cell Dev. Biol.*, **16**, 521-555.
- Chen,J., Anderson,J.B., DeWeese-Scott,C., Fedorova,N.D., Geer,L.Y., He,S., Hurwitz,D.I., Jackson,J.D., Jacobs,A.R., Lanczycki,C.J., Liebert,C.A., Liu,C., Madej,T., Marchler-Bauer,A., Marchler,G.H., Mazumder,R., Nikolskaya,A.N., Rao,B.S., Panchenko,A.R., Shoemaker,B.A., Simonyan,V., Song,J.S., Thiessen,P.A., Vasudevan,S., Wang,Y., Yamashita,R.A., Yin,J.J., and Bryant,S.H. (2003). MMDB: Entrez's 3D-structure database. *Nucleic Acids Res.*, **31**, 474-477.
- Colledge,M., Dean,R.A., Scott,G.K., Langeberg,L.K., Huganir,R.L., and Scott,J.D. (2000). Targeting of PKA to glutamate receptors through a MAGUK-AKAP complex. *Neuron*, **27**, 107-119.
- Colledge,M. and Scott,J.D. (1999). AKAPs: from structure to function. *Trends Cell Biol.*, **9**, 216-221.
- Conti,M., Richter,W., Mehats,C., Livera,G., Park,J.Y., and Jin,C. (2003). Cyclic AMP-specific PDE4 phosphodiesterases as critical components of cyclic AMP signaling. *J. Biol. Chem.*, **278**, 5493-5496.
- de Groot,B.L., Frigato,T., Helms,V., and Grubmuller,H. (2003). The mechanism of proton exclusion in the aquaporin-1 water channel. *J. Mol. Biol.*, **333**, 279-293.
- Deen,P.M. and Knoers,N.V. (1998). Vasopressin type-2 receptor and aquaporin-2 water channel mutants in nephrogenic diabetes insipidus. *Am. J. Med. Sci.*, **316**, 300-309.
- Ding,G.H., Franki,N., Condeelis,J., and Hays,R.M. (1991). Vasopressin depolymerizes F-actin in toad bladder epithelial cells. *Am. J. Physiol.*, **260**, C9-16.
- Diviani,D., Abuin,L., Cotecchia,S., and Pansier,L. (2004). Anchoring of both PKA and 14-3-3 inhibits the Rho-GEF activity of the AKAP-Lbc signaling complex. *EMBO J.*, **23**, 2811-2820.
- Diviani,D., Soderling,J., and Scott,J.D. (2001). AKAP-Lbc anchors protein kinase A and nucleates Galphai 2-selective Rho-mediated stress fiber formation. *J. Biol. Chem.*, **276**, 44247-44257.
- Dodge,K.L., Khouangathiene,S., Kapiloff,M.S., Mouton,R., Hill,E.V., Houslay,M.D., Langeberg,L.K., and Scott,J.D. (2001). mAKAP assembles a protein kinase A/PDE4 phosphodiesterase cAMP signaling module. *EMBO J.*, **20**, 1921-1930.
- Dousa,T.P. (1999). Cyclic-3',5'-nucleotide phosphodiesterase isozymes in cell biology and pathophysiology of the kidney. *Kidney Int.*, **55**, 29-62.
- Dransfield,D.T., Bradford,A.J., Smith,J., Martin,M., Roy,C., Mangeat,P.H., and Goldenring,J.R. (1997). Ezrin is a cyclic AMP-dependent protein kinase anchoring protein. *EMBO J.*, **16**, 35-43.

Referenzen

- Fischer,v.M., Stahl,B., Walch-Solimena,C., Takei,K., Daniels,L., Khoklatchev,A., De Camilli,P., Sudhof,T.C., and Jahn,R. (1994). Localization of Rab5 to synaptic vesicles identifies endosomal intermediate in synaptic vesicle recycling pathway. *Eur. J. Cell Biol.*, **65**, 319-326.
- Francis,S.H., Poteet-Smith,C., Busch,J.L., Richie-Jannetta,R., and Corbin,J.D. (2002). Mechanisms of autoinhibition in cyclic nucleotide-dependent protein kinases. *Front Biosci.*, **7**, d580-d592.
- Fraser,I.D., Tavalin,S.J., Lester,L.B., Langeberg,L.K., Westphal,A.M., Dean,R.A., Marion,N.V., and Scott,J.D. (1998). A novel lipid-anchored A-kinase Anchoring Protein facilitates cAMP-responsive membrane events. *EMBO J.*, **17**, 2261-2272.
- Frokiaer,J., Marples,D., Valtin,H., Morris,J.F., Knepper,M.A., and Nielsen,S. (1999). Low aquaporin-2 levels in polyuric DI +/- severe mice with constitutively high cAMP-phosphodiesterase activity. *Am. J. Physiol.*, **276**, F179-F190.
- Fushimi,K., Sasaki,S., and Marumo,F. (1997). Phosphorylation of serine 256 is required for cAMP-dependent regulatory exocytosis of the aquaporin-2 water channel. *J. Biol. Chem.*, **272**, 14800-14804.
- Gasman,S., Chasserot-Golaz,S., Bader,M.F., and Vitale,N. (2003). Regulation of exocytosis in adrenal chromaffin cells: focus on ARF and Rho GTPases. *Cell Signal.*, **15**, 893-899.
- Gouraud,S., Laera,A., Calamita,G., Carmosino,M., Procino,G., Rossetto,O., Mannucci,R., Rosenthal,W., Svelto,M., and Valenti,G. (2002). Functional involvement of VAMP/synaptobrevin-2 in cAMP-stimulated aquaporin 2 translocation in renal collecting duct cells. *J. Cell Sci.*, **115**, 3667-3674.
- Gray,P.C., Scott,J.D., and Catterall,W.A. (1998). Regulation of ion channels by cAMP-dependent protein kinase and A-kinase anchoring proteins. *Curr. Opin. Neurobiol.*, **8**, 330-334.
- Gray,P.C., Tibbs,V.C., Catterall,W.A., and Murphy,B.J. (1997). Identification of a 15-kDa cAMP-dependent protein kinase-anchoring protein associated with skeletal muscle L-type calcium channels. *J. Biol. Chem.*, **272**, 6297-6302.
- Gustafson,C.E., Katsura,T., McKee,M., Bouley,R., Casanova,J.E., and Brown,D. (2000). Recycling of AQP2 occurs through a temperature- and bafilomycin-sensitive trans-Golgi-associated compartment. *Am. J. Physiol Renal Physiol*, **278**, F317-F326.
- Hall,A. (1998). Rho GTPases and the actin cytoskeleton. *Science*, **279**, 509-514.
- Henn,V., Edemir,B., Stefan,E., Wiesner,B., Lorenz,D., Theilig,F., Schmitt,R., Vossebein,L., Tamme,G., Beyermann,M., Krause,E., Herberg,F.W., Valenti,G., Bachmann,S., Rosenthal,W., and Klussmann,E. (2004). Identification of a novel A-kinase anchoring protein 18 isoform and evidence for its role in the vasopressin-induced aquaporin-2 shuttle in renal principal cells. *J. Biol. Chem.*, **279**, 26654-26665.
- Hoffmann,R., Abdel'AI,S., and Engels,P. (1998). Differential distribution of rat PDE-7 mRNA in embryonic and adult rat brain. *Cell Biochem. Biophys.*, **28**, 103-113.

Referenzen

- Holroyd,C., Kistner,U., Annaert,W., and Jahn,R. (1999). Fusion of endosomes involved in synaptic vesicle recycling. *Mol. Biol. Cell*, **10**, 3035-3044.
- Homma,S., Gapstur,S.M., Coffey,A., Valtin,H., and Dousa,T.P. (1991). Role of cAMP-phosphodiesterase isozymes in pathogenesis of murine nephrogenic diabetes insipidus. *Am. J. Physiol*, **261**, F345-F353.
- Houslay,M.D. and Adams,D.R. (2003). PDE4 cAMP phosphodiesterases: modular enzymes that orchestrate signalling cross-talk, desensitization and compartmentalization. *Biochem. J.*, **370**, 1-18.
- Hulme,J.T., Ahn,M., Hauschka,S.D., Scheuer,T., and Catterall,W.A. (2002). A novel leucine zipper targets AKAP15 and cyclic AMP-dependent protein kinase to the C terminus of the skeletal muscle Ca²⁺ channel and modulates its function. *J. Biol. Chem.*, **277**, 4079-4087.
- Jo,I., Ward,D.T., Baum,M.A., Scott,J.D., Coghlan,V.M., Hammond,T.G., and Harris,H.W. (2001). AQP2 is a substrate for endogenous PP2B activity within an inner medullary AKAP-signaling complex. *Am. J. Physiol Renal Physiol*, **281**, F958-F965.
- Johnson,D.A., Akamine,P., Radzio-Andzelm,E., Madhusudan,M., and Taylor,S.S. (2001). Dynamics of cAMP-dependent protein kinase. *Chem. Rev.*, **101**, 2243-2270.
- Kamsteeg,E.J., Heijnen,I., van Os,C.H., and Deen,P.M. (2000). The subcellular localization of an aquaporin-2 tetramer depends on the stoichiometry of phosphorylated and nonphosphorylated monomers. *J. Cell Biol.*, **151**, 919-930.
- Katsura,T., Gustafson,C.E., Ausiello,D.A., and Brown,D. (1997). Protein kinase A phosphorylation is involved in regulated exocytosis of aquaporin-2 in transfected LLC-PK1 cells. *Am. J. Physiol*, **272**, F817-F822.
- King,L.S., Kozono,D., and Agre,P. (2004). From structure to disease: the evolving tale of aquaporin biology. *Nat. Rev. Mol. Cell Biol.*, **5**, 687-698.
- Klussmann,E., Edemir,B., Pepperle,B., Tamma,G., Henn,V., Klauschenz,E., Hundsrucker,C., Maric,K., and Rosenthal,W. (2001). Ht31: the first protein kinase A anchoring protein to integrate protein kinase A and Rho signaling. *FEBS Lett.*, **507**, 264-268.
- Klussmann,E., Maric,K., and Rosenthal,W. (2000). The mechanisms of aquaporin control in the renal collecting duct. *Rev. Physiol Biochem. Pharmacol.*, **141**, 33-95.
- Klussmann,E., Maric,K., Wiesner,B., Beyermann,M., and Rosenthal,W. (1999). Protein kinase A anchoring proteins are required for vasopressin-mediated translocation of aquaporin-2 into cell membranes of renal principal cells. *J. Biol. Chem.*, **274**, 4934-4938.
- Klussmann,E., Tamma,G., Lorenz,D., Wiesner,B., Maric,K., Hofmann,F., Aktories,K., Valenti,G., and Rosenthal,W. (2001). An inhibitory role of Rho in the vasopressin-mediated translocation of aquaporin-2 into cell membranes of renal principal cells. *J. Biol. Chem.*, **276**, 20451-20457.

- Kuwahara,M., Fushimi,K., Terada,Y., Bai,L., Marumo,F., and Sasaki,S. (1995). cAMP-dependent phosphorylation stimulates water permeability of aquaporin-collecting duct water channel protein expressed in Xenopus oocytes. *J. Biol. Chem.*, **270**, 10384-10387.
- Lamaze,C., Chuang,T.H., Terlecky,L.J., Bokoch,G.M., and Schmid,S.L. (1996). Regulation of receptor-mediated endocytosis by Rho and Rac. *Nature*, **382**, 177-179.
- Lamaze,C., Fujimoto,L.M., Yin,H.L., and Schmid,S.L. (1997). The actin cytoskeleton is required for receptor-mediated endocytosis in mammalian cells. *J. Biol. Chem.*, **272**, 20332-20335.
- Lande,M.B., Jo,I., Zeidel,M.L., Somers,M., and Harris,H.W., Jr. (1996). Phosphorylation of aquaporin-2 does not alter the membrane water permeability of rat papillary water channel-containing vesicles. *J. Biol. Chem.*, **271**, 5552-5557.
- Liebenhoff,U. and Rosenthal,W. (1995). Identification of Rab3-, Rab5a- and synaptobrevin II-like proteins in a preparation of rat kidney vesicles containing the vasopressin-regulated water channel. *FEBS Lett.*, **365**, 209-213.
- Lorenz,D., Krylov,A., Hahm,D., Hagen,V., Rosenthal,W., Pohl,P., and Maric,K. (2003). Cyclic AMP is sufficient for triggering the exocytic recruitment of aquaporin-2 in renal epithelial cells. *EMBO Rep.*, **4**, 88-93.
- Lu,H., Sun,T.X., Bouley,R., Blackburn,K., McLaughlin,M., and Brown,D. (2004). Inhibition of endocytosis causes phosphorylation (S256)-independent plasma membrane accumulation of AQP2. *Am. J. Physiol Renal Physiol*, **286**, F233-F243.
- Luttrell,L.M., Roudabush,F.L., Choy,E.W., Miller,W.E., Field,M.E., Pierce,K.L., and Lefkowitz,R.J. (2001). Activation and targeting of extracellular signal-regulated kinases by beta-arrestin scaffolds. *Proc. Natl. Acad. Sci. U. S. A.*, **98**, 2449-2454.
- MacKenzie,S.J., Baillie,G.S., McPhee,I., MacKenzie,C., Seamons,R., McSorley,T., Millen,J., Beard,M.B., van Heeke,G., and Houslay,M.D. (2002). Long PDE4 cAMP specific phosphodiesterases are activated by protein kinase A-mediated phosphorylation of a single serine residue in Upstream Conserved Region 1 (UCR1). *Br. J. Pharmacol.*, **136**, 421-433.
- Marchmont,R.J. and Houslay,M.D. (1980). A peripheral and an intrinsic enzyme constitute the cyclic AMP phosphodiesterase activity of rat liver plasma membranes. *Biochem. J.*, **187**, 381-392.
- Maric,K., Oksche,A., and Rosenthal,W. (1998). Aquaporin-2 expression in primary cultured rat inner medullary collecting duct cells. *Am. J. Physiol.*, **275**, F796-F801.
- Marples,D., Christensen,S., Christensen,E.I., Ottosen,P.D., and Nielsen,S. (1995). Lithium-induced downregulation of aquaporin-2 water channel expression in rat kidney medulla. *J. Clin. Invest.*, **95**, 1838-1845.
- Marples,D., Frokiaer,J., Dorup,J., Knepper,M.A., and Nielsen,S. (1996). Hypokalemia-induced downregulation of aquaporin-2 water channel expression in rat kidney medulla and cortex. *J. Clin. Invest.*, **97**, 1960-1968.

Referenzen

- Marples,D., Frokiaer,J., and Nielsen,S. (1999). Long-term regulation of aquaporins in the kidney. *Am. J. Physiol.*, **276**, F331-F339.
- Marples,D., Schroer,T.A., Ahrens,N., Taylor,A., Knepper,M.A., and Nielsen,S. (1998). Dynein and dynactin colocalize with AQP2 water channels in intracellular vesicles from kidney collecting duct. *Am. J. Physiol.*, **274**, F384-F394.
- Marr,N., Bichet,D.G., Lonergan,M., Arthus,M.F., Jeck,N., Seyberth,H.W., Rosenthal,W., van Os,C.H., Oksche,A., and Deen,P.M. (2002). Heterooligomerization of an Aquaporin-2 mutant with wild-type Aquaporin-2 and their misrouting to late endosomes/lysosomes explains dominant nephrogenic diabetes insipidus. *Hum. Mol. Genet.*, **11**, 779-789.
- Matteoli,M., Takei,K., Cameron,R., Hurlbut,P., Johnston,P.A., Sudhof,T.C., Jahn,R., and De Camilli,P. (1991). Association of Rab3A with synaptic vesicles at late stages of the secretory pathway. *J. Cell Biol.*, **115**, 625-633.
- Michel,J.J. and Scott,J.D. (2002). AKAP mediated signal transduction. *Annu. Rev. Pharmacol. Toxicol.*, **42**, 235-257.
- Morello,J.P. and Bichet,D.G. (2001). Nephrogenic diabetes insipidus. *Annu. Rev. Physiol.*, **63**, 607-630.
- Moses,A.M. and Scheinman,S.J. (1993). Hereditary vasopressin resistance in man and mouse. *Ann. N. Y. Acad. Sci.*, **689**, 233-249.
- Mulders,S.M., Bichet,D.G., Rijss,J.P., Kamsteeg,E.J., Arthus,M.F., Lonergan,M., Fujiwara,M., Morgan,K., Leijendekker,R., van der,S.P., van Os,C.H., and Deen,P.M. (1998). An aquaporin-2 water channel mutant which causes autosomal dominant nephrogenic diabetes insipidus is retained in the Golgi complex. *J. Clin. Invest.*, **102**, 57-66.
- Murata,K., Mitsuoka,K., Hirai,T., Walz,T., Agre,P., Heymann,J.B., Engel,A., and Fujiyoshi,Y. (2000). Structural determinants of water permeation through aquaporin-1. *Nature*, **407**, 599-605.
- Murphy,C., Saffrich,R., Grummt,M., Gournier,H., Rybin,V., Rubino,M., Auvinen,P., Lutcke,A., Parton,R.G., and Zerial,M. (1996). Endosome dynamics regulated by a Rho protein. *Nature*, **384**, 427-432.
- Nejsum,L.N., Zelenina,M., Aperia,A., Frokiaer,J., and Nielsen,S. (2004). Bidirectional regulation of AQP2 trafficking and recycling: involvement of AQP2-S256 phosphorylation. *Am. J. Physiol Renal Physiol.* Dec 29
- Nielsen,S., Frokiaer,J., Marples,D., Kwon,T.H., Agre,P., and Knepper,M.A. (2002). Aquaporins in the kidney: from molecules to medicine. *Physiol Rev.*, **82**, 205-244.
- Noda,Y., Horikawa,S., Furukawa,T., Hirai,K., Katayama,Y., Asai,T., Kuwahara,M., Katagiri,K., Kinashi,T., Hattori,M., Minato,N., and Sasaki,S. (2004). Aquaporin-2 trafficking is regulated by PDZ-domain containing protein SPA-1. *FEBS Lett.*, **568**, 139-145.

- Novick,P. and Guo,W. (2002). Ras family therapy: Rab, Rho and Ral talk to the exocyst. *Trends Cell Biol.*, **12**, 247-249.
- Oksche,A. and Rosenthal,W. (1998). The molecular basis of nephrogenic diabetes insipidus. *J. Mol. Med.*, **76**, 326-337.
- Perry,S.J., Baillie,G.S., Kohout,T.A., McPhee,I., Magiera,M.M., Ang,K.L., Miller,W.E., McLean,A.J., Conti,M., Houslay,M.D., and Lefkowitz,R.J. (2002). Targeting of cyclic AMP degradation to beta 2-adrenergic receptors by beta-arrestins. *Science*, **298**, 834-836.
- Preston,G.M. and Agre,P. (1991). Isolation of the cDNA for erythrocyte integral membrane protein of 28 kilodaltons: member of an ancient channel family. *Proc. Natl. Acad. Sci. U. S. A.*, **88**, 11110-11114.
- Procino,G., Carmosino,M., Marin,O., Brunati,A.M., Contri,A., Pinna,L.A., Mannucci,R., Nielsen,S., Kwon,T.H., Svelto,M., and Valenti,G. (2003). Ser-256 phosphorylation dynamics of Aquaporin 2 during maturation from the ER to the vesicular compartment in renal cells. *FASEB J.*, **17**, 1886-1888.
- Qualmann,B. and Mellor,H. (2003). Regulation of endocytic traffic by Rho GTPases. *Biochem. J.*, **371**, 233-241.
- Richter,W. and Conti,M. (2002). Dimerization of the type 4 cAMP-specific phosphodiesterases is mediated by the upstream conserved regions (UCRs). *J. Biol. Chem.*, **277**, 40212-40221.
- Richter W, Jin SL, Conti M. (2005) Splice variants of the cyclic nucleotide phosphodiesterase PDE4D are differentially expressed and regulated in rat tissue. *Biochem J.* 2005 Feb 17.
- Rosenthal,W., Seibold,A., Antaramian,A., Lonergan,M., Arthus,M.F., Hendy,G.N., Birnbaumer,M., and Bichet,D.G. (1992). Molecular identification of the gene responsible for congenital nephrogenic diabetes insipidus. *Nature*, **359**, 233-235.
- Sasaki,S., Ishibashi,K., and Marumo,F. (1998). Aquaporin-2 and -3: representatives of two subgroups of the aquaporin family colocalized in the kidney collecting duct. *Annu. Rev. Physiol.*, **60**, 199-220.
- Schweda,F. and Kurtz,A. (2004). Cellular mechanism of renin release. *Acta Physiol Scand.*, **181**, 383-390.
- Sette,C. and Conti,M. (1996). Phosphorylation and activation of a cAMP-specific phosphodiesterase by the cAMP-dependent protein kinase. Involvement of serine 54 in the enzyme activation. *J. Biol. Chem.*, **271**, 16526-16534.
- Shepherd,M.C., Baillie,G.S., Stirling,D.I., and Houslay,M.D. (2004). Remodelling of the PDE4 cAMP phosphodiesterase isoform profile upon monocyte-macrophage differentiation of human U937 cells. *Br. J. Pharmacol.*, **142**, 339-351.
- Shukla,A., Hager,H., Corydon,T.J., Bean,A.J., Dahl,R., Vajda,Z., Li,H., Hoffmann,H.J., and Nielsen,S. (2001). SNAP-25-associated Hrs-2 protein colocalizes

- with AQP2 in rat kidney collecting duct principal cells. *Am. J. Physiol Renal Physiol*, **281**, F546-F556.
- Simon,H., Gao,Y., Franki,N., and Hays,R.M. (1993). Vasopressin depolymerizes apical F-actin in rat inner medullary collecting duct. *Am. J. Physiol*, **265**, C757-C762.
- Skalhegg,B.S. and Tasken,K. (2000). Specificity in the cAMP/PKA signaling pathway. Differential expression, regulation, and subcellular localization of subunits of PKA. *Front Biosci.*, **5**, D678-D693.
- Smith,F.D. and Scott,J.D. (2002). Signaling complexes: junctions on the intracellular information super highway. *Curr. Biol.*, **12**, R32-R40.
- Soderling,S.H. and Beavo,J.A. (2000). Regulation of cAMP and cGMP signaling: new phosphodiesterases and new functions. *Curr. Opin. Cell Biol.*, **12**, 174-179.
- Storm,R., Klussmann,E., Geelhaar,A., Rosenthal,W., and Maric,K. (2003). Osmolality and solute composition are strong regulators of AQP2 expression in renal principal cells. *Am. J. Physiol Renal Physiol*, **284**, F189-F198.
- Sudhof,T.C. (2004). The synaptic vesicle cycle. *Annu. Rev. Neurosci.*, **27**, 509-547.
- Tamma,G., Klussmann,E., Maric,K., Aktories,K., Svelto,M., Rosenthal,W., and Valenti,G. (2001). Rho inhibits cAMP-induced translocation of aquaporin-2 into the apical membrane of renal cells. *Am. J. Physiol Renal Physiol*, **281**, F1092-F1101.
- Tamma,G., Klussmann,E., Procino,G., Svelto,M., Rosenthal,W., and Valenti,G. (2003a). cAMP-induced AQP2 translocation is associated with RhoA inhibition through RhoA phosphorylation and interaction with RhoGDI. *J. Cell Sci.*, **116**, 1519-1525.
- Tamma,G., Wiesner,B., Furkert,J., Hahm,D., Oksche,A., Schaefer,M., Valenti,G., Rosenthal,W., and Klussmann,E. (2003b). The prostaglandin E2 analogue sulprostone antagonizes vasopressin-induced antidiuresis through activation of Rho. *J. Cell Sci.*, **116**, 3285-3294.
- Tasken,K. and Aandahl,E.M. (2004). Localized effects of cAMP mediated by distinct routes of protein kinase A. *Physiol Rev.*, **84**, 137-167.
- Tasken,K.A., Collas,P., Kemmner,W.A., Witczak,O., Conti,M., and Tasken,K. (2001). Phosphodiesterase 4D and protein kinase a type II constitute a signaling unit in the centrosomal area. *J. Biol. Chem.*, **276**, 21999-22002.
- Trotter,K.W., Fraser,I.D., Scott,G.K., Stutts,M.J., Scott,J.D., and Milgram,S.L. (1999). Alternative splicing regulates the subcellular localization of A-kinase anchoring protein 18 isoforms. *J. Cell Biol.*, **147**, 1481-1492.
- Valenti,G., Procino,G., Carmosino,M., Frigeri,A., Mannucci,R., Nicoletti,I., and Svelto,M. (2000). The phosphatase inhibitor okadaic acid induces AQP2 translocation independently from AQP2 phosphorylation in renal collecting duct cells. *J. Cell Sci.*, **113 (Pt 11)**, 1985-1992.

Referenzen

- van Balkom,B.W., Savelkoul,P.J., Markovich,D., Hofman,E., Nielsen,S., van der,S.P., and Deen,P.M. (2002). The role of putative phosphorylation sites in the targeting and shuttling of the aquaporin-2 water channel. *J. Biol. Chem.*, **277**, 41473-41479.
- Verbavatz,J.M., Brown,D., Sabolic,I., Valenti,G., Ausiello,D.A., Van Hoek,A.N., Ma,T., and Verkman,A.S. (1993). Tetrameric assembly of CHIP28 water channels in liposomes and cell membranes: a freeze-fracture study. *J. Cell Biol.*, **123**, 605-618.
- Verde,I., Pahlke,G., Salanova,M., Zhang,G., Wang,S., Coletti,D., Onuffer,J., Jin,S.L., and Conti,M. (2001). Myomegalin is a novel protein of the golgi/centrosome that interacts with a cyclic nucleotide phosphodiesterase. *J. Biol. Chem.*, **276**, 11189-11198.
- Wade,J.B., Stetson,D.L., and Lewis,S.A. (1981). ADH action: evidence for a membrane shuttle mechanism. *Ann. N. Y. Acad. Sci.*, **372**, 106-117.
- Wang,W., Li,C., Kwon,T.H., Knepper,M.A., Frokiaer,J., and Nielsen,S. (2002). AQP3, p-AQP2, and AQP2 expression is reduced in polyuric rats with hypercalcemia: prevention by cAMP-PDE inhibitors. *Am. J. Physiol Renal Physiol*, **283**, F1313-F1325.
- Wong,W. and Scott,J.D. (2004). AKAP signalling complexes: focal points in space and time. *Nat. Rev. Mol. Cell Biol.*, **5**, 959-970.
- Zaccolo,M. and Pozzan,T. (2002). Discrete microdomains with high concentration of cAMP in stimulated rat neonatal cardiac myocytes. *Science*, **295**, 1711-1715.
- Zerial,M. and McBride,H. (2001). Rab proteins as membrane organizers. *Nat. Rev. Mol. Cell Biol.*, **2**, 107-117.
- Zhang,J., Ma,Y., Taylor,S.S., and Tsien,R.Y. (2001). Genetically encoded reporters of protein kinase A activity reveal impact of substrate tethering. *Proc. Natl. Acad. Sci. U. S. A.*, **98**, 14997-15002.
- Zhang,K.Y., Card,G.L., Suzuki,Y., Artis,D.R., Fong,D., Gillette,S., Hsieh,D., Neiman,J., West,B.L., Zhang,C., Milburn,M.V., Kim,S.H., Schlessinger,J., and Bollag,G. (2004). A glutamine switch mechanism for nucleotide selectivity by phosphodiesterases. *Mol. Cell*, **15**, 279-286.