

8. Literaturverzeichnis

ABBOTT, A. (2004):

Microbiology: gut reaction.

Nature 427, 284-286

ABE, F.; N. ISHIBASHI u. S. SHIMAMURA (1995):

Effect of administration of bifidobacteria and lactic acid bacteria to newborn calves and piglets.

J. Dairy Sc. 78, 2838-2846

AGUNGPRIYONO, S., A.A. MACDONALD, K.Y.G. LEUS, N.KITAMURA, I.K.M. ADNYANE, G.P.

GOODALL, E.HONDO u. J.YAMADA (2000):

Immunohistochemical Study on the Distribution of Endocrine Cells in the Gastrointestinal Tract of the Babirusa, *Babyrousa babyrussa* (Suidae).

Anat. Histol. Embryol. 29; 173-178

AHMAD, M., D. G. SMITH u. S. MAHBOOB (2002):

Effect of NaCl on Heat Tolerance of *Enterococcus faecium* and *Enterococcus faecalis*.

OnLine J. Biol. Sc. 2, 483-484

ALEXOPOULOS, C.; I. E. GEORGIOULAKIS; A. TZIVARA; C. S. KYRIAKIS; A.GOVARIS u. S. C.

KYRIAKIS (2004):

Field evaluation of the effect of a probiotic-containing *Bacillus licheniformis* and *Bacillus subtilis* spores on the health status, performance, and carcass quality of grower and finisher pigs.

J. Vet. Med. A Physiol. Pathol. Clin. Med. 51, 306-312

ANLAUF, M., R. EISSELE, M.K.H. SCHAFFER, L.E. EIDEN, R. ARNOLD, U. PAUSER, G. KLOPPEL u.

E. WEIHE (2003):

Expression of the two isoforms of the vesicular monoamine transporter (Vmat1 and Vmat2) in the endocrine pancreas and pancreatic endocrine tumors.

J Histochem Cytochem 2003;51:1027-40.

AUDISIO, C., G. OLIVER u. M. C. APPELLA (1999):

Antagonistic effect of *Enterococcus faecium* J96 against human and poultry pathogenic *Salmonella* spp.

J. Food Prot. 62, 751-755

BABINSKA, I., T. ROTKIEWICZ u. I. OTROCKA-DOMAGALA (2005):

The effect of *Lactobacillus acidophilus* and *Bifidobacterium* spp. administration on the morphology of the gastrointestinal tract, liver and pancreas in piglets.

Pol J Vet Sc. 8(1);29-35

BAKKE-MCKELLEP, A.M., C.M. PRESS, G. BAEVERFJORD, A. KROGDAHL u. T. LANDSVERK (2000):

Changes in immune and enzyme histochemical phenotypes of cells in the intestinal mucosa of atlantic Salmon, *Salmo salar* L., with soybean meal-induced enteritis.

Journal of fish diseases, 23:(2) 115-127

BARKA, T. u. P.J. ANDERSON (1963):

Histochemistry. Theory, Practice and Bibliography.

Harper and Row Inc. New York Evanston London

BARRETT, R. E., A. DE LAHUNTA, W.J. ROENICK, R.E. HOFFER u. F.M. COONS (1976):

Four cases of portocaval shunt in dogs.

J. Small Anim. Pract. 17, 71-75

BARRETT, A. (1969):

Properties of lysosomal enzymes.

Lysosomes in biology and pathology II, 245-312

BAUM, B., E. M. LIEBLER-TENORIO, M. L. ENSS, J. F. POHLENZ u. G. BREVES (2002):

Saccharomyces Boulardii and *Bacillus Cereus* var. *voyoi* Influence the Morphology and the Mucins of the Intestine of Pigs.

Z. Gastroenterol. 40, 277-284

BAYLISS, W.M. u. E.H. STARLING (1902):

On the causation of the so-called "peripheral reflex secretion" of the pancreas.

Proc. Roy. Soc. B69, 352-353

BENNO, Y. u. T. MITSUOKA (1992):

Impact of *Bifidobacterium longum* on human fecal microflora.

Microbiol. Immunol. 36, 683-694

BENYACOUB, J., G. L. CZARNECKI-MAULDEN, C. CAVADINI, T. SAUTHIER, R. E. ANDERSON, E. J.

SCHIFFRIN u. T. VON DER WEID. (2003):

Supplementation of food with *Enterococcus faecium* (SF68) stimulates immune functions in young dogs.

J.Nutr. 133 (4), 1158-1162.

BERG, R. (1988):

Angewandte und topographische Anatomie der Haustiere.

3. Auflage, VEB Gustav Fischer Verlag Jena

BISPING, W. u. G. AMTSBERG (1988):

Farbatlas zur Diagnose bakterieller Infektionserreger der Tiere.

Parey, Berlin, Hamburg

BJÖRKMAN, N. u. M. SIBALIN (1967):

Acid phosphatase activity in cytoplasmic bodies of the absorbing intestinal cells from suckling pigs.

Biomedical and Life Sciences 23(5)

BLANKENSHIP, L.C., J.S. BAILEY, N.A. COX, N.J. STERN, R. BREWER, u. O. WILLIAMS (1993):

Two-step mucosal competitive exclusion flora treatment to diminish *Salmonellae* in commercial broiler chickens.

Poult. Sc. 72, 1667-1672

BOSI, P., M. MAZZONI, S. FILIPPI, P. TREVISI, L. CASINI, G. PETROSINO u. G. LALATTA-COSTERBOSA (2006):

A continuous dietary supply of free calcium formate negatively affects the parietal cell population and gastric RNA expression for H⁺/K⁺-ATPase in weaning pigs.

Jour. Of Nutrition 136(5), 1229-1235

- BOUDRY, G., J.P. LALLES, C.H. MALBERT, E. BOBILLIER u. B. SEVE (2002):
Diet-Related Adaption of the Small Intestine at Weaning in Pigs Is Functional Rather Than Structural.
Journal of Pediatric Gastroenterology and Nutrition, 34(2), 180-187
- BRADY, L.J., D.D. GALLAHER u. F.F. BUSTA (2000):
The role of probiotic cultures in the prevention of colon cancer.
J. Nutr. 130, 410S-414S
- BRANDAO, R.L., I.M. CASTRO, E.A. BAMBIRRA, S.C. AMARAL, L.G. FIETTO, M.J. TROPIA, M.J. NEVES, R.G. DOS SANTOS, N.C. GOMES u. J.R. NICOLI (1998):
Intracellular signal triggered by Cholera toxin in *Saccharomyces boulardii* and *Saccharomyces cerevisiae*.
Appl. Environ. Microbiol. 64, 564-568
- BRAUN, M., A. WAHEED u. K. VON FIGURA (1989):
Lysosomal acid phosphatase is transported to lysosomes via the cell surface.
EMBO J. 8, 3633-3640
- BREVES, G.; C. WALTER; M. BURMESTER u. B. SCHROEDER (2000):
In vitro studies on the effects of *Saccharomyces boulardii* and *Bacillus cereus* var. *toyoi* on nutrient transport in pig jejunum.
J. Anim Physiol Anim Nutr. (Berl) 84, 9-20
- BREVES, G. (2004):
Probiotische Mikroorganismen und ihre zellulären Wirkungsmechanismen.
Mikrobiologie und Tierernährung: 20. Hülsenberger Gespräche, Lübeck, Schriftenreihe der Wilhelm Schaumann Stiftung, 9-13
- BULL, H., P.G. MURRAY, D. THOMAS, A.M. FRASER u. P.N. NELSON (2001):
Acid phosphatases
J. Clin. Pathol. 55: 65-72
- BURGALETA, C., u. D.W. GOLDE (1977):
Effect of glucan on granulopoiesis and macrophage genesis in mice.
Cancer Res. 37, 1739-1742

BURNETT, G.S., u. J. HANNA (1963):

Effect of dietary calcium lactate and lactic acid on faecal *Escherichia coli* counts in pigs.
Nature 197, 815

BURSTONE, M.S. (1962):

Enzyme Histochemistry and its Application in the Study of Neoplasms
Academic Press, New York, London

BUTS, J. P., P. BERNASCONI, M. VAN CRAYNEST, P. MALDAGUE u. R. DE MEYER (1986):

Response of human and rat small intestinal mucosa to oral administration of *Saccharomyces boulardii*.
Pediatr. Res. 20, 192-196

BUTS, J.P., P. BERNASCONI, J. VAERMAN u. C. DIVE (1990):

Stimulation of secretory IgA and secretory component of immunoglobulins in small intestine of rats treated with *Saccharomyces boulardii*.
Dig. Dis. Sc. 35, 251-256

BYRNE, B.M., u. J. DANKERT (1979):

Volatile fatty acids and aerobic flora in the gastrointestinal tract of mice under various conditions.
Infect.Immun. 23, 559-563

CAPELLA, C. u. E. SOLCIA (1972):

The endocrine cells of the pig gastrointestinal mucoa and pancreas.
Arch. Histol. Jap.35;1-29

CASTAGLIUOLO, I., J.T. LAMONT, S.T. NIKULASSON u. C. POTHOUKAKIS (1996):

Saccharomyces boulardii protease inhibits *Clostridium difficile* toxin A effects in the rat ileum.
Infect. Immun. 64, 5225-5232

CASTAGLIUOLO, I., M.F. RIEGLER, L. VALENICK, J.T. LAMONT u. C. POTHOUKAKIS (1999):

Saccharomyces boulardii protease inhibits the effect of *Clostridium difficile* toxin A and B in human colonic mucosa.
Infect. Immun. 67, 302-307

CHENOWETH, C. u. D. SCHABERG (1990):

The epidemiology of enterococci.

European Journal of Clinical Microbiology and Infectious Diseases 9, 80-89.

CHOPRA, P., D. VERMA, M. KHULLAR, S. SAPRU u. S. MAHMOOD (2006):

Shiga Toxin Exposure Modulates Intestinal Brush Border Membrane Functional Proteins in Rabbit Ileum.

Molecular and Cellular Biochemistry 283(1-2), 85-92

CHRISTENSEN, H. R.; H. FROKIAER u. J. J. PESTKA (2002):

Lactobacilli differentially modulate expression of cytokines and maturation surface markers in murine dendritic cells.

J. Immunol. 168, 171-178

COLLINS, F:M. u. P.B. CARTER (1978):

Growth of *Salmonellae* in orally infected germfree mice

Infect. Immun. 21, 41-47

COLLINS, M. D. u. G. R. GIBSON (1999):

Probiotics, prebiotics and synbiotics: approaches for modulating the microbial ecology of the gut.

Am. J. Clin. Nutr. 69 1052S-1057S

COOKE, H.J., M. SIDHU u. Y.Z. WANG (1997):

Activation of 5-HT1P receptors on submucosal afferents subsequently triggers VIP neurons and chloride secretion in the guinea-pig colon.

J. Auton. Nerv. Syst. 66, 105-110.

COOKE, H.J.(2000):

Neurotransmitters in neuronal reflexes regulating intestinal secretion.

Ann. N. Y. Acad. Sc. 915, 77-80

CRAVEN, S.E. u. D.D. WILLIAMS (1997):

Inhibition of *Salmonella typhimurium* attachment to chicken cecal mucus by intestinal isolates of enterobacteriaceae and lactobacilli.

Avian Dis. 41, 548-558

DAVIDSON, J.N. u. D.C. HIRSH (1976):

Bacterial competition as a means of preventing neonatal diarrhea in pigs.

Infect. Immun. 13, 1773-1774

DELLMANN, H.D. u. J.A. EURELL (1998):

Textbook of veterinary histology

5th Edition, Williams & Wilkins, a waverly company, Baltimore, Philadelphia, London, Paris, Bangkok, Buenos Aires, Hong Kong, Munich, Sydney, Tokyo, Wroclaw

DEMLING, L. (1976):

Gastrointestinale Hormone.

Z. Gastroenterol. 14, 63-69

DE SIMONE, C., VESELY, R., NEGRI, R. Et al., (1988):

Enhancement of immune response of murine Peyer`s patches by a diet supplemented with yogurt.

Immunopharmacol Immuntoxicol 9:87-100

DEVRIESE, L.A., L.LAURIER, P. DE HERDT u. F. HAESEBROUCK (1992):

Enterococcal and streptococcal species isolated from faeces of calves, young cattle and dairy cows.

Journal of Applied Bacteriology 72, 29-31.

DEZFULI, B.S., L. GIARI, S. ARRIGHI, C. DOMENEGHINI u. G.BOSI (2003):

Influence of enteric helminths on the distribution of intestinal endocrine cells belonging to the diffuse endocrine system in brown trout, *Salmo trutta* L.

Journal of Fish Diseases 26; 155-166

DOBBINS, W. O. u. L.L. AUSTIN (1991):

Electron microscopic definition of intestinal endocrine cells: immunogold localization and review.

Ultrastruct. Path. 15, 15-39

DOCKRAY, G.J. (1989):

Gastrin, cholecystokinin (CCK), and the leukosulfakinins.

Biol. Bull. 177, 195-197

DOHMS, J. (2004):

Aspekte der Darmgesundheit und Chancen für den Einsatz von Probiotika.

Lohmann - Information 4, 27-30

DUGGAN, C., GANNON J., u. W. ALLAN WALKER, , (2002):

Protective nutrients and functional foods for the gastrointestinal tract.

Am J Clin Nutr 75, 789-808

DYCE K.M., W.O. SACK , C.J.G. WENSING (1991):

Anatomie der Haustiere, S.143-148

Ferdinand Enke Verlag Stuttgart 1991

EFTHIMIOU, S. (1996):

Dietary intake of β -1,3/1,6 glucans in juvenile dentex (*Dentex dentex*), Sparidae:

Effects on growth performance, mortalities and non-specific defense mechanisms.

J. Appl. Ichthyl. 12, 1-7

EGGEBRECHT, S. (2006):

Mündl. Mitteilung

Berlin, F.U. Berlin

EICHINGER-LEIN, E. (1992):

Laktobazillen zur Verdrängung von Enterobacteriaceae im Darm von Psittaciformes.

München, Univ., Diss.

FAN, M.Z., O.ADEOLA u. E.K. ASEM (1999):

Characterization of brush border membrane-bound alkaline phosphatase activity in different segments of the porcine small intestine.

J Nutr Biochem.;10(5):299-305.

FAN, M.Z., B. STOLL, R.JIANG u. D.G. BURRIN (2001):

Enterocyte Digestive Enzyme Activity Along the Crypt-Villus and Longitudinal Axes in the neonatal Pig Small Intestine.

Journal of Animal Sc. 79(2), 371-381

FAN, M.Z., O.ADEOLA,, E.K. ASEM und D. KING (2002):

Postnatal ontogeny of kinetics of porcine jejunal brush border membrane-bound alkaline phosphatase, aminopeptidase N and sucrase activities.

Comp Biochem Physiol A Mol Integr Physiol.;132(3):599-607.

FEHLHABER, K., u. P. JANETSCHKE (1992):

Ursachen von Gesundheitsschädigungen durch Lebensmittel.

In: K. Fehlhaber u. P. Janetschke (Hrsg.): Veterinärmedizinische Lebensmittelhygiene.

Fischer, Jena, Stuttgart, S. 23-126

FLACHOWSKY, G., K. TIROKE u. M. MATTHEY (1992):

Influence of yeast (*Saccharomyces cerevisiae* as Yea-sacc or Levaferm) on in sacco dry matter degradability and ruminal parameters of variously fed small ruminants.

Arch. Anim. Nutr. 42, 159-169

FLACHOWSKY, G. u. R. DAENICKE (1996):

Probiotika in der Rinderfütterung.

Übers. Tierernährung. 24, 62-68

FOX, S. M. (1988):

Probiotics: Intestinal inoculants for production animals.

Vet. Med. 83, 806-830

FUJITA, T. (1973):

Gastro-Entero-Pancreatic Endocrine System. A Cell-Biological Approach.

Igaku Shoin, Tokyo

FUJITA, T. (1988):

The endocrine cell system of digestive tract.

P.M. Motta and H. Fujita: Ultrastructure of Digestive Tract.

Martinius Nijhoff, Boston, 213-228

FULLER, R. (1977):

The importance of lactobacilli in maintaining normal microbial balance in the crop.

Br. Poult. Sc. 18, 85-94

FULLER, R. (1989):

Probiotics in man and animals.

J. appl. Bacteriol. 66, 365-378

Futtermittelverordnung (2006)

www.bmelv.de

GALDEANO, C. M. u. G. PERDIGON (2004):

Role of viability of probiotic strains in their persistence in the gut and in mucosal immune stimulation.

J. Appl. Microbiol. 97, 673-681

GEBERT, A., H.J. ROTHKÖTTER u. R. PABST (1994):

Cytokeratin 18 is an M-cell marker in porcine Peyer`s patches.

Cell Tissue Res. 276, 213-221

GEDEK, B. u. W. AMSELGRUBER (1990):

Mikrobieller Antagonismus: Zur Eliminierung von enteropathogenen *E. coli*-Keimen und Salmonellen aus dem Darm durch *Saccharomyces boulardii*.

In: R. Ottenjann, J. Müller u. Seifert (Hrsg.): Ökosystem Darm II .

Springer Verlag, Berlin, S.180-192

GEDEK, B., M. KIRCHGESSNER, S. WIEHLER, A. BOTT, U. EIDELSBURGER u. F. X. ROTH (1993):

Zur nutritiven Wirksamkeit von *Bacillus cereus* als Probiotikum in der Ferkelaufzucht.

2. Mitteilung – Einfluss auf Keimzahlen, Zusammensetzung und Resistenzeigenschaften der gastrointestinalen und faecalen Mikroflora.

Arch. Anim. Nutr. 44, 215-226

GEDEK, B. (1999):

Mode of actions of probiotics in poultry diets.

Proceedings 12th European Symposium on Poultry Nutrition, Veldhoven, The Netherlands, August 15-19, 83-90

GIBSON, G.R., u. M.R. ROBERFROID (1995):

Dietary modulation of the human colonic microbiota: Introducing the concept of prebiotics.

J. Nutr. 125, 1401-1412

GIBSON, G.R., u. A.L. MCCARTNEY (1998):

Modification of the gut flora by dietary means.

Biochem. Soc. Trans. 26, 222-228

GIONCHETTI, P.; F. RIZZELLO; A. VENTURI; P. BRIGIDI; D. MATTEUZZI; G. BAZZOCCHI; G.

POGGIOLI; M. MIGLIOLI u. M. CAMPIERI (2000):

Oral bacteriotherapy as maintenance treatment in patients with chronic pouchitis: a double-blind, placebo-controlled trial.

Gastroenterology 119, 305-309

GIONCHETTI, P.; F. RIZZELLO; U. HELWIG; A. VENTURI; K. M. LAMMERS; P. BRIGIDI; B. VITALI;

G. POGGIOLI; M. MIGLIOLI u. M. CAMOIERI (2003):

Prophylaxis of pouchitis onset with probiotic therapy: a double-blind, placebo-controlled trial.

Gastroenterology 124, 1202-1209

GISMONDO, M. R., L. DRAGO u. A. LOMBARDI (1999):

Review of probiotics available to modify gastrointestinal flora.

Int. J. Antimicrob. Agents. 12(4), 287-292

GÖRKE, B. (2000):

Untersuchung der Schleimhautmorphologie in Dünn- und Dickdarm nach oraler Applikation von *Saccharomyces boulardii* und *Bacillus cereus* var. *toyoi* beim Schwein.

Hannover, Tierärztl. Hochsch., Diss.

GÖRKE, B. u. E. LIEBLER-TENORIO (2001):

Probiotika: Gibt es wissenschaftliche Grundlagen für ihre Wirkung?

Dtsch. Tierärztl. Wochenschr. 108, 249-251

GOLDIN, B. R. (1998):

Health benefits of probiotics.

Br. J. Nutr. 80, S203-S207

GOLDSTONE, R. T., R.D. WILKES u. I.M. SEYBOLD (1981):

Hepatic function test-1: hepatic enzymology.

Vet. Med. Small Anim. Clin. 76, 642-645

GOMORI, G. (1952):

Microscopic histochemistry, principles and practice.
Chicago, Univ. Chicago Press

GOSSRAU, R. (1975):

Die Lysosomen des Darmepithels. Eine entwicklungsgeschichtliche Untersuchung.
Ergebn. Anat. Entwickl. Gesch. 51, H.5

GOTTSCHALK, S., A. WAHEED, B.SCHMIDT, P. LAIDLER u. K. VON FIGURA (1989):

Sequential processing of lysosomal acid phosphatase by a cytoplasmic thiol proteinase and a lysosomal aspartyl proteinase.
EMBO J. 8, 3215-3219

GOTZ, V., J.A. ROMANKIEWICZ, J. MOSS u. H.W. MURRAY (1979):

Prophylaxis against ampicillin- associated diarrhea with a lactobacillus preparation.
Am. J. Hosp. Pharm. 36: 754-757

GOYAL, R.K. u. I. HIRANO (1996):

The enteric nervous system.
N. Engl. J. Med. 334, 1106-1115

GRAM, L., J. MELCHIORSEN, B. SPANGGAARD, I. HUBER u. T.F. NIELSEN (1999):

Inhibition of *Vibrio anguillarum* by *Pseudomonas fluorescens* AH2, a possible probiotic treatment of fish.
Appl. Environ. Microbiol. 65, 969-973

GREEN, S.A., K.P. ZIMMER, G.GRIFFITHS u. I. MELLMANN (1987):

Kinetics of intracellular transport and sorting of lysosomal membrane and plasma membrane proteins.
J. Cell Biol., 105, 1227-1240

GROSSMANN, M.I. (1972):

Gastrointestinale hormones: some thoughts about clinical applications.
Scand. J. Gastroenterol. 7, 97-104

GRUBE, D. u. W.G. FORSSMANN (1979):

Morphology and function of the enterocrine cells.
Horm. Metabol. Res. 11, 589-606

GRUBE, D. (1986):

The endocrine cells of digestive system: amines, peptides, and modes of action.
Anat. Embryol. 175, 151-162

GU, X. u. D.LI (2004):

Effect of Dietary Crude Protein Level on Villous Morphology, Immune Status and
Histochemistry Parameters of Digestive Tract in Weaning Piglets.
Animal Feed Science and Technology 114(1-4); 113-126

GUPTA, P.; H. ANDREW; B. S. KIRSCHNER u. S. GUANDALINI (2000):

Is lactobacillus GG helpful in children with Crohn's disease? Results of a preliminary, open-
label study.
J. Pediatr. Gastroenterol. Nutr. 31, 453-457

HÄUSSINGER, D. u. W. GEROK, (1995):

Diagnostische Verfahren.
In: GEROK, W. und BLUM, H. E. (Hrsg.): Hepatologie. 2. Aufl., Urban & Schwarzenberg,
München, Wien, Baltimore, 3-44

HAHN, H. u. K. VOGT (1994):

Bacillus.
In: H. Hahn, D. Falke u. P. Klein (Hrsg.): Medizinische Mikrobiologie.
Springer, Berlin, S. 406-411

HAKANSON, R., J. ALUMETS u. F. SUNDLER (1978):

Classification of peptide-hormone-producing cells.
Lancet 1978;1:997.

HALL, R. L. (1985):

Laboratory evaluation of liver disease.
Vet. Clin. North. Am. Small Anim. Pract. 15, 3-18

- HALLER, D.; S. BLUM; C. BODE; W. P. HAMMES u. E. J. SCHIFFRIN (2000):
Activation of human peripheral blood mononuclear cells by nonpathogenic bacteria in vitro:
evidence of NK cells as primary targets.
Infect. Immun. 68, 752-759
- HAMPSON, D.J. (1986):
Alterations in piglet small intestinal structure at weaning.
Res. Vet. Sc. 40, 32-40
- HARRISON, G.A., R.W. HEMKEN, K.A. DAWSON, R.J. HARMON u. K.B. BARKER (1988):
Influence of addition of yeast culture supplement to diets of lactating cows on ruminal
fermentation and microbial populations.
Dairy Sc. 71, 2967-2975
- HARTMANN, J. (2000):
Darmflora stabilisieren, Durchfall verhindern!
Dtsch. Geflügelwirtsch. Magazin 35, 44-47
- HATCHER, G. E. u. R. S. LAMBRECHT (1993):
Augmentation of macrophage phagocytic activity by cell-free extracts of selected lactic
acidproducing bacteria.
J. Dairy Sc. 76, 2485-2492
- HAVENAAR, R. u. M. HUIS IN 'T VELD (1992):
Probiotics : a general view. 1. Aufl.
Verlag Elsevier Applied Science Publisher, Amsterdam
- HAYWARD, A.F. (1967):
Changes in fine structure of developing intestinal epithelium associated with pinocytosis.
J. Anat. 102, 57-90
- HELANDER, H.F. (1975):
Enzyme patterns and protein absorption in rat colon during development.
Acta. Anat. 91, 330-349

HELIN, T.; S. HAAHTELA u. T. HAAHTELA (2002):

No effect of oral treatment with an intestinal bacterial strain, *Lactobacillus rhamnosus* (ATCC 53103), on birch-pollen allergy: a placebo-controlled double-blind study.
Allergy 57, 243-246

HILGENBRINK, R., (1987):

Vorkommen, Bedeutung und Funktion der Paneth-Zelle. Eine Literaturübersicht.
Dissertation TiHo Hannover

HINRICHS, M. (2005):

Auswirkungen von *Enterococcus faecium* als Futterzusatzstoff auf die Keimflora und Zusammensetzung des Darminhalts sowie auf die Leistung von Broilern bei gleichzeitigem Einsatz eines Kokzidiostatikums
Diss. TiHo Hannover

HUIS IN'T VELD, J.H.J. (1998):

Probiotics and the control of pathogens: what do we know?
In: Naggy and Mulder (Hrsg.):
COST action 97-pathogenic micro-organisms in poultry and eggs, 5. Poultry and food safety. Proceedings of the workshop. Budapest, Hungary, 103-108

HUNGATE, R.E. (1984):

Microbes of nutritional importance in the alimentary tract.
Proc. Nutr. Soc. 43, 1-11

IBEN, C. u. J. LEIBETSEDER (1989):

Untersuchung der leistungsfördernden Wirkung von Toyocerin in der Ferkelaufzucht.
Wien. Tierärztl. Mschr. 76, 363-366

IJI, P.A. u. D.R. TIVEY (1998):

Natural and synthetic oligosaccharides in broiler chicken diets.
World's Poultry Sc. J. 54, 129-143

IJI, P.A. u. D.R. TIVEY (1999):

The use of oligosaccharides in broiler chicken diets.
Proceedings 12th European Symposium on Poultry Nutrition, Veldhoven, The Netherlands, August 15-19, 193-201

IZADNIA, F., C.T.WONG u. S.A. KOCOSHS (1998):

Brewer's yeast and *Saccharomyces boulardii* both attenuate *Clostridium difficile*-induced colonic secretion in rat.

Dig. Dis. Sc. 43, 2055-2060

JADAMUS, A. (2001):

Untersuchungen zur Wirksamkeit und Wirkungsweise des sporenbildenden *Bac. Cereus* var. *toyoi* im Verdauungstrakt von Broilern und Ferkeln.

FU- Berlin, Diss.

JAHN, H.U., R. ULLRICH, T.SCHNEIDER, R.M. LIEHR, H.L.SCHIEFERDECKER, H.HOLST u. M. ZEITZ (1996):

Immunological and trophical effects of *Saccharomyces boulardii* on the small intestine in healthy human volunteers.

Digestion.57(2); 95-104.

JIN, L.Z., Y.W. HO, N. ABDULLAH, u. S. JALALUDIN (1997):

Probiotics in poultry: Modes of action.

World's Poul. Sc. J. 53, 351-368

JIN, L.Z., R.R. MARQUARDT, X. ZHAO (2000):

A strain of *Enterococcus faecium* (18C23) inhibits adhesion of enterotoxigenic *Escherichia coli* K88 to porcine small intestine mucus.

Applied and Environmental Microbiology 66, 4200-4204.

JUNG, K., K. AHN u. C. CHAE (2006):

Decreased Activity of Brush Border Membrane-Bound Digestive Enzymes in Small Intestines from Pigs Experimentally Infected With Porcine Epidemic Diarrhea Virus.

Research in Veterinary Science 81(3), 310-315

KAHRS, D. (1991):

5 Jahre Erfahrung mit Toyocerin.

Lohmann Information, November/Dezember, 1-8

-
- KAILA, M.; E. ISOLAURI; E. SOPPI; E. VIRTANEN; S. LAINE u. H. ARVILOMMI (1992):
Enhancement of the circulating antibody secreting cell response in human diarrhea by a human *Lactobacillus* strain.
Pediatr. Res. 32, 141-144
- KIRCHGESSNER, M.; F. X. ROTH; U. EIDELSBURGER u. B. GEDEK (1993):
The nutritive efficiency of *Bacillus cereus* as a probiotic in the raising of piglets. 1. Effect on the growth parameters and gastrointestinal environment.
Arch. Tierernähr. 44, 111-121
- KIRCHHELLE, A., N. FRÜHWEIN u. D. TOBÜREN (1996):
Behandlung der persistierenden Diarrhö mit *S. boulardii* bei Reiserückkehrern.
Fortschr. Med. 11, 136-140
- KIRJAVAINEN, P. V.; S. J. SALMINEN u. E. ISOLAURI (2003):
Probiotic bacteria in the management of atopic disease: underscoring the importance of viability.
J. Pediatr. Gastroenterol. Nutr. 36, 223-227
- KLEIN, U. (1995):
Untersuchungen zur Wirksamkeit des Bioregulators Paciflor® in der Kälber- sowie Bullenmast. In: Schubert, R., Flachowsky, G. und Bitsch, R. (eds.): Vitamine und weitere Zusatzstoffe in der Ernährung von Mensch und Tier.
5. Symposium, 28./29.09.1995 in Jena/Thüringen, S. 494-499
- KLEIN, U. u. H.L. SCHMIDTS (1997):
Zum Einfluss des Bioregulators Paciflor auf die Morphologie der Dünndarmmukosa beim Schwein.
Proc. Soc. Nutr. Physiol. 6:41
- KNEIFEL, W. (2005):
Entwicklung und Qualitätsmerkmale von Probiotika.
Lohmann information 1/2005, 17-20
- KOOPMAN, J.P., G.W. WELLING, A.W. HUYBREGTS, J.W. MULLINK u. R.A. PRINS (1979):
Association of germ-free mice with intestinal microfloras.
Z. Versuchstierk. 23, 145-154

KOT, E. u. A. BEZKOROVAINY (1999):

Binding of ferric iron to the cell walls and membranes of *Bifidobacterium thermophilum*: effect of free radicals.

J. Agric. Food Chem. 47, 4606-4610

KRÜGER, W.F., J.W. BRADLEY u. R.H. PATTERSON (1977):

The interaction of gentian violet and lactobacillus organisms in the diet of leghorn hens.

Poultry Sc. 56, 1729 (abstract)

KRUIS, W.; E. SCHUTZ; P. FRIC; B. FIXA; G. JUDMAIER u. M. STOLTE (1997):

Double-blind comparison of an oral *Escherichia coli* preparation and mesalazine in maintaining remission of ulcerative colitis.

Aliment. Pharmacol. Ther. 11, 853-858

KRUIS, W., P. FRIC u. M. STOLTE (2001):

Maintenance of remission in ulcerative colitis is equally effective with *Escherichia coli* Nissle 1917 and with standard Mesalazine.(Abstr).

Gastroenterology 120, A127

KURUMAYA, H., G. OHTA u. Y.NAKANUMAS (1989):

Endocrine cells in the intrahepatic biliary tree in normal livers and hepatolithiasis.

Arch. Path. Lab. Med. 113, 143-147

KURZAK, P., M.A. EHRMANN, u. R.F. VOGEL (1998):

Diversity of lactic acid bacteria associated with ducks.

System. Appl. Microbiol. 21, 588-592

KUTSCHER W. u. H.WOLBERGS (1935):

Prostataphosphatase

Z. Physiol. Chem. 236: 237-240

LANGHENDRIES, J. P.; J. DETRY; J. VAN HEES; J. M. LAMBORAY; J. DARIMONT; M. J. MOZIN; M.

C. SECRETIN u. J. SENTERRE (1995):

Effect of a fermented infant formula containing viable bifidobacteria on the fecal flora composition and pH of healthy full-term infants.

J. Pediatr. Gastroenterol. Nutr. 21, 177-181

LAUN, T., F. LÄPPE u. B. PTOK (2002):

Zwischen Hoffnung und Humbug.

www.pharmazeutische-zeitung.de/fileadmin/pza/2002-47/titel.htm (Zugriff März 2007)

LECLERK, H., L.A. DEVRIESE u. D.A.A.MOSSEL (1996):

Taxonomical changes in intestinal (faecal) enterococci and streptococci: Consequences on their use as indicators of faecal contamination in drinking water.

Journal of Applied Bacteriology 81, 459-466.

LENHARDT, L. u. S. MOZES (2003):

Morphological and Functional Changes of Small Intestine in Growth-Stunted Broilers.

Acta Veterinaria Brno. 72(3); 353-+

LETTNER, F. (1987):

Zum Einsatz von Toyocerin im Milchaustauschfutter für Kälbermast.

Kraftfutter 5, 157-159

LIEBICH, H.G. (1992):

Funktionelle Histologie. Farbatlas und Kurzlehrbuch der mikroskopischen Anatomie der Haussäugetiere. 2. Aufl.

Verlag Schattauer, Stuttgart, New York

LILLEY, D. M. u. R. J. STILWELL (1965):

Probiotics: growth promoting factors produced by microorganisms.

Science 147, 747-748

LING, W. H.; R. KORPELA; H. MYKKANEN; S. SALMINEN u. O. HANNINEN (1994):

Lactobacillus strain GG supplementation decreases colonic hydrolytic and reductive enzyme activities in healthy female adults.

J. Nutr. 124, 18-23

LOH, C. (2003):

Ak Ernährungsmythen. 2. Probiotika

Hertzfeld, H. (HG): Tatort und Tatsache

Ferienakademie der Stipendiatinnen und Stipendiaten der Rosa-Luxemburg-Stiftung

Karl Dietz Verlag Berlin

- MACHA, M., TARAS, D., VAHJEN, W., ARINI, A. u. O.SIMON (2004):
Specific Enumeration of the probiotic strain *Enterococcus faecium* NCIMB 10415 in the intestinal tract and in faeces of piglets and sows.
Archives of Animal Nutrition, 58(6), 443-452
- MACHADO-CAETANO, J.A., M.T. PARAMÉS, M.J. BABO, A. SANTOS, A. BANDEIRA FERREIRA, A.A. FREITAS, M.R. CLEMENTE COELHO u. A. MATTHIOLI MATHEUS (1986):
Immunopharmacological effects of *Saccharomyces boulardii* in healthy human volunteers.
Int. J. Immunopharmacol. 8, 245-259
- MADSEN, K.; A. CORNISH; P. SOPER; C. MCKAIGNEY; H. JIJON; C. YACHIMEC; J.DOYLE; L. JEWELL u. C. DE SIMONE (2001):
Probiotic bacteria enhance murine and human intestinal epithelial barrier function.
Gastroenterology 121, 580-591
- MÄNNER, K. u. A. SPIELER (1997):
Probiotics in piglets – an alternative to traditional growth promoters.
Micoecol. Ther. 26, 243-256.
- MÄNNER, K., A. JADAMUS, W. VAHJEN, U. FRACKENPOHL u. O. SIMON (2002):
Effekte probiotischer Zusätze bei Puten auf Leistungsmerkmale und intestinale Mikroflora.
In: Rodehutsord, Markus (Hrsg): 7. Tagung Schweine- und Geflügelernährung, 78-80
Universitätsdruckerei, Halle (Saale)
- MAJAMAA, H. u. E. ISOLAURI (1997):
Probiotics: a novel approach in the management of food allergy.
J. Allergy Clin. Immunol. 99, 179-185
- MATHEW, A.G., S.E. CHATTIN, C.M. ROBBINS, u. D.A. Golden (1998):
Effect of direct-fed yeast culture on enteric microbial populations, fermentation acids and performance of weanling pigs.
J. Anim. Sc. 76, 2138-2145

-
- McFARLAND, L. V.; C. M. SURAWICZ; R. N. GREENBERG; R. FEKETY; G. W. ELMER; K. A. MOYER; S. A. MELCHER; K. E. BOWEN; J. L. COX u. Z. NOORANI (1995):
Prevention of beta-lactam-associated diarrhea by *Saccharomyces boulardii* compared with placebo.
Am. J. Gastroenterol. 90, 439-448
- MERRITT, A.M. u. B.E. KOLTS (1983):
Simultaneous pancreatic, gastric and hormonal responses to 2-deoxyglucose in the conscious pig.
Quart. J. exper. Physiol. 68, 69-75
- METSCHNIKOW, E.(1901):
Sur la flore du corps humain.
Lit. Philos. Soc. 45, 1-38
- METSCHNIKOW, E.(1907):
The prolongation of life: optimistic studies.
In: Chalmer, M. (Ed.), Heinemann, London, 161-183
- MEYDANI, S.N. u. W. K.HA (2000):
Immunological effects of yoghurt
Am. J. Clin Nutr. 71, 861- 872
- MIAO, Y.-G. (1988):
Study on the alkaline phosphatase in the midgut of domestic silkworm. *Brombyx mori*.
Acta Sericologica Sinica 14(3), 154-158
- MIAO, Y.-G. (2002):
Studies on the activity of the alkaline phosphatase in the midgut of infected silkworm, *Brombyx mori* L.
J. Appl. Ent. 126; 138-142
- MILLER, T.A., O.L. LLANOS, J.S.SWIERCZED, P.L. RAYFORD u. J.C.THOMPSON (1978):
Concentrations of gastrin and secretin in the alimentary tract of the cat.
Surgery 83, 90-93

- MIMURA, T.; F. RIZZELLO; U. HELWIG; A. VENTURI; P. BRIGIDI; D. MATTEUZZI; G. BAZZOCCHI; G. POGGIOLI; M. MIGLIOLI u. M. CAMPIERI (2004):
Once daily high dose probiotics therapy (VSL#3) for maintaining remission in recurrent or refractory pouchitis.
Gut 53, 108-114
- MOHAN, B., R. KADIRVEL, M. BHASKARAN, U. A. NATARAJAN (1995):
Effect of probiotic supplementation on serum / yolk cholesterol and on egg shell thickness in layers.
Br. Poult. Sc. 36, 799-803
- MOHAN, B., R. KADIRVEL, A. NATARAJAN, u. M. BHASKARAN (1996):
Effect of probiotic supplementation on growth, nitrogen utilisation and serum cholesterol in broilers.
Br. Poult. Sc. 37, 395-401
- MOLITOR, D (1996):
In vitro- und in vivo-Effekte eines Probiotikums (*Enterococcus faecium*) als Futterzusatz bei Hunden.
Hannover, Tierärztl. Hochsch., Diss.
- MONTOYA, C.A., P.LETERME u. J.P. LALLES (2006):
A Protein-Free Diet Alters Small Intestinal Architecture and Digestive Enzyme Activities in Rats.
Reproduction Nutrition Development 46(1), 49-56
- MOUWEN, J.M.V.M. (1971):
White scours in piglets. I. Stereomicroscopy of the mucosa of the small intestine.
Vet. Path. 8, 364-380
- MURALIDHARA, K.S., W.E. SANDINE, D.C. ENGLAND u. P.R. ELLIKER (1973):
Colonization of *Escherichia coli* and *Lactobacillus* in intestines of pigs.
J. Dairy Sc. 56, 1973
- NARISAWA, S., L. HUANG, A. IWASAKI, H.HASEGAWA, D.H. ALPERS u. J.L. MILLAN (2003):
Accelerated Fat Absorption in Intestinal Alkaline Phosphatase Knockout Mice.
Molecular and Cellular Biology 23(21): 7525-7530

NEWBOLD, C.J. (1995):

Microbial feed additives for ruminants.

In: R.J. Wallace u. A. Chesson (Hrsg.): Biotechnology in Animal Feeds and Animal Feeding.
Verlag Chemie, Weinheim, New York S. 260-278

NICHOLL, C.G., J.M. POLAK u. S.R. BLOOM (1985):

The hormonal regulation of food intake, digestion and absorption.

Annu. Rev. Nutr. 5, 213-239

NICKEL, R., A. SCHUMMER u. E. SEIFERLE (1987):

Lehrbuch der Anatomie der Haustiere.

Band III, 6. Auflage Verlag Paul Parey

NOBLE, C.J. (1978):

Carriage of group D streptococci in the human bowel.

Journal of Clinical Pathology 4, 1182-1186.

NOGOSSEK, M. (2001):

Untersuchungen zur Wirksamkeit von Pro- und Prebiotika gegen eine Infektion des
Verdauungstraktes mit *Escherichia coli* bei Kanarienvögeln (*Serinus canaria*)

Hannover, Tierärztl. Hochsch., Diss.

NOUSIAINEN, J. (1991):

Comperative observations on selected probiotics and olanquinox as feed additives for piglets
around weaning. 2. Effects on villus length and crypt depth in the jejunum, ileum, caecum and
colon.

J. Anim. Physiol. Anim. Nutr. 66: 224-230

NURMI, E. u. M. RANTALA (1973):

New aspects of Salmonella infection in broiler production.

Nature Vol. 241, 210-211

ONO, K. U. u. Y. SATOH (1981):

Ultrastructural lokalization of acid phosphatase activity in the small intestinal absorptive cells
of postnatal rats.

Histochem. 71, 501-512

- ORCI, L., D. BAETENS, M.P. DUBOIS u. C. RUFENER (1975):
Evidence for the D-cell of the pancreas secreting somatostatin.
Horm. Metab. Res. 7; 400-402
- PAN, H. u. M.D. GERSHON (2000):
Activation of intrinsic afferent pathways in submucosal ganglia of the guinea pig small intestine.
J. Neurosci. 20(9), 3295-3309.
- PARKER, R. B. (1974):
Probiotics, the other half of the antibiotic story.
Anim. Nutr. Health 29, 4-8
- PEARSE, A.G.E. (1976):
Neurotransmission and the APUD concept. In: R.E. Coupland and T. Fujita (Eds.):
Chromaffin, Enterochromaffin and Related Cells.
Elsevier, Amsterdam-Oxford-New York, 147-154
- PELTO, L.; E. ISOLAURI; E. M. LILIUS; J. NUUTILA u. S. SALMINEN (1998):
Probiotic bacteria down-regulate the milk-induced inflammatory response in milk-hypersensitive subjects but have an immunostimulatory effect in healthy subjects.
Clin. Exp. Allergy 28, 1474-1479
- PERDIGON, G.; M. E. DE MACIAS; S. ALVAREZ; G. OLIVER u. A. A. RUIZ HOLGADO (1986):
Effect of perorally administered lactobacilli on macrophage activation in mice.
Infect. Immun. 53, 404-410
- PERDIGON, G.; M. E. DE MACIAS; S. ALVAREZ; G. OLIVER u. A. P. RUIZ HOLGADO (1988):
Systemic augmentation of the immune response in mice by feeding fermented milks with *Lactobacillus casei* and *Lactobacillus acidophilus*.
Immunology 63, 17-23
- PETZOLD, K., u. E. MÜLLER (1986):
Tierexperimentelle und zellbiologische Untersuchungen zur Wirkung von *Saccharomyces cerevisiae* Hansen CBS 5926 bei der unspezifischen Steigerung des Immunsystems.
Arzneim. Forsch. / Drug Res. 36, 1085-1088

PLUSKE, J.R., D.J. HAMPSON u. I.H. WILLIAMS (1997):

Factors influencing the structure and function of the small intestine in the weaned pig: a review.

Livestock Prod. Sc. 51, 215-236

POLAK, J.M., L. GRIMELIUS, A.G.E. PEARSE, S.R. BLOOM u. A. ARIMURA (1975):

Growth-hormone release inhibiting hormone in gastrointestinal and pancreatic D-cells.

Lancet 1; 1220-1222

POLLMANN, D.S., D.M. DANIELSON, W.B. WREN, E.R. PEO, u. K.M. SHAHANI (1980):

Influence of *Lactobacillus acidophilus* inoculum on gnotobiotic and conventional pigs.

J. Anim. Sc. 51, 629-637

POTHOULAKIS, C., C.P. KELLY, M.A. JOSHI, N. GAO, C.J. O'KEANE, I. CASTAGLIUOLO u.

J.T. LAMONT (1993):

Saccharomyces boulardii inhibits *Clostridium difficile* toxin A binding and enterotoxicity in rat ileum.

Gastroenterology 104, 1108-1115

PRITCHARD, D.M. u. A.J.M. WATSON (1996):

Apoptosis and gastrointestinal pharmacology.

Pharmacol. Ther. 72, 149-169

QUEDNAU, M., S. AHRNÉ, G. MOLIN (1999):

Genomic relationships between *Enterococcus faecium* strains from different sources and with different antibiotic resistance profiles evaluated by restriction endonuclease analysis of total chromosomal DNA using EcoRI and PvuII.

Appl. Environ. Microbiol. 65, 1777-1780

RADECKI, S.V., P.K. KU, M.R. BENNINK, M.T. YOKOYAMA u. E.R. MILLER (1992):

Effect of Dietary Copper on Intestinal-Mucosa Enzyme-Activity, Morphology, and Turnover Rates in Weanling Pigs.

Journal of Animal Science 70(5), 1424-1431

RAPPORT, M.M., A.A. GREEN u. I.H. PAGE (1947):

Purification of the substance which is responsible for the vasoconstrictor activity of serum.

Fed. Proc. 6, 184

RECHT, J. (2005):

Einfluss Seltener Erden in Verbindung mit phyto-genen Zusatzstoffen auf Leistungsparameter beim Ferkel.

München, Tierärztl. Fakultät der LMU, Diss.

REDDY, B. S. (1998):

Prevention of colon cancer by pre- and probiotics: evidence from laboratory studies.

Br. J. Nutr. 80, S219-S223

REITER, K. (2005):

Einfluss des Probiotikums *Enterococcus faecium* SF 68 (NCIMB 10415) auf die Morphologie der Darmschleimhaut des Schweines.

Berlin, Vet-med. Diss. FU Berlin

REMBACKEN, B. J.; A. M. SNELLING; P. M. HAWKEY; D. M. CHALMERS u. A. T. AXON (1999):

Non-pathogenic *Escherichia coli* versus mesalazine for the treatment of ulcerative colitis: a randomised trial.

Lancet 354, 635-639

RICHTER-DAHLFORS, A., U. HECZKO, R.M. MELOCHE, B.B. FINLAY u. A.M. BUCHAN (1998):

Helicobacter pylori-infected human antral primary cell cultures: effect on gastrin cell function.

Am. J. Physiol. 275(3); 393-401

RIGGI, S.J., u. N.R. DI LUZIO (1961):

Identification of a reticuloendothelial stimulating agent in zymosan.

Am. J. Physiol. 200, 297-300

RINKINEN, M., K. JALAVA, E. WESTERMARCK, S. SALMINEN u. A. C. OUWEHAND (2003):

Interaction between probiotic lactic acid bacteria and canine enteric pathogens: a risk factor for intestinal *Enterococcus faecium* colonization?

Vet. Microbiol. 92, 111-119

ROEDIGER, W. E. (1980):

Role of anaerobic bacteria in the metabolic welfare of the colonic mucosa in man.

Gut 21, 793-798

ROLFE, R. D. (2000):

The Role of Probiotic Cultures in the Control of Gastrointestinal Health.

J. Nutr. 130, 396S-402S

ROLLE, M. (1993):

Medizinische Mikrobiologie, Infektions- und Seuchenlehre für Tierärzte, Biologen,
Agrarwissenschaftler und Interessierte aus anderen Fachgebieten.

Verlag Enke, Stuttgart

S.534

ROSEN, G. D. (1992):

Antibacterials in disease control and performance enhancement. Part II.

Feed Compounder 12, 20-23

ROTH, F. X. u. M. KIRCHGESSNER (1988a):

Nutritive Wirksamkeit von Toyocerin.

2.Mitteilung: Kälbermast

Landwirtsch. Forschung 41, 63-70

ROTH, F. X. u. M. KIRCHGESSNER (1988):

Nutritive Wirksamkeit von Toyocerin.

1.Mitteilung: Ferkelaufzucht

Landwirtsch. Forschung 41, 58-62

RYCHEN, G., u. C. SIMOES NUNES (1995):

Effects of three microbial probiotics on postprandial porto-arterial concentration differences of
glucose, galactose and amino-nitrogen in the young pig.

Br. J. Nutr. 74, 19-26

SANDERS, M.E. (2000):

Considerations for use of probiotic bacteria to modulate human health.

J. Nutr. 130, 384S-390S

SAVAGE, D.C. (1969):

Microbial interference between indigenous yeast and lactobacilli in the rodent stomach.

J. Bacteriol. 98, 1278-1283

SCHÄFER, A. u. P. HÖHN (1975):

Funktionelle Morphologie der Darmschleimhaut – Struktur und Enzymmuster bei der Ratte.
Gastroent. Stoffw. 8, 1-78

SCHENCK, M., u. E. KOLB (1990):

Grundriß der physiologischen Chemie.
8. Aufl. Fischer, Jena

SCHEUERMANN, S.E. (1993):

Effect of the probiotic Paciflor® (CIP 5832) on energy and protein metabolism in growing pigs.
Anim. Feed Sc. Technol. 41, 181-189

SCHIFFRIN, E. J.; F. ROCHAT; H. LINK-AMSTER; J. M. AESCHLIMANN u. A. DONNET-HUGHES
(1995):

Immunomodulation of human blood cells following the ingestion of lactic acid bacteria.
J. Dairy Sc. 78, 491-497

SCHMÖGER, R. (1960):

Bactisubtil zur Behandlung der Ernährungsstörung der Säuglinge und der Dysbakterie bei enteralen Antibiotikagaben.
Münchn. Med. Wochenschr. 24, 1213-1217

SCHWARZBACHER, S., (2004):

Vorlesung zum Anatomiekurs II. Unterer Verdauungstrakt.
Anatomisches Institut I, Universität Frankfurt

SCHWARZE, L. (1962):

Kompendium der Veterinär-Anatomie
Band II- Eingeweidesystem, S.83 – 107
VEB Gustav Fischer Verlag Jena

SEIFERT, H. S. u. F. GESSLER (1997):

Continuous oral application of probiotic *Bacillus cereus* - an alternative to the prevention of enterotoxemia?
Anim. Res. Dev. 46: 30-38

SHARMA, R. u. U. SCHUMACHER (1995):

Morphometric analysis of intestinal mucins under different dietary conditions and gut flora in rats.

Dig Dis Sc. 40(12);2532-9.

SHARMA, R. u. U. SCHUMACHER (1996):

The diet and gut microflora influence the distribution of enteroendocrine cells in the rat intestine.

Experientia 52(7); 664-70.

SHEIL, B.; J. MCCARTHY; L. O'MAHONY; M. W. BENNETT; P. RYAN; J. J. FITZGIBBON; B. KIELY; J. K. COLLINS u. F. SHANAHAN (2004):

Is the mucosal route of administration essential for probiotic function? Subcutaneous administration is associated with attenuation of murine colitis and arthritis.

Gut 53, 694-700

SHIDOJI, Y. u. S.H. KIM (2004):

Mutually Distinctive Gradients of Three Types of Intestinal Alkaline Phosphatase Along the longitudinal Axis of the Rat Intestine.

Digestion 70(1); 10-15

SIMON, O., A. JADAMUS u. W. VAHJEN (2001):

Probiotic feed additives – effectiveness and expected modes of action.

J. Anim. Feed Sc. 10, Suppl. 1, 51-67

SIMON, O., W. VAHJEN u. D. TARAS (2004):

Ernährung und intestinale Mikrobiota bei Schwein und Geflügel.

Mikrobiologie und Tierernährung: 20. Hülsenberger Gespräche, Lübeck, Schriftenreihe der Wilhelm Schaumann Stiftung, 112-124

SISSONS, J. W. (1989):

Potential of probiotic organisms to prevent diarrhoea and promote digestion in farm animals – a review.

J. Sc. Food Agric. 49, 1-13

SMOLLICH, A., u. G. MICHEL (1992):

Mikroskopische Anatomie der Haustiere, S.152- 170

Gustav Fischer Verlag Jena, Stuttgart

SOGABE, N., L.MIZOI, K.ASAHI, I. EZAWA u. M. GOSEKI-SONE (2004):

Enhancement by Lactose of Intestinal Alkaline Phosphatase Expression in Rats.

Bone 35(1): 249-255

SPILLMANN, H. (1997):

Probiotika und probiotische Mikroorganismen: Lebensmittel oder Heilmittel? - Ein Vergleich.

Dtsch. Molkerei-Zeitschrift 12, 515-525

STANLEY, V.G., R. OJO, S. WOLDESENBET, D.H. HUTCHINSON, u. L.F. KUBENA (1993):

The use of *Saccharomyces cerevisiae* to suppress the effects of aflatoxicosis in broiler chicks.

Poult. Sc. 72, 1867-1872

STAVRIC, S., u. E.T. KORNEGAY (1995):

Microbial probiotics for pigs and poultry.

In: R.J. Wallace u. A. Chesson (Hrsg.): Biotechnology in Animal Feeds and Animal Feeding.

Verlag Chemie, Weinheim, New York ,S. 205-231

STECCHINI, M. L.; M. DEL TORRE u. M. Munari (2001):

Determination of peroxy radical-scavenging of lactic acid bacteria.

Int. J. Food Microbiol. 64, 183-188

SURAWICZ, C. M., ELMER, G.W., SPEELMAN, P., MCFARLAND, L.V., CHIN, J. u. G. VAN BELLE (1989):

Prevention of antibiotic- associated diarrhea by *Saccharomyces boulardii*: a prospective study.

Gastroenterology 96, 981- 988

SURAWICZ, C. M. (2003):

Probiotics, antibiotic-associated diarrhoea and *Clostridium difficile* diarrhoea in humans.

Best. Pract. Res. Clin. Gastroenterol. 17, 775-783

SZAJEWSKA, H.; M. KOTOWSKA; J. Z. MRUKOWICZ; M. ARMANSKA u. W. MIKOLAJCZYK (2001):

Efficacy of *Lactobacillus GG* in prevention of nosocomial diarrhea in infants.

J. Pediatr. 138, 361-365

-
- TANG, M., B. LAARVELD, A.G. VAN KESSEL, D.L. HAMILTON, A. ESTRADA u. J.F. PATIENCE (1999):
Effect of Segregated Early Weaning on Postweaning Small Intestinal Development in Pigs.
Journal of Animal Science, 77(12), 3191-3200
- TARAS, D., VAHJEN, W., MACHA, M. u. O. SIMON (2005):
Response of performance characteristics and fecal consistency to long-lasting dietary supplementation with the probiotic strain *Bacillus cereus* var. toyoi to sows and piglets.
Archives of Animal Nutrition, 59(06), 405-417
- TARAS, D. (2007):
Mündl. Mitteilung
Institut für Tierernährung
Berlin, F.U. Berlin
- TEITELBAUM, J.E. u. W.A. WALKER (2002):
Nutritional Impact of Pre- and Probiotics as Protective Gastrointestinal Organisms.
Annu. Rev. Nutr. 22, 107-138
- TEJADA-SIMON, M.V., J.H. LEE, Z. USTUNUL, u. J.J. PESTKA (1995):
Indigestion of yogurt containing *Lactobacillus acidophilus* and *Bifidobacterium* to potentiate immunoglobulin A response to cholera toxin in mice.
J. Dairy Sc. 82, 649-660
- THELEN, U. (1997):
Einfluss zweier Varianten von *Bacillus cereus* als Probiotika beim frühentwöhnten Ferkel unter Einbeziehung ernährungsphysiologischer, mikrobiologischer, histologischer und hämatologischer Aspekte.
Giessen, Univ., Diss.
- TOSSENBERGER, J., Z. GARBACZ, Z. GYURMÁN, Z. SZEGEDI u. I. SZEGNER (1994):
Wirkung von Probiotika auf die Leistung von säugenden Sauen und deren Ferkel.
Znan. Prak. Poljopr. Tehnol. 24, 99-106
Zit. Nach C. Winckler (1997)

ULBRICH, M., M. HOFFMANN u. W. DROCHNER (2004):

Fütterung und Tiergesundheit.
Verlag Eugen Ulmer, Stuttgart

UNICOMP, L., (2006):

Low-Level Fluoroquinolone Resistance among *Campylobacter jejuni* Isolates in Australia.
Clinical Infectious Diseases 42,Nr.10, 1368-1374

VAGNE, M. (1970):

Les hormones gastro-intestinales. I.La gastrine et la sécrétine.
Path. Biol. 18, 887-907

VAHJEN, W., A. JADAMUS u. O. SIMON (2002):

Influence of a probiotic *Enterococcus faecium* strain on selected bacterial groups in the small intestine of growing turkey poults.
Arch. Anim. Nutr. 56, 419-429

VAHJEN, W., u. K. MÄNNER (2003):

The effect of a probiotic *Enterococcus faecium* product in diets of healthy dogs on bacteriological counts of *Salmonella spp.*, *Campylobacter spp.* and *Clostridium spp.* in faeces.
Arch. Anim. Nutr. 57 (3), 229-233

VAHJEN, W., M. MACHA, D. TARAS u. O. SIMON (2004):

Effects of an *Enterococcus faecium* probiotic in pigs: modification of intestinal bacterial metabolic activities.
Anim. Res. 53, 338

VAN BRIEL, C. (2002):

Veränderung der Anzahl und Verteilung von Plasmazellen und Lymphozytenpopulationen in der Darmschleimhaut des Schweines nach Applikation von Probiotika.
Hannover, Tierärztl. Hochsch., Diss.

VANDERHOOF, J. A.; D. B. WHITNEY; D. L. ANTONSON; T. L. HANNER; J. V. LUPO u. R. J. YOUNG (1999):

Lactobacillus GG in the prevention of antibiotic-associated diarrhea in children.
J. Pediatr. 135, 564-568

VENTURI, A.; P. GIONCHETTI; F. RIZZELLO; R. JOHANSSON; E. ZUCCONI; P. BRIGIDI; D.

MATTEUZZI u. M. CAMPIERI (1999):

Impact on the composition of the faecal flora by a new probiotic preparation: preliminary data on maintenance treatment of patients with ulcerative colitis.

Aliment. Pharmacol. Ther. 13, 1103-1108

VOLLRATH, L. (1969):

Über die Entwicklung des Dünndarms der Ratte. Morphologische, histochemische und experimentelle Untersuchungen.

Erg. Anant. Entw. Gesch. 41, H.2

VORBERG, U. (1994):

Einsatz von Probiotika als Futterzusatzstoff bei Hund und Katze.

Wien, Veterinärmed. Universität, Diss.

WAGNER, B. und K. LANDFRIED (1999):

Bullenmast mit Mikroorganismen.

Veredlungsproduktion 4: 80-81

WEBER, R. (2000):

Prüfung wechselseitiger Hemmeffekte verschiedener *Campylobacter jejuni*-Stämme bei der Kolonisation des Hühnerdarmes.

Hannover, Tierärztl. Hochsch., Diss.

WEYRAUCH, K.-D., S. BLÄHSER u. J. PERSCHBACHER (1987):

Somatostatin cells in the gastric mucosa of small ruminants.

Immunohistochemical study.

Acta anat. 128, 188-193

WEYRAUCH, K.D. B.SCHNORR u. H.S. GLASER (1989):

Development of gastrin-and somatostatin-containing cells in the pyloric gland region of the bovine abomasum.

Acta. Anat. (Basel) 135(3), 255-260.

WHITE, F., G. WENHAM, G.A. SHARMAN, A.S. JONES, E.A. RATTRAY u. I. McDONALD (1969):

Stomach function in relation to a scour syndrome in the piglet.

Br. J. Nut. 23, 847-857

WIESNER, E. u. R. RIBBECK (2000):

Lexikon der Veterinärmedizin 4.Aufl.

Enke, Stuttgart

WILHELM, C. u. P. M. SHAH (2002):

In-vitro-Aktivität von Linezolid

Chemother. J. 11, 113-116

WILLE, K.H. (2001):

On the functional morphology of the large intestinal mucosa of the ox (*Bos primigenius f. Taurus*), sheep (*Ovis ammon f. aries*) and goat (*Capra aegagrus f. hircus*).

Anat. Hist. Embryol. J. Veter. Med. Ser. C, 30(2); 65-76

WILLIAMS, P.E.V., C.A.G. TAIT, G.M. INNES, u. C.J. NEWBOLD (1991):

Effects of the inclusion of yeast culture (*Saccharomyces cerevisiae* plus growth medium) in the diet of dairy cows on milk yield and forage degradation and fermentation patterns in the rumen of sheep and steers.

Anim. Sc. 69, 3016-3026

WILSON, K.H., u. F. PERINI (1988):

Role of competition for nutrients in suppression of *Clostridium difficile* by the colonic microflora.

Infect. Immun. 56, 2610-2614

WUNDERLICH, P. F., L. BRAUN, I. FUMAGALLI, V. D'APUZZO, F. HEIM, M. KARLY, R. LODI, G.

POLITTA, F. VONBANK, u. L. ZELTNER (1989):

Double-blind report on the efficacy of lactic acid-producing *Enterococcus* SF68 in the prevention of antibiotic-associated diarrhoea and in the treatment of acute diarrhoea.

J. Int. Med. Res. 17(4), 333-338

YAMAUCHI, K. u. P. TARACHAI (2000):

Changes in intestinal villi, cell area and intracellular autophagic vacuoles related to intestinal function in chickens.

Br. Poult. Sc., 41(4): 416-423

YASUI, H.; N. NAGAOKA; A. MIKE; K. HAYAKAWA u. M. OHWAKI (1992):

Detection of Bifidobacterium strains that induce large quantities of IgA.
Microb. Ecol. Health Dis. 5, 155-162

ZIMMERMANN, B., E. BAUER u. R. MOSENTHIN (2001):

Pro- and prebiotics in pig nutrition- potential modulator of gut health?
J. Anim Feed Sc. 10, 47-56

ZENTEK, J., D. MOLITOR, u. J. KAMPHUES (1998):

Prüfung intestinaler Effekte eines Probiotikums (*Enterococcus faecium*) bei Hunden.
Kleintierpraxis 43, 187-197