

## A. Literaturverzeichnis

1. Abe, T., Sugihara, H., Nawa, H., Shigemoto, R., Mizuno, N. & Nakanishi, S. (1992). Molecular characterization of a novel metabotropic glutamate receptor mGluR5 coupled to inositol phosphate:Ca<sup>2+</sup> signal transduction. *Journal of Biological Chemistry* 267, 13361–13368.
2. Aguado, F., Carmona, M.A., Pozas, E., Aguijo, A., Martinez-Guijarro, F.J., Alcantara, S., Borrell, V., Yuste, R., Ibanez, C. F. & Soriano, E. (2003). BDNF regulates spontaneous correlated activity at early developmental stages by increasing synaptogenesis and expression of the K(+)/Cl(-) co-transporter KCC2. *Development* 130, 1267-1280.
3. Aguirre, J.A., Andbjær, B., Gonzalez-Baron, S., Hansson, A., Stromberg, I., Agnati, L.F. & Fuxe, K. (2001). Group I mGluR antagonist AIDA protects nigral DA cells from MPTP-induced injury. *Neuroreport* 12, 2615-2617.
4. Akyeli, J., Meier, J. & Grantyn R. (2004). BDNF promotes the maturation of GABAergic synapses by advancing KCC2 membrane integration. *Abstr., FENS*, 2003
5. Alagarsamy S., Sorensen S.D. & Conn P.J. (2001). Coordinate regulation of metabotropic glutamate receptors. *Current Opinion in Neurobiology* 11, 357-362.
6. Allen, J.W., Vicini, S. & Faden, A.I. (2001). Exacerbation of neuronal cell death by activation of group I metabotropic glutamate receptors: role of NMDA receptors and arachidonic acid release. *Exp.Neurol.* 169, 449-460.
7. Alsina, B., Vu, T. & Cohen-Cory, S. (2001). Visualizing synapse formation in arborizing optic axons in vivo: dynamics and modulation by BDNF. *Nat.Neurosci.* 4, 1093-1101.
8. Ango, F., Pin, J.P., Tu, J.C., Xiao, B., Worley, P.F., Bockaert, J. & Fagni, L. (2000). Dendritic and axonal targeting of type 5 metabotropic glutamate receptor is regulated by homer1 proteins and neuronal excitation. *J.Neurosci.* 20, 8710-8716.
9. Ango, F., Robbe, D., Tu, J.C., Xiao, B., Worley, P.F., Pin, J.P., Bockaert, J. & Fagni, L. (2002). Homer-dependent cell surface expression of metabotropic glutamate receptor type 5 in neurons. *Mol.Cell Neurosci.* 20, 323-329.

10. Aniksztejn, L., Otani, S. & Ben Ari, Y. (1992). Quisqualate Metabotropic Receptors Modulate NMDA Currents and Facilitate Induction of Long-Term Potentiation Through Protein Kinase C. *Eur.J.Neurosci.* 4, 500-505.
11. Anwyl, R. (1999). Metabotropic glutamate receptors: electrophysiological properties and role in plasticity. *Brain Res.Brain Res.Rev.* 29, 83-120.
12. Aramori, I. & Nakanishi, S. (1992). Signal transduction and pharmacological characteristics of a metabotropic glutamate receptor, mGluR1, in transfected CHO cells. *Neuron* 8, 757–765.
13. Awad, H., Hubert, G.W., Smith, Y., Levey, A.I. & Conn, P.J. (2000). Activation of metabotropic glutamate receptor 5 has direct excitatory effects and potentiates NMDA receptor currents in neurons of the subthalamic nucleus. *J.Neurosci.* 20, 7871-7879.
14. Baldauf, Z.B., Wang, X.P., Wang, S. & Bickford, M.E. (2003). Pretectotectal pathway: an ultrastructural quantitative analysis in cats. *J.Comp Neurol.* 464, 141-158.
15. Baldelli, P., Novara, M., Carabelli, V., Hernández-Guijo, J. M. & Carbone, E. (2002). BDNF upregulates evoked GABAergic transmission in developing hippocampus by potentiating presynaptic N- and P/Q-type  $\text{Ca}^{2+}$  channels signalling. *Eur.J.Neurosci.* 16, 2297–2310.
16. Balduzzi, R., Cupello, A. & Robello, M. (1992). Modulation of the expression of GABA(A) receptors in rat cerebellar granule cells by protein tyrosine kinases and protein kinase C. *Biochim.Biophys.Acta.* 1564, 263-270.
17. Banks, M.I., Hardie, J.B. & Pearce, R.A. Development of GABA(A) (2002). Receptor-Mediated Inhibitory Postsynaptic Currents in Hippocampus. *J.Neurophysiol.* 88, 3097-3107.
18. Barmashenko, G., Eysel, U.T. & Mittmann, T. (2001). Intracellular calcium signals in the surround of rat visual cortex lesions. *Neuroreport* 12, 3023-3028.
19. Basarsky, T.A., Parpura, V. & Haydon, P.G. (1994). Hippocampal synaptogenesis in cell culture: developmental time course of synapse formation, calcium influx, and synaptic protein distribution. *J.Neurosci.* 14, 6402-6411.
20. Battaglia, G., Bruno, V., Pisani, A., Centonze, D., Catania, M.V., Calabresi, P. & Nicoletti, F. (2001). Selective blockade of type-1 metabotropic glutamate receptors induces neuroprotection by enhancing gabaergic transmission. *Mol.Cell Neurosci.* 17, 1071-1083.

21. Baude, A., Nusser, Z., Roberts, J.D., Mulvihill, E., McIlhinney, R.A. & Somogyi, P. (1993) The metabotropic glutamate receptor (mGluR1 alpha) is concentrated at perisynaptic membrane of neuronal subpopulations as detected by immunogold reaction. *Neuron* 11, 771-787.
22. Ben-Ari, Y. (2001). Developing networks play a similar melody. *Trends.Neurosci.* 24, 353-360.
23. Ben-Ari, Y., Khazipov, R., Leinekugel, X., Caillard, O. & Gaiarsa, J.L. (1997). GABA<sub>A</sub>, NMDA and AMPA receptors: a developmentally regulated 'menage a trois'. *Trends.Neurosci.* 20, 523-529.
24. Benson, D.L., Colman, D.R. & Huntley, G.W. (2001). Molecules, maps and synapse specificity. *Nat.Rev.Neurosci.* 2, 899-909.
25. Benveniste, H., Drejer, J., Schousboe, A. & Diemer, N.H. (1984). Elevation of the extracellular concentrations of glutamate and aspartate in rat hippocampus during transient cerebral ischemia monitored by intracerebral microdialysis. *J.Neurochem.* 43, 1369-1374.
26. Binns, K.E. (1999). The synaptic pharmacology underlying sensory processing in the superior colliculus. *Prog.Neurobiol.* 59, 129-159.
27. Bizon, J.L., Lauterborn, J.C. & Gall, C.M. (1999). Subpopulations of striatal interneurons can be distinguished on the basis of neurotrophic factor expression. *J.Comp.Neurol.* 408, 283-98.
28. Boller, M. & Schmidt, M. (2001). Postnatal maturation of GABA(A) and GABA(C) receptor function in the mammalian superior colliculus. *Eur.J Neurosci.* 14, 1185-1193.
29. Bordi, F. & Ugolini, A. (1999). Group I metabotropic glutamate receptors: implications for brain diseases. *Prog.Neurobiol.* 59, 55-79.
30. Bormann, J. (2000) The 'ABC' of GABA receptors. *Trends.Pharmacol.Sci.* 21, 16-19.
31. Boxal A.R. (2000). GABAergic mIPSCs in rat cerebellum Purkinje cells are modulated by TrkB and mGluR1-mediated stimulation of Src. *Journal of Physiology* 524, 677-684.
32. Brandon, N.J., Delmas, P., Kittler, J.T., McDonald, B.J., Sieghart, W., Brown, D. A., Smart, T.G. & Moss, S. J. (2000). GABAA receptor phosphorylation and functional modulation in cortical neurons by a protein kinase C-dependent pathway. *J.Biol.Chem.* 275, 38856-38862.
33. Brandon, N.J., Jovanovic, J.N., Colledge, M., Kittler, J.T., Brandon, J.M., Scott, J.D. & Moss, S.J. (2003). A-kinase anchoring protein 79/150 facilitates the phosphorylation of

- GABA(A) receptors by cAMP-dependent protein kinase via selective interaction with receptor beta subunits. *Mol.Cell Neurosci.* 22, 87-97.
34. Brandon, N.J., Jovanovic, J.N., Smart, T.G. & Moss, S.J. (2002). Receptor for activated C kinase-1 facilitates protein kinase C-dependent phosphorylation and functional modulation of GABA(A) receptors with the activation of G-protein-coupled receptors. *J.Neurosci.* 22, 6353-6361.
35. Brandon, N.J., Uren, J.M., Kittler, J.T., Wang, H., Olsen, R., Parker, P.J. & Moss, S.J. (1999). Subunit-specific association of protein kinase C and the receptor for activated C kinase with GABA type A receptors. *J.Neurosci.* 19, 9228-9234.
36. Breysse, N., Baunez, C., Spooren, W., Gasparini, F. & Amalric, M. (2002). Chronic but not acute treatment with a metabotropic glutamate 5 receptor antagonist reverses the akinetic deficits in a rat model of parkinsonism. *J.Neurosci.* 22, 5669-5678.
37. Brünig, I., Penschuck, S., Berninger, B., Benson, J. & Fritschy, J. M.. (2001). BDNF reduces miniature inhibitory postsynaptic currents by rapid downregulation of GABA<sub>A</sub> receptor surface expression. *Eur.J.Neurosci.* 13, 1320-1328.
38. Brünig, I., Suter, A., Knuese, I., Luscher, B. & Fritschy, J.M. (2002). GABAergic terminals are required for postsynaptic clustering of dystrophin but not of GABA(A) receptors and gephyrin. *J.Neurosci.* 22, 4805-4813.
39. Bruno, V., Battaglia, G., Copani, A., Cespedes, V. M., Galindo, M. F., Cena, V., Sanchez-Prieto, J., Gasparini, F., Kuhn, R., Flor, P. J. & Nicoletti, F. (2001a). An activity-dependent switch from facilitation to inhibition in the control of excitotoxicity by group I metabotropic glutamate receptors. *Eur.J.Neurosci.* 13, 1469-1478.
40. Bruno, V., Battaglia, G., Copani, A., D'Onofrio, M., Di Iorio, P., De Blasi, A., Melchiorri, D., Flor, P. J. & Nicoletti, F. (2001b). Metabotropic glutamate receptor subtypes as targets for neuroprotective drugs. *J.Cereb.Blood Flow Metab* 21, 1013-1033.
41. Bruno, V., Ksiazek, I., Battaglia, G., Lukic, S., Leonhardt, T., Sauer, D., Gasparini, F., Kuhn, R., Nicoletti, F. & Flor, P. J. (2000). Selective blockade of metabotropic glutamate receptor subtype 5 is neuroprotective. *Neuropharmacology* 39, 2223-2230.
42. Buchner, K., Adamec, E., Beermann, M. L. & Nixon, R. A. (1999). Isoform-specific translocation of protein kinase C following glutamate administration in primary hippocampal neurons. *Brain Res.Mol.Brain Res.* 64, 222-235.

43. Bulleit, R. F. & Hsieh T. (2000). MEK inhibitors block BDNF-dependent and -independent expression of GABA<sub>A</sub> receptor subunit mRNAs in cultured mouse cerebellar granule neurons. *Dev.Brain Res.*, 119, 1–10.
44. Bushell, T.J., Lee, C.C., Shigemoto, R. & Miller, R.J. (1999). Modulation of synaptic transmission and differential localisation of mGlus in cultured hippocampal autapses. *Neuropharmacol.* 38, 1553-1567.
45. Bustó, R., Globus, M. Y., Neary, J. T. & Ginsberg, M. D. (1994). Regional alterations of protein kinase C activity following transient cerebral ischemia: effects of intraischemic brain temperature modulation. *J.Neurochem.* 63, 1095-1103.
46. Cadusseau J. & Roger M. (1985). Afferent projections to the superior colliculus in the rat, with special attention to the deep layers. *J.Hirnforsch.* 26(6), 667-81.
47. Calabresi, P., Centonze, D., Pisani, A. & Bernardi, G. (1999a). Metabotropic glutamate receptors and cell-type-specific vulnerability in the striatum: implication for ischemia and Huntington's disease. *Exp.Neurol.* 158, 97-108.
48. Calabresi, P., Marfia, G.A., Centonze, D., Pisani, A. & Bernardi, G. (1999b). Sodium influx plays a major role in the membrane depolarization induced by oxygen and glucose deprivation in rat striatal spiny neurons. *Stroke.* 30, 171-179.
49. Cartmell, J. & Schoepp, D.D. (2000). Regulation of neurotransmitter release by metabotropic glutamate receptors. *J.Neurochem.* 75, 889-907.
50. Catania, M.V., Landwehrmeyer, G.B., Testa, C.M., Standaert, D.G., Penney, J.B. Jr. & Young, A.B. (1994). Metabotropic glutamate receptors are differentially regulated during development. *Neuroscience* 61, 481–495.
51. Chakravarthy, B., Morley, P. & Whitfield, J. (1999). Ca<sup>2+</sup>-calmodulin and protein kinase Cs: a hypothetical synthesis of their conflicting convergences on shared substrate domains. *Trends Neurosci.* 22, 12-16.
52. Chavis, P., Fagni, L., Bockaert, J. & Lansman, J. B. (1995) Modulation of calcium channels by metabotropic glutamate receptors in cerebellar granule cells. *Neuropharmacology* 34, 929-937.
53. Chebib, M. & Johnston, G.A. (1999) The 'ABC' of GABA receptors: A brief review. *Clin.Exp.Pharmacol.Physiol.* 26, 937-940.

54. Cherubini, E. & Conti, F. (2001). Generating diversity at GABAergic synapses. *Trends Neurosci.*, 24(3), 155–162.
55. Choe, Y., Lee, B. J. & Kim, K. (2002). Participation of protein kinase C alpha isoform and extracellular signal-regulated kinase in neurite outgrowth of GT1 hypothalamic neurons. *J.Neurochem.* 83, 1412-1422.
56. Choi, S. & Lovinger, D. M. (1996) Metabotropic glutamate receptor modulation of voltage-gated  $\text{Ca}^{2+}$  channels involves multiple receptor subtypes in cortical neurons. *J.Neurosci.* 16, 36-45.
57. Chu, Z. & Hablitz, J.J. (1998). Activation of group I mGluRs increases spontaneous IPSC frequency in rat frontal cortex. *J.Neurophysiol.* 80, 621-627.
58. Cirone, J., Pothecary, C.A., Turner, J. P. & Salt, T.E. (2002a). Group I metabotropic glutamate receptors (mGluRs) modulate visual responses in the superficial superior colliculus of the rat. *J.Physiol.* 541, 895-903.
59. Cirone, J., Sharp, C., Jeffery, G. & Salt, T.E. (2002b). Distribution of metabotropic glutamate receptors in the superior colliculus of the adult rat, ferret and cat. *Neuroscience* 109, 779–786.
60. Clark, S.E., Garret, M. & Platt, B. (2001). Postnatal alterations of GABA receptor profiles in the rat superior colliculus. *Neuroscience*, 104(2), 441–454.
61. Conolly, C.N., Kittler, J.T., Thomas, P., Uren, J.M., Brandon, N.J., Smart, T.G. & Moss, S.J. (1999). Cell surface stability of gamma-aminobutyric acid type A receptors. Dependence on protein kinase C activity and subunit composition. *J.Biol.Chem.* 274, 36565–36572.
62. Cozzi, A., Meli, E., Carla, V., Pellicciari, R., Moroni, F. & Pellegrini-Giampietro, D. E. (2002). Metabotropic glutamate 1 (mGlu1) receptor antagonists enhance GABAergic neurotransmission: a mechanism for the attenuation of post-ischemic injury and epileptiform activity? *Neuropharmacology* 43, 119-130.
63. Covolan, L., Ribeiro, L.T., Longo, B.M. & Mello, L.E. (2000). Cell damage and neurogenesis in the dentate granule cell layer of adult rats after pilocarpine- or kainate-induced status epilepticus. *Hippocampus*. 10, 169-180.
64. Craig, A.M. & Boudin, H. (2001). Molecular heterogeneity of central synapses: afferent and target regulation. *Nature Neurosci.* 4, 569–578.

65. Cynader, M. & Berman, N. (1972). Receptive-field organization of monkey superior colliculus. *J.Neurophysiol.* 35, 187-201.
66. Dale, L.B., Bhattacharya, M., Anborgh, P.H., Murdoch, B., Bhatia, M., Nakanishi, S. & Ferguson, S.S. (2000). G protein-coupled receptor kinase-mediated desensitization of metabotropic glutamate receptor 1A protects against cell death. *J.Biol.Chem.* 275, 38213-38220.
67. Davies, P., Anderton, B., Kirsch, J., Konnerth, A., Nitsch, R. & Sheetz, M. (1998). First one in, last one out: the role of GABAergic transmission in generation and degeneration. *Prog. Neurobiol.* 55, 651–658.
68. Defagot, M.C., Villar, M.J. & Antonelli, M.C. (2002). Differential localization of metabotropic glutamate receptors during postnatal development. *Dev.Neurosci.* 24, 272-282.
69. Deller, T., Frotscher, M. & Nitsch, R. (1995). Morphological evidence for the sprouting of inhibitory commissural fibers in response to the lesion of the excitatory entorhinal input to the rat dentate gyrus. *J.Neurosci* 15, 6868-6878.
70. Drew, G.M. & Vaughan, C.W. (2004). Multiple metabotropic glutamate receptor subtypes modulate GABAergic neurotransmission in rat periaqueductal grey neurons in vitro. *Neuropharmacol.* 46, 927-934.
71. Dumoulin, A., Lévi, S., Riveau, B., Gasnier, B. & Triller, A.. (2000). Formation of mixed glycine and GABAergic synapses in cultured spinal cord neurons. *Eur.J.Neurosci.* 12, 3883–3892.
72. Dumoulin, A., Rostaing, P., Bedet, C., Levi, S., Isambert, M.F., Henry, J.P., Triller, A. & Gasnier, B. (1999). Presence of the vesicular inhibitory amino acid transporter in GABAergic and glycinergic synaptic terminal boutons. *J. Cell Sci.* 112, 811–823.
73. Durand, G.M., Kovalchuk, Y. & Konnerth, A. (1996). Long-term potentiation and functional synapse induction in developing hippocampus. *Nature* 381, 71–75.
74. Durkin, J. P., Tremblay, R., Buchan, A., Blosser, J., Chakravarthy, B., Mealing, G., Morley, P. & Song, D. (1996). An early loss in membrane protein kinase C activity precedes the excitatory amino acid-induced death of primary cortical neurons. *J.Neurochem.* 66, 951-962.
75. Durkin, J. P., Tremblay, R., Chakravarthy, B., Mealing, G., Morley, P., Small, D. & Song, D. (1997). Evidence that the early loss of membrane protein kinase C is a necessary step in the

- excitatory amino acid-induced death of primary cortical neurons. *J.Neurochem.* 68, 1400-1412.
76. Ebihara, S., Shirato, K., Harata, N. & Akaike, N. (1995) Gramicidin-perforated patch recording: GABA response in mammalian neurones with intact intracellular chloride. *J.Physiol* (Lond) 434:77– 86.
77. Edwards, F.A., Konnerth, A., Sakmann, B. & Takahashi, T. (1989). A thin slice preparation for patch clamp recordings from neurones of the mammalian central nervous system. *Pflugers Arch.* 414, 600-612.
78. Edwards, M.A. & Jacobson, M. (1984). Effects of permeability of midtectal barriers in goldfish on compression of the visuotectal projection rostrally and regenerative escape caudally. *J.Comp.Neurol.* 226, 141-153.
79. Edwards, M.D., White, A.M. & Platt, B. (2002). Characterisation of rat superficial superior colliculus neurones: firing properties and sensitivity to GABA. *Neuroscience* 110, 93-104.
80. Essrich, C., Lorez, M., Benson, J.A., Fritschy, J.M., Lüscher, B., (1998). Postsynaptic clustering of major GABA<sub>A</sub> receptor subtypes requires the gamma 2 subunit and gephyrin. *Nature Neurosci.* 1, 563–571.
81. Eysel, U.T., Schweigart, G., Mittmann, T., Eyding, D., Qu, Y., Vandesande, F., Orban, G. & Arckens, L. (1999). Reorganization in the visual cortex after retinal and cortical damage. *Restor.Neurol.Neurosci.* 15, 153-164.
82. Fagni, L., Chavis, P., Ango, F. & Bockaert, J. (2000). Complex interactions between mGluRs, intracellular Ca<sup>2+</sup> stores and ion channels in neurons. *Trends.Neurosci.* 23, 80-88.
83. Faux, M. C. & Scott, J. D. (1997b). Regulation of the AKAP79-protein kinase C interaction by Ca<sup>2+</sup>/Calmodulin. *J Biol.Chem.* 272, 17038-17044.
84. Feng, J., Cai, X., Zhao, J. & Yan, Z. (2001). Serotonin receptors modulate GABA(A) receptor channels through activation of anchored protein kinase C in prefrontal cortical neurons. *J.Neurosci.* 21, 6502-6511.
85. Feng, J., Chi, P., Blanpied, T.A., Xu, Y., Magarinos, A.M., Ferreira, A., Takahashi, R.H., Kao, H.T., McEwen, B.S., Ryan, T.A., Augustine, G.J. & Greengard, P. (2002). Regulation of neurotransmitter release by synapsin III. *J.Neurosci.* 22, 4372-4380.

86. Fiala, A., Muller, U. & Menzel, R. (1999). Reversible downregulation of protein kinase A during olfactory learning using antisense technique impairs long-term memory formation in the honeybee, *Apis mellifera*. *J.Neurosci.* 19, 10125-10134.
87. Fiala, J.C., Kirov, S.A., Feinberg, M.D., Petrak, L.J., George, P., Goddard, C.A. & Harris, K.M. (2003). Timing of neuronal and glial ultrastructure disruption during brain slice preparation and recovery in vitro. *J.Comp.Neurol.* 465, 90-103.
88. Filippova, N., Sedelnikova, A., Zong, Y., Fortinberry, H. & Weiss D.S. (2000). Regulation of recombinant gamma-aminobutyric acid (GABA)(A) and GABA(C) receptors by protein kinase. *C.Mol.Pharmacol.* 57, 847-856.
89. Flint, A.C., Dammerman, R.S. & Kriegstein, A. R. (1999). Endogenous activation of metabotropic glutamate receptors in neocortical development causes neuronal calcium oscillations. *Proc.Natl.Acad.Sci.U.S.A.* 96, 12144-12149.
90. Fritschy, J.M. & Brunig, I. (2003). Formation and plasticity of GABAergic synapses: physiological mechanisms and pathophysiological implications. *Pharmacol.Ther.* 98, 299-323.
91. Galante, M. & Diana, M.A.. (2004). Group I metabotropic glutamate receptors inhibit GABA release at interneuron-Purkinje cell synapses through endocannabinoid production. *J.Neurosci.* 24, 4865-4874.
92. Gao, L. & Greenfield, L.J. (2005). Activation of protein kinase C reduces benzodiazepine potency at GABAA receptors in NT2-N neurons. *Neuropharmacology.* 48, 333-342.
93. Gao, X.B. & van den Pol, A.N., (2000). GABA release from mouse axonal growth cones. *J.Physiol.* 523, 629-637.
94. Gasparini, F., Lingenhohl, K., Stoehr, N., Flor, P.J., Heinrich, M., Vranesic, I., Biollaz, M., Allgeier, H., Heckendorf, R., Urwyler, S., Varney, M.A., Johnson, E.C., Hess, S.D., Rao, S.P., Sacaan, A.I., Santori, E.M., Velicelebi, G. & Kuhn, R. (1999). 2-Methyl-6-(phenylethynyl)-pyridine (MPEP), a potent, selective and systemically active mGlu5 receptor antagonist. *Neuropharmacology.* 38, 1493-1503.
95. Geddis, M. S. & Rehder, V. (2003). The phosphorylation state of neuronal processes determines growth cone formation after neuronal injury. *J.Neurosci.Res.* 74, 210-220.

96. Gereau, R.W. & Conn, P.J. (1995). Multiple presynaptic metabotropic glutamate receptors modulate excitatory and inhibitory synaptic transmission in hippocampal area CA1. *J.Neurosci.* 15, 6879-6889.
97. Grantyn, R. (1988). Gaze control through superior colliculus: Structure and function. *Anatomy of the oculomotor system*. Vol.2, Buettner-Ennever, J., Ed., Elsevier, Amsterdam, 273-331
98. Grantyn, R., Jüttner R. & Meier, J. (2004). Development and use-dependent modification of synaptic connections in the visual layers of the rodent superior colliculus. *The superior colliculus – New approaches for studying sensorimotor integration*. Ch. 8, Hall, W.C. & Moschovakis, A., Ed., CRC Press, 173-210.
99. Grunbaum, L. & Muller, U. (1998). Induction of a specific olfactory memory leads to a long-lasting activation of protein kinase C in the antennal lobe of the honeybee. *J.Neurosci.* 18, 4384-4392.
100. Gryniewicz, G., Poeni, M. & Tsin, R.Y. (1985). A new generation of  $\text{Ca}^{2+}$  indicators with greatly improved fluorescence properties. *J.Biol.Chem.* 6, 3440-3450.
101. Hannan, A.J., Blakemore, C., Katsnelson, A., Vitalis, T., Huber, K.M., Bear, M., Roder, J., Kim, D., Shin, H.S. & Kind, P.C. (2001). PLC-beta1, activated via mGluRs, mediates activity-dependent differentiation in cerebral cortex. *Nat.Neurosci.* 4, 282-288.
102. Hanson, J.E. & Smith, Y. (1999) Group I metabotropic glutamate receptors at GABAergic synapses in monkeys. *J.Neurosci.* 19, 6488-6496.
103. Harting, J.K., Updyke, B.V. & Van Lieshout, D.P. (1992). Corticotectal projections in the cat: anterograde transport studies of twenty-five cortical areas. *J.Comp.Neurol.* 324(3), 379-414.
104. Henneberger, C., Jüttner, R., Rothe, T. & Grantyn, R. (2002). Postsynaptic action of BDNF on GABAergic synaptic transmission in the superficial layers of the mouse superior colliculus. *J.Neurophysiol.* 88, 595–603.
105. Henneberger, C., Kirischuk, S. & Grantyn, R. (2005). Brain-derived neurotrophic factor modulates GABAergic synaptic transmission by enhancing presynaptic glutamic acid decarboxylase 65 levels, promoting asynchronous release and reducing the number of activated postsynaptic receptors. *Neuroscience*. 135, 749-763.

- 106.Hennou, S., Khalilov, I., Diabira, D., Ben-Ari, Y. & Gozlan, H. (2002). Early sequential formation of functional GABA(A) and glutamatergic synapses on CA1 interneurons of the rat foetal hippocampus. *Eur.J.Neurosci.* 16, 197-208.
- 107.Hilbig, H. (1991). The structure of the superior colliculus in the microphthalmic mouse strain 944. *J.Hirnforsch.* 32, 135-138.
- 108.Hobi, R., Studer, M., Ruch, F. & Kuenzle, C.C. (1984). The DNA content of cerebral cortex neurons. Determinations by cytophotometry and high performance liquid chromatography. *Brain Res.* 305, 209-219.
- 109.Hoffpauir, B. K. & Gleason, E. L. (2002). Activation of mGluR5 modulates GABA(A) receptor function in retinal amacrine cells. *J.Neurophysiol.* 88, 1766-1776.
- 110.Holmes, K. H., Keele, N. B. & Shinnick-Gallagher, P. (1996) Loss of mGluR-mediated hyperpolarizations and increase in mGluR depolarizations in basolateral amygdala neurons in kindling-induced epilepsy. *J.Neurophysiol.* 76, 2808-2812.
- 111.Houser, C.R., Hendry, S.H., Jones, E.G. & Vaughn, J.E. (1983). Morphological diversity of immunocytochemically identified GABA neurons in the monkey sensory-motor cortex. *J.Neurocytol.* 12, 617-638.
- 112.Huang, L.Q., Rowan, M.J. & Anwyl, R. (1999). Role of protein kinases A and C in the induction of mGluR-dependent long-term depression in the medial perforant path of the rat dentate gyrus in vitro. *Neurosci.Lett.* 274, 71-74.
- 113.Hubert, G.W., Paquet, M. & Smith, Y. (2001). Differential subcellular localization of mGluR1a and mGluR5 in the rat and monkey Substantia nigra. *J.Neurosci.* 21, 1838-1847.
- 114.Isa, T., Endo, T. & Saito, Y., (1998). The visuo-motor pathway in the local circuit of the rat superior colliculus. *J.Neurosci.* 18, 8496-8504.
- 115.Isa, T. & Y. Saito. (2001). The direct visuo-motor pathway in mammalian superior colliculus; novel perspective on the interlaminar connection. *Neurosci.Res.* 41(2), 107–113.
- 116.Jokel, E.S., Garduno, E.R., Ariano, M.A. & Levine, M.S. (2001). Metabotropic glutamate receptors mGluR1alpha and mGluR2/3 display dynamic expression patterns in developing rat striatum. *Dev.Neurosci.* 23, 1-6.
- 117.Jourdain, P., Nikonenko, I., Alberi, S. & Muller, (2002). Remodeling of hippocampal synaptic networks by a brief anoxia-hypoglycemia. *J.Neurosci.* 22, 3108-3116.

- 118.Jouvenceau, A., Dutar, P. & Billard, J.M. (1995) Presynaptic depression of inhibitory postsynaptic potentials by metabotropic glutamate receptors in rat hippocampal CA1 pyramidal cells. *Eur. J. Pharmac.* 281, 131-139.
- 119.Jovanovic, J.N., Benfenati, F., Siow, Y.L., Sihra, T.S., Sanghera, J.S., Pelech, S.L., Greengard, P. & Czernik, A.J, (1996) Neurotrophins stimulate phosphorylation of synapsin I by MAP kinase and regulate synapsin I-actin interactions. *Proc.Natl.Acad.Sci USA* 93, 3679–3683.
- 120.Jovanovic, J.N., Sihra, T.S., Nairn, A.C., Hemmings, H.C. Jr., Greengard, P. & Czernik, A.J. (2001). Opposing changes in phosphorylation of specific sites in synapsin I during Ca<sup>2+</sup>-dependent glutamate release in isolated nerve terminals. *J.Neurosci.* 21, 7944-7953
- 121.Jovanovic, J.N., Thomas, P., Kittler, J.T., Smart, T.G., & Moss, S.J. (2004). Brain-derived neurotrophic factor modulates fast synaptic inhibition by regulating GABA(A) receptor phosphorylation, activity, and cell-surface stability. *J.Neurosci.* 24, 522-530.
- 122.Jüttner, R., Henneberger, C., Grantyn R. & Rothe T. (2001a). Early onset of glutamatergic and GABAergic synaptic activity in the visual layers of the rodent superior colliculus. *Int.J.Devl.Neurosci.* 19(3), 255–261.
- 123.Jüttner, R., Meier, J. & Grantyn, R.. (2001 b). Slow IPSC kinetics, low levels of α1 subunit expression and paired-pulse depression are distinct properties of neonatal inhibitory GABAergic synaptic connections in the mouse superior colliculus. *Eur.J.Neurosci.*, 13(11), 2088–2098.
- 124.Kapur, A., Yeckel, M. & Johnston, D. (2001). Hippocampal mossy fiber activity evokes Ca<sup>2+</sup> release in CA3 pyramidal neurons via a metabotropic glutamate receptor pathway. *Neuroscience* 107, 59-69.
- 125.Katchman, A.N. & Hershkowitz, N. (1997). Nitric oxide modulates synaptic glutamate release during anoxia. *Neurosci.Lett.* 228, 50-54.
- 126.Kawabata, S., Tsutsumi, R., Kohara, A., Yamaguchi, T., Nakanishi, S. & Okada, M. (1996). Control of calcium oscillations by phosphorylation of metabotropic glutamate receptors. *Nature* 383, 89-92.
- 127.Kerner, J.A., Standaert, D.G., Penney, J.B. Jr, Young, A.B. & Landwehrmeyer, G.B. (1997). Expression of group one metabotropic glutamate receptor subunit mRNAs in

- neurochemically identified neurons in the rat neostriatum, neocortex, and hippocampus. *Brain.Res.Mol.Brain.Res.* 48, 259–269.
- 128.Kirischuk, S., Clements, J.D. & Grantyn, R. (2002). Presynaptic and postsynaptic mechanisms underlie paired pulse depression at single GABAergic boutons in rat collicular cultures. *J.Physiol.* 543, 99-116.
- 129.Kirov, S.A., Petrak, L.J., Fiala, J.C. & Harris, K.M. (2004). Dendritic spines disappear with chilling but proliferate excessively upon rewarming of mature hippocampus. *Neuroscience*. 127, 69-80.
- 130.Kirov, S.A., Sorra, K.E. & Harris, K.M., (1999). Slices have more synapses than perfusion-fixed hippocampus from both young and mature rats. *J.Neurosci.* 19, 2876–2886.
- 131.Kirsch, J. & Betz, H., (1998). Glycine-receptor activation is required for receptor clustering in spinal neurons. *Nature* 392, 717–720.
- 132.Kishimoto, A., Kajikawa, N., Shiota, M. & Nishizuka, Y. (1983). Proteolytic activation of calcium-activated, phospholipid-dependent protein kinase by calcium-dependent neutral protease. *J.Biol.Chem.* 258, 1156-1164.
- 133.Kishimoto, A., Mikawa, K., Hashimoto, K., Yasuda, I., Tanaka, S., Tominaga, M., Kuroda, T. & Nishizuka, Y. (1989). Limited proteolysis of protein kinase C subspecies by calcium-dependent neutral protease (calpain). *J.Biol.Chem.* 264, 4088-4092.
- 134.Kittler, J.T. & Moss, S.J. (2003). Modulation of GABA<sub>A</sub> receptor activity by phosphorylation and receptor trafficking: implications for the efficacy of synaptic inhibition. *Curr.Opin.Neurobiol.* 13, 341-347.
- 135.Kneussel, M., Brandstatter, J.H., Laube, B., Stahl, S., Muller, U. & Betz, H. (1999). Loss of postsynaptic GABA(A) receptor clustering in gephyrin-deficient mice. *J.Neurosci.* 19, 9289–9297.
- 136.Konur, S. & Yuste, R. (2004). Imaging the motility of dendritic protrusions and axon terminals: roles in axon sampling and synaptic competition. *Mol.Cell.Neurosci.* 27, 427-40.
- 137.Koponen, S., Kurkinen, K., Akerman, K. E., Mochly-Rosen, D., Chan, P. H. & Koistinaho, J. (2003). Prevention of NMDA-induced death of cortical neurons by inhibition of protein kinase C zeta. *J.Neurochem.* 86, 442-450.

- 138.Kraszewski, K. & Grantyn R. (1992). Unitary, quantal and miniature GABA-activated synaptic chloride currents in cultured neurons from the rat superior colliculus. *Neuroscience*. 47(3), 555–570.
- 139.Krishek, B.J., Xie, X., Blackstone, C., Huganir, R.L., Moss S.J. & Smart T.G. (1994). Regulation of GABA<sub>A</sub> receptor function by protein kinase C phosphorylation. *Neuron* 12, 1081–1095.
- 140.Kyrozis, A.& Reichling, D.B. (1995) Perforated-patch recording with gramicidin avoids artifactual changes in intracellular chloride concentration. *J.Neurosci.Methods* 57, 27–35
- 141.Lam, K., Sefton, A.J. & Bennet, M.R. (1982). Loss of axons from the optic nerve of the rat during early postnatal development. *Develop.Brain Res.* 3, 487–491.
- 142.Lauri, S.E., Lamsa, K., Pavlov, I., Riekki, R., Johnson, B.E., Molnar, E., Rauvala, H. & Taira, T. (2003). Activity blockade increases the number of functional synapses in the hippocampus of newborn rats. *Mol.Cell Neurosci.* 22, 107-117.
- 143.Law, R.M., Stafford, A. & Quick, M.W. (2000). Functional regulation of gamma-aminobutyric acid transporters by direct tyrosine phosphorylation. *J.Biol.Chem.* 275, 23986-23991.
- 144.Lee, P.H., Helms, M.C., Augustine, G.C. & Hall, W.C. (1997). Role of intrinsic synaptic circuitry in collicular sensorimotor integration. *Proc.Natl.Acad.Sci. USA* 94, 13299-13304.
- 145.Lehmann, A. & Jacobson, I. (1990). Ion Dependence and Receptor Mediation of Glutamate Toxicity in the Immature Rat Hippocampal Slice. *Eur.J.Neurosci.* 2, 620-628.
- 146.Lévi, S., Vannier, C. & Triller, A. (1998). Strychnine-sensitive stabilization of postsynaptic glycine receptor clusters. *J.Cell.Sci.* 111, 335–345.
- 147.Li, D., Field, P.M., Yoshioka, N. & Raisman, G. (1994). Axons regenerate with correct specificity in horizontal slice culture of the postnatal rat entorhino-hippocampal system. *Eur.J.Neurosci.* 6, 1026-1037.
- 148.Liu Y.-B., Disterhoft J.F. & Slater N.T. (1993) Activation of metabotropic glutamate receptors induces long-term depression of GABAergic inhibition in hippocampus. *J.Neurophysiol.* 69, 1000–1004.
- 149.Llano I. & Marty A. (1995) Presynaptic metabotropic glutamatergic regulation of inhibitory synapses in rat cerebellar slices. *J.Physiol. (Lond.)* 486, 163–176.
- 150.Lund, R.D. (1966). The occipitotectal pathway of the rat. *J.Anat* 100, 51–62.

- 151.Lund, R. D. & Lund, J. S. (1972). Development of synaptic patterns in the superior colliculus of the rat. *Brain Res.* 42, 1–20.
- 152.Mannaioni, G., Carla, V. & Moroni, F. (1996). Pharmacological characterization of metabotropic glutamate receptors potentiating NMDA responses in mouse cortical wedge preparations. *Br.J.Pharmacol.* 118,1530-1536.
- 153.Marino, M.J., Wittmann, M., Bradley, S.R., Hubert, G.W., Smith, Y. & Conn, P.J. (2001). Activation of group I metabotropic glutamate receptors produces a direct excitation and disinhibition of GABAergic projection neurons in the substantia nigra pars reticulata. *J.Neurosci.* 21, 7001-7012.
- 154.Martin, L.J., Blackstone, C.D., Huganir, R.L. & Price, D. L. (1992). Cellular localization of a metabotropic glutamate receptor in rat brain. *Neuron* 9, 259–270.
- 155.Masu, M., Tanabe, Y., Tsuchida, K., Shigemoto, R. & Nakanishi, S. (1991). Sequence and expression of a metabotropic glutamate receptor. *Nature* 349, 760–765.
- 156.McDonald, B.J., Amato, A., Connolly, C.N., Benke, D., Moss, S.J. & Smart, T.G. (1998). Adjacent phosphorylation sites on GABA<sub>A</sub> receptor beta subunits determine regulation by cAMP-dependent protein kinase. *Nat.Neurosci.* 1, 23-28.
- 157.McDonald, B. J. & Moss, S. J. (1997). Conserved phosphorylation of the intracellular domains of GABA<sub>A</sub> receptor β2 and β3 subunits by cAMP-dependent protein kinase, cGMP-dependent protein kinase protein kinase C and Ca<sup>2+</sup>/calmodulin type II-dependent protein kinase. *Neuropharmacology*, 36, 1377–1385.
- 158.McGraw, C.F. & McLaughlin, B.J. (1980) Fine structural studies of synaptogenesis in the superficial layers of the chick optic tectum. *J.Neurocytol.* 9, 79-93.
- 159.McGraw, C.F., McLaughlin, B.J. & Boykins, L.G. (1980). A freeze-fracture study of synaptic junction development in the superficial layers of the chick optic tectum. *J.Neurocytol.* 9, 95-106.
- 160.McNally, H.A. & Borgens, R.B. (2004). Three-dimensional imaging of living and dying neurons with atomic force microscopy. *J.Neurocytol.* 33, 251-258.
- 161.Mehta, A.K. & Ticku, M.K. (1999). An update on GABA<sub>A</sub> receptors. *Brain Res.Rev.* 29, 196–217.
- 162.Meier, J. (2003). The enigma of transmitter-selective receptor accumulation at developing inhibitory synapses. *Cell Tissue Res.* 311, 271-276.

- 163.Meier, J., Akyeli, J., Kirischuk, S. & Grantyn, R. (2003). GABA(A) receptor activity and PKC control inhibitory synaptogenesis in CNS tissue slices. *Mol.Cell Neurosci.* 23, 600-613.
- 164.Meier, J. & Grantyn, R. (2004). A gephyrin-related mechanism restraining glycine receptor anchoring at GABAergic synapses. *J.Neurosci.* 24, 1398-1405.
- 165.Meier, J., Jüttner, R., Kirischuk, S. & Grantyn, R. (2002). Synaptic anchoring of glycine receptors in developing collicular neurons under control of metabotropic glutamate receptor activity. *Mol.Cell Neurosci.* 21, 324-340.
- 166.Meredith, M.A. & Stein, B.E. (1996). Spatial determinants of multisensory integration in cat superior colliculus neurons. *J.Neurophysiol.* 75, 1843-1857.
- 167.Mize, R.R. (1992). The organization of GABAergic neurons in the mammalian superior colliculus. *Progress in Brain Research.*, 90, 219–248.
- 168.Morimoto, K., Fahnestock, M. & Racine, R.J. (2004). Kindling and status epilepticus models of epilepsy: rewiring the brain. *Prog.Neurobiol.* 73, 1-60.
- 169.Morishita, W., Kirov, S. A. & Alger, B. E. (1998). Evidence for metabotropic glutamate receptor activation in the induction of depolarization-induced suppression of inhibition in hippocampal CA1. *J.Neurosci.* 18, 4870-4882.
- 170.Moroni, F., Lombardi, G., Thomsen, C., Leonardi, P., Attucci, S., Peruginelli, F., Torregrossa, S.A., Pellegrini-Giampietro, D.E., Luneia, R. & Pellicciari, R. (1997). Pharmacological characterization of 1-aminoindan-1,5-dicarboxylic acid, a potent mGluR1 antagonist. *J.Pharmacol.Exp.Ther.* 281, 721-729.
- 171.Moss, S. J. & Smart, T. G. (2001). Constructing inhibitory synapses. *Nat.Rev.Neurosci.* 2, 240-250.
- 172.Mozhayeva, M.G., Sara, Y., Liu, X. & Kavalali, E.T. (2002). Development of vesicle pools during maturation of hippocampal synapses. *J.Neurosci.* 22, 654-665.
- 173.Nakanishi, S. (1992). Molecular diversity of glutamate receptors and implications for brain function. *Science* 258, 597–603.
- 174.Nakanishi, S., Masu, M., Bessho, Y., Nakajima, Y., Hayashi, Y. & Shigemoto, R. (1994) Molecular diversity of glutamate receptors and their physiological functions. *Exs* 71, 71-80.
- 175.Nakanishi, S., Nakajima, Y., Masu, M., Ueda, Y., Nakahara, K., Watanabe, D., Yamaguchi, S., Kawabata, S. & Okada, M. (1998) Glutamate receptors: brain function and signal transduction. *Brain Res. Brain Res. Rev.* 26, 230-235.

176. Netzeband, J.G., Parsons, K.L., Sweeney, D.D. & Gruol, D.L. (1997). Metabotropic glutamate receptor agonists alter neuronal excitability and Ca<sup>2+</sup> levels via the phospholipase C transduction pathway in cultured Purkinje neurons. *J.Neurophysiol.* 78, 63-75.
177. Niell, C.M., Meyer, M.P. & Smith, S.J. (2004). In vivo imaging of synapse formation on a growing dendritic arbor. *Nat.Neurosci.* 7, 254-60.
178. Nicoletti, F., Bruno, V., Copani, A., Casabona, G. & Knopfel, T. (1996). Metabotropic glutamate receptors: a new target for the therapy of neurodegenerative disorders? *Trends Neurosci.* 19, 267-271.
179. Nishizawa Y. (2001). Glutamate release and neuronal damage in ischemia. *Life Sci.* 69, 369-381. Review.
180. Numberger, M. & Draguhn A. (1996) Patch-Clamp Technik. *Labor im Fokus*, Spektrum Akademischer Verlag Heidelberg
181. Okada, Y. (1992). The distribution and function of gamma-aminobutyric acid (GABA in the superior colliculus. *Prog.Brain.Res.*, 90, 249-262.
182. Olsen, R.W., DeLorey, T.M., Gordey, M. & Kang, M.H. (1999). GABA receptor function and epilepsy. *Adv.Neurol.* 79, 499-510.
183. Ottersen O.P. & Landsend A.S. (1997) Organization of glutamate receptors at the synapse. *Eur.J.Neurosci.* 9, 2219-2224.
184. Owens, D.F., Boyce, L.H., Davis, M.B. & Kriegstein, A.R. (1996). Excitatory GABA responses in embryonic and neonatal cortical slices demonstrated by gramicidin perforated-patch recordings and calcium imaging. *J.Neurosci.* 16, 6414-6423.
185. Pagano, A., Ruegg, D., Litschig, S., Stoehr, N., Stierlin, C., Heinrich, M., Floersheim, P., Prezeau, L., Carroll, F., Pin, J.P., Cambria, A., Vranesic, I., Flor, P.J. & Gasparini, F., Kuhn, R. (2000). The non-competitive antagonists 2-methyl-6-(phenylethynyl)pyridine and 7-hydroxyiminocyclopropan[b]chromen-1a-carboxylic acid ethyl ester interact with overlapping binding pockets in the transmembrane region of group I metabotropic glutamate receptors. *J.Biol.Chem.* 275, 33750-33758.
186. Pellegrini-Giampietro, D.E. (2003). The distinct role of mGlu1 receptors in post-ischemic neuronal death. *Trends Pharmacol.Sci.* 24, 461-470.
187. van Pelt, J., van Ooyen, A. & Corner, M.A. (1996). Growth cone dynamics and activity-dependent processes in neuronal network development. *Prog.Brain.Res.* 108, 333-346.

- 188.Pieribone, V.A., Porton, B., Rendon, B., Feng, J., Greengard, P. & Kao, H.T. (2002). Expression of synapsin III in nerve terminals and neurogenic regions of the adult brain. *J.Comp.Neurol.* 454,105-114.
- 189.Pin J.-P. & Duvoisin R. (1994). The Metabotropic Glutamate Receptors: Structure and Functions. *Neuropharmacology* 34, 1-26.
- 190.Pisani, A., Calabresi, P., Tozzi, A., D'Angelo, V. & Bernardi, G. (1998). L-type Ca<sup>2+</sup> channel blockers attenuate electrical changes and Ca<sup>2+</sup> rise induced by oxygen/glucose deprivation in cortical neurons. *Stroke*. 29, 196-201.
- 191.Poisbeau, P., Cheney, M. C., Browning, M. D. & Mody, I. (1999). Modulation of synaptic GABA<sub>A</sub> receptor function by PKA and PKC in adult hippocampal neurons. *J.Neurosci.* 19, 674-683.
- 192.Poisik, O.V., Mannaioni, G., Traynelis, S., Smith, Y. & Conn, P.J. (2003). Distinct functional roles of the metabotropic glutamate receptors 1 and 5 in the rat globus pallidus. *J.Neurosci.* 23, 122-130.
- 193.van den Pol, A.N., Obrietan, K., Belousov, A.B., Yang, Y. & Heller, (1998). H.C. Early synaptogenesis in vitro: role of axon target distance. *J.Comp.Neurol.* 399, 541-560.
- 194.Popoli, P., Frank, C., Tebano, M.T., Potenza, R.L., Pintor, A., Domenici, M.R., Nazziccone, V., Pezzola, A. & Reggio, R. (2003). Modulation of glutamate release and excitotoxicity by adenosine A2A receptors. *Neurology*.61, 69-71.
- 195.Popoli, P., Pintor, A., Tebano, M.T., Frank, C., Pepponi, R., Nazziccone, V., Grieco, R., Pezzola, A., Reggio, R., Minghetti, L., De Berardinis, M.A., Martire, A., Potenza, R.L., Domenici, M.R. & Massotti, M. (2004). Neuroprotective effects of the mGlu5R antagonist MPEP towards quinolinic acid-induced striatal toxicity: involvement of pre- and post-synaptic mechanisms and lack of direct NMDA blocking activity. *J.Neurochem* 89, 1479-1489.
- 196.Quick, M.W., Hu, J., Wang, D. & Zhang, H.Y. (2004). Regulation of a gamma-aminobutyric acid transporter by reciprocal tyrosine and serine phosphorylation. *J.Biol.Chem.* 279, 15961-15967.
- 197.Richerson, G.B. & Messer, C. (1995). Effect of composition of experimental solutions on neuronal survival during rat brain slicing. *Exp.Neurol.* 131, 133-143.

- 198.Rivera, C., Voipio, J. & Kaila, K. (2005). Two developmental switches in GABAergic signalling: the K<sup>+</sup>-Cl<sup>-</sup> cotransporter KCC2 and carbonic anhydrase CAVII. *J.Physiol.* 562, 27-36. Erratum in: (2005). *J.Physiol.* 564, 953.
- 199.Rivera, C., Voipio, J., Thomas-Crusells, J., Li, H., Emri, Z., Sipila, S., Payne, J.A., Minichiello, L., Saarma, M. & Kaila, K. (2004). Mechanism of activity-dependent downregulation of the neuron-specific K-Cl cotransporter KCC2. *J.Neurosci.* 24, 4683-4691.
- 200.Romano, C., Sesma, M.A., McDonald, C.T., O'Malley, K., Van Den Pol, A.N. & Olney, J.W. (1995). Distribution of metabotropic glutamate receptor mGluR5 immunoreactivity in rat brain. *J.Comp.Neurol.* 355, 455-469.
- 201.Rothe, T., Bahring, R., Carroll, P. & Grantyn, R. (1999). Repetitive firing deficits and reduced sodium current density in retinal ganglion cells developing in the absence of BDNF. *J.Neurobiol.* 40, 407-419.
- 202.Rothe, T., Bigl, V. & Grantyn, R. (1994). Potentiating and depressant effects of metabotropic glutamate receptor agonists on high-voltageactivated calcium currents in cultured retinal ganglion cells from postnatal mice. *Pflugers Archiv* 426, 161-170.
- 203.Rumpel, S., Hatt, H. & Gottmann, K. (1998). Silent synapses in the developing rat visual cortex: evidence for postsynaptic expression of synaptic plasticity. *J.Neurosci.* 18, 8863-8874.
- 204.Rumpel, S., Kattenstroth, G. & Gottmann, K. (2004). Silent synapses in the immature visual cortex: layer-specific developmental regulation. *J.Neurophysiol.* 91, 1097-1101.
- 205.Salt, T.E & Turner, J.P. (1998). Functions of metabotropic glutamate receptors in sensory processing in the thalamus. in: F. Moroni, F. Nicoletti, Pellegrini-Giampietro Eds., *Metabotropic Glutamate Receptors and Brain Function*, Portland Press Proceedings, pp. 49-58.
- 206.Sanes, J.R. & Lichtman, J.W. (2001). Induction, assembly, maturation and maintenance of a postsynaptic apparatus. *Nature Rev. Neurosci.* 2, 791-805.
- 207.Schauwecker, P.E. & McNeill, T.H. (1996). Dendritic remodeling of dentate granule cells following a combined entorhinal cortex/fimbria fornix lesion. *Exp.Neurol.* 141, 145-153.
- 208.Schiller, P.H., Sandell, J.H. & Maunsell, J.H. (1987) The effect of frontal eye field and superior colliculus lesions on saccadic latencies in the rhesus monkey. *J.Neurophysiol.* 57, 1033-1049.

- 209.Schmidt, S.A., Jüttner, R., Meier, J., Rothe, T. & Grantyn, R. (2002). NMDA receptor activation accelerates the decay of GABAergic IPSCs and increases the expression of the GABA<sub>A</sub> receptor subunit α1 in developing neurons of the rodent superior colliculus. In *FENS Abstr.* Vol 1, A011.13.
- 210.Schoepp, D.D. & Conn, P.J. (1993). Metabotropic glutamate receptors in brain function and pathology. *Trends in Pharmacol. Science* 14, 13–20.
- 211.Schoepp, D. D., Jane, D. E. & Monn, J. A. (1999). Pharmacological agents acting at subtypes of metabotropic glutamate receptors. *Neuropharmacology* 38, 1431-1476.
- 212.Schurr, A., Payne, R.S., Heine, M.F. & Rigor, B.M. (1995). Hypoxia, excitotoxicity, and neuroprotection in the hippocampal slice preparation. *J.Neurosci Methods*. 59,129-38.
- 213.Schwartz, N.E. & Alford, S. (2000). Physiological activation of presynaptic metabotropic glutamate receptors increases intracellular calcium and glutamate release. *J.Neurophysiol.* 84, 415-427.
- 214.Schwartz-Bloom, R.D. & Sah, R. (2001). gamma-Aminobutyric acid(A) neurotransmission and cerebral ischemia. *J.Neurochem.* 77, 353-71. Review.
- 215.Segal, M. (2001). Rapid plasticity of dendritic spine: hints to possible functions? *Prog.Neurobiol.* 63, 61-70.
- 216.Shigemoto, R., Kinoshita, A., Wada, E., Nomura, S., Ohishi, H., Takada, M., Flor, P.J., Neki, A., Abe, T., Nakanishi, S. & Mizuno, N. (1997). Differential presynaptic localization of metabotropic glutamate receptor subtypes in the rat hippocampus. *J.Neurosci.* 17, 7503-7522.
- 217.Shigemoto, R., Nakanishi, S. & Mizuno, N. (1992) Distribution of the mRNA for a metabotropic glutamate receptor (mGluR1) in the central nervous system: an in situ hybridization study in adult and developing rat. *J.Comp.Neurol.* 322, 121–135.
- 218.Shigemoto, R., Nomura, S., Ohishi, H., Sugihara, H., Nakanishi, S. & Mizuno, N. (1993) Immunohistochemical localization of a metabotropic glutamate receptor, mGluR5, in the rat brain. *Neurosci.Lett.* 163, 53–57.
- 219.Simburger, E., Plaschke, M., Fritschy, J.M. & Nitsch, R. (2001). Localization of two major GABA(A) receptor subunits in the dentate gyrus of the rat and cell type-specific up-regulation following entorhinal cortex lesion. *Neuroscience*. 102, 789-803.

- 220.Simburger, E., Plaschke, M., Kirsch, J. & Nitsch, R. (2000). Distribution of the receptor-anchoring protein gephyrin in the rat dentate gyrus and changes following entorhinal cortex lesion. *Cereb,Cortex.* 10, 422-432.
- 221.Siminoff, R., Schwassmann, H.O. & Kruger, L. (1966), An electrophysiological study of the visual projection to the superior colliculus of the rat. *J.Comp.Neurol.* 127, 435-4444.
- 222.Spooren W., Ballard T., Gasparini F., Amalric M., Mutel V. & Schreiber R. (2003) Insight into the function of Group I and Group II metabotropic glutamate (mGlu) receptors: behavioural characterization and implications for the treatment of CNS disorders. *Behav.Pharmacol* 14, 257-277.
- 223.Stein, B.E. & Meredith, M.A. (1993) The merging of the senses. *MIT Press*, Cambridge, MA.
- 224.Stefani, A., Pisano, A., Mercuri, N., Bernadi, G. & Calabresi, P. (1994). Activation of metabotropic glutamate receptors inhibits calcium currents and GABA-mediated synaptic potentials in striatal neurons. *J.Neurosci.* 14, 6734–6743.
- 225.Stone, L.M., Browning, M.D. 6 Finger, T.E. (1994). Differential distribution of the synapsins in the rat olfactory bulb. *J.Neurosci.* 14, 301-309.
- 226.Strasser, U., Lobner, D., Behrens, M. M., Canzoniero, L. M. & Choi, D. W. (1998). Antagonists for group I mGluRs attenuate excitotoxic neuronal death in cortical cultures. *Eur.J.Neurosci.* 10, 2848-2855.
- 227.Sugiyama, S. & Nakamura, H. (2003). The role of Grg4 in tectal laminar formation. *Development* 130, 451-462.
- 228.Swope, S.L., Moss, S.J., Raymond, L.A. & Huganir, R.L. (1999). Regulation of ligand-gated ion channels by protein phosphorylation. *Adv.Second.Messenger Phosphoprotein Res.* 33 49-78.
- 229.Takayama, C. & Inoue, Y. (2004). Extrasynaptic localization of GABA in the developing mouse cerebellum. *Neurosci.Res.* 50, 447-458.
- 230.Tallaksen-Greene, S.J., Kaatz, K.W., Romano, C. & Albin, R.L. (1998). Localization of mGluR1a-like immunoreactivity and mGluR5-like immunoreactivity in identified populations of striatal neurons. *Brain.Res.* 780, 210–217.
- 231.Tapia, J.C., Espinoza, F. & Aguayo, L.G. (1997). Differential intracellular regulation of cortical GABA(A) and spinal glycine receptors in cultured neurons. *Brain.Res.* 769, 203-210.

- 232.Tartaglia, N., Du, J., Tyler, W. J., Neale, E., Pozzo-Miller, L. D. & Lu, B. (2001). Protein synthesis-dependent and -independent regulation of hippocampal synapses by brain-derived neurotrophic factor. *J.Biol.Chem.*, 276, 37585–37593.
- 233.Testa, C.M., Standaert, D.G., Young, A.B. & Penney, J.B. Jr. (1994). Metabotropic glutamate receptors mRNA expression in the basal ganglia of the rat. *J.Neurosci.* 14, 3005–3018.
- 234.Tighilet, B. & Lacour, M. (2001). Gamma amino butyric acid (GABA) immunoreactivity in the vestibular nuclei of normal and unilateral vestibular neurectomized cats. *Eur.J.Neurosci.* 13, 2255–2267.
- 235.Tong, J.X., Vogelbaum, M.A., Drzymala, R.E. & Rich, K.M. (1997). Radiation-induced apoptosis in dorsal root ganglion neurons. *J.Neurocytol.* 26, 771-777.
- 236.Tremblay, R., Chakravarthy, B., Hewitt, K., Tauskela, J., Morley, P., Atkinson, T. & Durkin, J. P. (2000). Transient NMDA receptor inactivation provides long-term protection to cultured cortical neurons from a variety of death signals. *J.Neurosci.* 20, 7183-7192.
- 237.Tremblay, R., Hewitt, K., Lesiuk, H., Mealing, G., Morley, P. & Durkin, J. P. (1999). Evidence that brain-derived neurotrophic factor neuroprotection is linked to its ability to reverse the NMDA-induced inactivation of protein kinase C in cortical neurons. *J.Neurochem.* 72, 102-111.
- 238.Vajda, F.J. (2002). Neuroprotection and neurodegenerative disease. *J.Clin.Neurosci.* 9, 4-8.
- 239.Veng, L.M., Mesches, M.H. & Browning, M.D. (2003). Age-related working memory impairment is correlated with increases in the L-type calcium channel protein alpha1D (Cav1.3) in area CA1 of the hippocampus and both are ameliorated by chronic nimodipine treatment. *Brain.Res.Mol.Brain.Res.* 110, 193-202.
- 240.Waites, C.L., Craig, A.M. & Garner, C.C. (2005). Mechanisms of vertebrate synaptogenesis. *Annu.Rev.Neurosci.* 28, 251-274.
- 241.Wallace, M.T. & Stein, B.E. (1996). Sensory organization of the superior colliculus in cat and monkey. *Prog. Brain. Res.* 112, 301-311.
- 242.Wang, J. & Johnson, K.M. (1995) Regulation of striatal cyclic-3',5'-adenosine monophosphate accumulation and GABA release by glutamate metabotropic and dopamine D1 receptors. *J.Pharmacol.Exp.Ther.* 275, 877–884.

- 243.Wang, J., Lonart, G. & Johnson, K.M. (1996) Glutamate receptor activation induces carrier mediated release of endogenous GABA from rat striatal slices. *J. Neural Transm.* 103, 31–43.
- 244.Warton, S.S., Perouansky, M. & Grantyn, R. (1990). Development of GABAergic synaptic connections in vivo and in cultures from the rat superior colliculus. *Dev.Brain Res.* 52(1), 95–111.
- 245.Wetsel, W.C., Khan, W.A., Merchenthaler, I., Rivera, H., Halpern, A.E., Phung, H.M., Negro-Vilar, A. & Hannun, Y.A. (1992). Tissue and cellular distribution of the extended family of protein kinase C isoenzymes. *J.Cell.Biol.* 117, 121-133.
- 246.Wiener, S.I. (1986). Laminar distribution and patchiness of cytochrome oxidase in mouse superior colliculus. *J.Comp.Neurol.* 244, 137-148.
- 247.Wisden, W., Laurie, D. J., Monyer, H. & Seburg, P. H. (1992). The distribution of 13 GABA<sub>A</sub> receptor subunit mRNAs in the rat brain. I. Telencephalon, diencephalon, mesencephalon. *J.Neurosci.* 12, 1040–1062.
- 248.Yavin, E., Inserte-Igual, J. & Gil, S. (1995). Ischemia-triggered translocation and inactivation of protein kinase C isoforms in the fetal brain. *J.Neurochem.* 65, 2594-2602.
- 249.Zepeda, A., Sengpiel, F., Guagnelli, M.A., Vaca, L. & Arias, C. (2004). Functional reorganization of visual cortex maps after ischemic lesions is accompanied by changes in expression of cytoskeletal proteins and NMDA and GABA(A) receptor subunits. *J.Neurosci.* 24, 1812-1821.
- 250.Zheng, F. & Johnson, S.W. (2003). Dual modulation of gabaergic transmission by metabotropic glutamate receptors in rat ventral tegmental area. *Neuroscience*. 119, 453-640.
- 251.Zhuo, F.-M. & Hablitz, J.J. (1997a). Metabotropic glutamate receptor enhancement of spontaneous IPSCs in neocortical interneurons, *J.Neurophysiol.* 78, 2287–2295.
- 252.Zhuo, F.-M. & Hablitz, J.J. (1997b). Direct excitation of cortical interneurons by metabotropic glutamate receptors, *Soc.Neurosci.Abstr.* 23, 1997 2025.
- 253.Ziemka-Nalecz, M., Zalewska, T., Zajac, H., & Domanska-Janik, K. (2003). Decrease of PKC precedes other cellular signs of calpain activation in area CA1 of the hippocampus after transient cerebral ischemia. *Neurochem.Int.* 42, 205-214.