

- Abdu F, Hicks GA, Hennig G, Allen JP, Grundy D. Somatostatin sst(2) receptors inhibit peristalsis in the rat and mouse jejunum. *Am J Physiol Gastrointest Liver Physiol* 2002;282:G624-33.
- Ahmann GB, Sachs DH, Hodes RJ. Requirement for an Ia-bearing accessory cell in Con A-induced T cell proliferation. *J Immunol* 1978;121(5):1981-9.
- Ait-Azzouzene D, Gendron MC, Houdayer M, Langkopf A, Burki K, Nemazee D, Kanellopoulos-Langevin C. Maternal B lymphocytes specific for paternal histocompatibility antigens are partially deleted during pregnancy. *J Immunol*. 1998;161:2677-83.
- Aktories K, Schmidt G, Just I. Rho GTPases as targets of bacterial protein toxins. *Biol Chem* 2000;381:421-6.
- Albuquerque FC Jr, Smith EH, Kellum JM. 5-HT induces cAMP production in crypt colonocytes at a 5-HT₄ receptor. *J Surg Res* 1998;77:137-40.
- Alfieri AB, Cubeddu LX. Treatment with para-chlorophenylalanine antagonises the emetic response and the serotonin-releasing actions of cisplatin in cancer patients. *Br J Cancer* 1995;71:629-32.
- Ameisen JC, Meade R, Askenase PW. A new interpretation of the involvement of serotonin in delayed-type hypersensitivity. Serotonin-2 receptor antagonists inhibit contact sensitivity by an effect on T cells. *J Immunol* 1989;142(9):3171-9.
- Anlauf M, Eissele R, Schafer MKH, Eiden LE, Arnold R, Pauser U, Kloppel G, Weihe E. Expression of the two isoforms of the vesicular monoamine transporter (Vmat1 and Vmat2) in the endocrine pancreas and pancreatic endocrine tumors. *J Histochem Cytochem* 2003;51:1027-40.
- Aune TM, Golden HW, McGrath KM. Inhibitors of serotonin synthesis and antagonists of serotonin 1A receptors inhibit T lymphocyte function in vitro and cell-mediated immunity in vivo. *J Immunol*. 1994 Jul 15;153(2):489-98.
- Aune TM, McGrath KM, Sarr T, Bombara MP, Kelley KA. Expression of 5HT_{1a} receptors on activated human T cells. Regulation of cyclic AMP levels and T cell proliferation by 5-hydroxytryptamine. *J Immunol* 1993;151(3):1175-83.
- Austin MC, O'Donnel SM. Regional distribution and cellular expression of tryptophan hydroxylase messenger RNA in postmortem human brainstem and pineal gland. *J Neurochem* 1999;72:2065-73.
- Baban B, Chandler P, McCool D, Marshall B, Munn DH, Mellor AL. Indoleamine 2,3-dioxygenase expression is restricted to fetal trophoblast giant cells during murine gestation and is maternal genome specific. *J Reprod Immunol* 2004;61(2):67-77.
- Baggiolini M, Dahinden CA. CC chemokines in allergic inflammation. *Immunol Today* 1994;15(3):127-33.
- Barnes NM, Sharp T. A review of central 5-HT receptors and their function. *Neuropharmacology* 1999;38:1083-1152.
- Barone FC, Deegan JF, Price WJ, Fowler PJ, Fondacaro JD, Ormsbee HS 3rd. Cold-restraint stress increases rat fecal pellet output and colonic transit. *Am J Physiol* 1990;258:G329-37.
- Bearcroft CP, Andre EA, Farthing MJ. In vivo effects of the 5-HT₃ antagonist alosetron on basal and cholera toxin-induced secretion in the human jejunum: a segmental perfusion study. *Aliment Pharmacol Ther* 1997;11(6):1109-14.
- Bellivier F, Leboyer M, Courtet P, Buresi C, Beaufile B, Samolyk D, Allilaire J-F, Feingold J, Mallet J, Malafosse A. Association between the tryptophan hydroxylase gene and manic-depressive illness. *Arch Gen Psychiatry* 1998;55:33-7.

V LITERATURVERZEICHNIS

- Bengel D, Murphy DL, Andrews AM, Wichems CH, Feltner D, Heils A, Mossner R, Westphal H, Lesch KP. Altered brain serotonin homeostasis and locomotor insensitivity to 3,4-methylenedioxymethamphetamine („Ecstasy“) in serotonin transporter-deficient mice. *Mol Pharmacol* 1998;53:649-55.
- Bennet JP, Snyder SH. Serotonin and lysergic acid diethylamide binding in rat brain membranes: relationship to postsynaptic serotonin receptors. *Mol Pharmacol* 1976;12:373-89.
- Berk BC, Alexander RW, Brock TA, Gimbrone MA Jr, Webb RC. Vasoconstriction: a new activity for platelet-derived growth factor. *Science* 1986;232(4746):87-90.
- Bertazzo A, Ragazzi E, Biasiolo M, Costa CVL, Allegri G. Enzyme activities involved in tryptophan metabolism along the kynurenine pathway in rabbits. *Biochim Biophys Acta*. 2001;1527:167-75.
- Beubler E, Horina G. 5-HT₂ and 5-HT₃ receptor subtypes mediate cholera toxin-induced intestinal fluid secretion in the rat. *Gastroenterology* 1990;99:83-9.
- Bian X, Ren J, DeVries M, Schnegelsberg B, Cockayne D, Ford A, Galligan J. Peristalsis is impaired in the small intestine of mice lacking the P2X₃ subunit. *J Physiol* 2003;551(1):309-22.
- Billingham RE, Brent L, Medawar PB. ‚Actively acquired tolerance‘ of foreign cells. *Nature* 1953;172:603-6.
- Birnboim HC, Doly J. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* 1979;7(6):1513-23.
- Bischoff SC, Krieger M, Brunner T, Rot A, von Tschanner V, Baggiolini M, Dahinden CA. RANTES and related chemokines activate human basophil granulocytes through different G protein-coupled receptors. *Eur J Immunol* 1993;23(3):761-7.
- Border WA, Noble NA. Transforming growth factor beta in tissue fibrosis. *N Engl J Med* 1994;331:1286-92.
- Boucek RJ, Alvarez TR. 5-hydroxytryptamine: a cytospecific growth stimulator of cultured fibroblasts. *Science*. 1970 Feb 6;167(919):898-9.
- Boullin DJ. Observations on the significance of 5-hydroxytryptamine in relation to the peristaltic reflex of the rat. *Br J Pharmacol* 1964;23:14-33.;23:14-33.
- Brindley LL, Sweet JM, Goetzl EJ. Stimulation of histamine release from human basophils by human platelet factor 4. *J Clin Invest* 1983;72(4):1218-23.
- Buchheit KH. Inhibition of cholera toxin-induced intestinal secretion by the 5-HT₃ receptor antagonist ICS 205-930. *Naunyn Schmiedebergs Arch Pharmacol* 1989;339(6):704-5.
- Bulbring E, Crema A. The release of 5-hydroxytryptamine in relation to pressure exerted on the intestinal mucosa. *J Physiol* 1959a;146:18-28.
- Bulbring E, Crema A. The action of 5-hydroxytryptamine, 5-hydroxytryptophan and reserpine on intestinal peristalsis in anaesthetized guinea-pigs. *J Physiol* 1959b;146(1):29-53.
- Cady SG, Sono M. 1-methyl-DL-tryptophan, β -(3-Benzofuranyl)-DL-alanine (the oxygen analog of tryptophan), and β -[3-benzo(b)thienyl]-DL-alanine (the sulfur analog of tryptophan) are competitive inhibitors for indoleamine 2,3-dioxygenase. *Arch Biochem Biophys*. 1991;291:326.
- Cameron JS. Platelets in glomerular disease. *Annu Rev Med*. 1984;35:175-80.
- Carabelli V, Giancippoli A, Baldelli P, Carbone E, Artalejo AR. Distinct potentiation of L-type currents and secretion by cAMP in rat chromaffin cells. *Biophys J* 2003;85:1326-37.

- Cases O, Seif I, Grimsby J, Gaspar P, Chen K, Pournin S, Muller U, Aguet M, Babinet C, Shih JC et al. Aggressive behavior and altered amounts of brain serotonin and norepinephrine in mice lacking MAOA. *Science* 1995;268:1763-3.
- Cash CD, Vayer P, Mandel P, Maitre M. Tryptophan 5-hydroxylase. Rapid purification from whole brain and production of a specific antiserum. *Eur J Biochem* 1985;149:239-45.
- Centurion D, Glusa E, Sanchez-Lopez A, Valdivia LF, Saxena PR, Villalon CM. 5-HT₇, but not 5-HT_{2B}, receptors mediate hypotension in vagosympathectomized rats. *Eur J Pharmacol* 2004;502:239-42.
- Cetin Y, Kuhn M, Kulaksiz H, Adermann K, Bargsten G, Grube D, Forssmann WG. Enterochromaffin cells of the digestive system: cellular source of guanylin, a guanylate cyclase-activating peptide. *Proc Natl Acad Sci U S A* 1994;91(8):2935-9.
- Citron BA, Gregory EJ, Steigerwalt DS, Qin F, Festoff BW. Regulation of the dual function tissue transglutaminase/G α_h during murine neuromuscular development: gene and enzyme isoform expression. *Neurochem Int* 2000;37:337-49.
- Choquet D, Korn H. Dual effects of serotonin on a voltage-gated conductance in lymphocytes. *Proc Natl Acad Sci U S A* 1988;85(12):4557-61.
- Chung YI, Park DH, Kim M, Baker H, Joh TH. Immunochemical characterization of brain and pineal tryptophan hydroxylase. *J Korean Med Sci* 2001;16:489-97.
- Clench MH, Pineiro-Carrero VM, Mathias JR. Migrating myoelectric complex demonstrated in four avian species. *Am J Physiol* 1989;256:G598-603.
- Cohen ML, Wittenauer LA. Relationship between serotonin and tryptamine receptors in the rat stomach fundus. *J Pharmacol Exp Ther* 1985;233(1):75-9.
- Compan V, Zhou M, Grailhe R, Gazzara RA, Martin R, Gingrich J, Dumuis A, Brunner D, Bockaert J, Hen R. Attenuated response to stress and novelty and hypersensitivity to seizures in 5-HT₄ receptor knock-out mice. *J Neurosci* 2004;24:412-9.
- Cooke HJ. "Enteric Tears": Chloride Secretion and Its Neural Regulation. *News Physiol Sci* 1998;13:269-274.
- Cooke HJ. Neurotransmitters in neuronal reflexes regulating intestinal secretion. *Ann N Y Acad Sci* 2000;915:77-80.
- Cooke HJ, Sidhu M, Wang YZ. Activation of 5-HT_{1P} receptors on submucosal afferents subsequently triggers VIP neurons and chloride secretion in the guinea-pig colon. *J Auton Nerv Syst* 1997;66:105-10.
- Costa M, Furness JB, Cuello AC, Verhofstad AAJ, Steinbusch HJW, Elde RP. Neurons with 5-hydroxytryptamine-like immunoreactivity in the enteric nervous system: their visualization and reactions to drug treatment. *Neurosci* 1982;7:351-63.
- Cote F, Thevenot E, Fligny C, Fromes Y, Darmon M, Ripoche AM, Bayard E, Hanoun N, Saurini F, Lechat P, Dandolo L, Hamon M, Mallet J, Vojdani G. Disruption of the nonneuronal tph1 gene demonstrates the importance of peripheral serotonin in cardiac function. *PNAS* 2003;100:13525-30.
- Craig DA, Clarke DE. Peristalsis evoked by 5-HT and renzapride: evidence for putative 5-HT₄ receptor activation. *Br J Pharmacol* 1991;102(3):563-4.
- Dabire H, Cherqui C, Safar M, Schmitt H. Haemodynamic aspects and serotonin. *Clin Physiol Biochem*. 1990;8:56-63.

V LITERATURVERZEICHNIS

- Dale GL, Friese P, Batar B, Hamilton SF, Reed GL, Jackson KW, Clementson KJ, Alberio L. Stimulated platelets use serotonin to enhance their retention of procoagulant proteins on the cell surface. *Nature* 2002;415:175-9.
- Dahlström A, Fuxe K. Evidence for the existence of monoamine-containing neurons in the central nervous system. I. Demonstration of monoamines in the cell bodies of brain stem neurons. *Acta Physiol Scand Suppl* 1964;232:1-55.
- D'Andrea G, Welch KM, Nagel-Leiby S, Grunfeld S, Joseph R. Platelet catecholamines in migraine. *Cephalalgia* 1989;9(1):3-5.
- Day M, Vane JR. An analysis of the direct and indirect actions of drugs on the isolated guinea pig ileum. *Br J Pharmacol* 1963;20:150-70.
- Derkach V, Surprenant A, North RA. 5-HT₃ receptors are membrane ion channels. *Nature* 1989;339(6227):706-9.
- Deuel TF, Senior RM, Chang D, Griffin GL, Henrikson RL, Kaiser ET. Platelet factor 4 is chemotactic for neutrophils and monocytes. *Proc Natl Acad Sci U S A* 1981;78(7):4584-7.
- Diacovo TG, Catalina MD, Siegelman MH, von Andrian UH. Circulating activated platelets reconstitute lymphocyte homing and immunity in L-selectin-deficient mice. *J Exp Med*. 1998 Jan 19;187(2):197-204.
- Doggrell SA. The role of 5-HT on the cardiovascular and renal systems and the clinical potential of 5-HT modulation. *Expert Opin Investig Drugs*. 2003 May;12(5):805-23.
- Doggrell SA, Chen YY. Responsiveness, affinity constants and receptor reserves for serotonin on aortae of aged normotensive and hypertensive rats. *J Pharm Pharmacol* 2001;53:1403-8.
- Donaldson DM, Tew JG. beta-Lysin of platelet origin. *Bacteriol Rev*. 1977;41(2):501-13.
- Doye A, Mettouchi A, Bossis G, Clement R, Buisson-Touati C, Flatau G, Gagnoux L, Piechaczyk M, Boquet P, Lemichez E. CNF1 exploits the ubiquitin-proteasome machinery to restrict Rho GTPase activation for bacterial host cell invasion. *Cell* 2002;111:553-64.
- Dukat M, Smith C, Herrick-Davis K, Teitler M, Glennon RA. Binding of tryptamine analogs at h5-HT_{1E} receptors: a structure-affinity investigation. *Bioorg Med Chem* 2004;12(10):2545-52.
- Dumas S, Darmon MC, Delort J, Mallet J. Differential control of tryptophan hydroxylase expression in raphe and in pineal gland: evidence for a role of translation efficiency. *J Neurosci Res* 1989;24:537-47.
- Durham PL, Russo AF. New insights into the molecular actions of serotonergic antimigraine drugs. *Pharmacol Ther* 2002;94:77-92.
- Eichler HG, Ford GA, Blaschke TF, Swislocki A, Hoffman BB. Responsiveness of superficial hand veins to phenylephrine in essential hypertension. Alpha adrenergic blockade during prazosin therapy. *J Clin Invest* 1989;83(1):108-12.
- Enck P, Merlin V, Erckenbrecht JF, Wienbeck M. Stress effects on gastrointestinal transit in the rat. *Gut* 1989;30:455-9.
- Erlebacher A. Why isn't the fetus rejected? *Curr Opin Immunol*. 2001;13:590-3.
- Erspamer V. *Arch Sci Biol Napoli* 1946;31:86.
- Erspamer V, Asero B. Identification of enteramine, the specific hormone of the enterochromaffin cell system, as 5-hydroxytryptamine. *Nature* 1952;169:800-1.

- Ferrara A, Zinner MJ, Jaffe BM. Intraluminal release of serotonin, substance P, and gastrin in the canine small intestine. *Dig Dis Sci* 1987;32:289-94.
- Fetkovska N, Amstein R, Ferracin F, Regenass M, Buhler FR, Pletscher A. 5-Hydroxytryptamine kinetics and activation of blood platelets in patients with essential hypertension. *Hypertension* 1990 Mar;15(3):267-73.
- Feyer PC, Stewart AL, Titlbach OJ. Aetiology and prevention of emesis induced by radiotherapy. *Support Care Cancer* 1998;6:253-60.
- Finocchiaro LME, Arzt ES, Fernandez-Castelo S, Criscuolo M, Finkielman S, Nahmod VE. Serotonin and melatonin synthesis in peripheral blood mononuclear cells: Stimulation by interferon-gamma as part of an immunomodulatory pathway. *J Interferon Res* 1988;8:705-15.
- Fiorica-Howells E, Hen R, Gingrich J, Li Z, Gershon MD. 5-HT(2A) receptors: location and functional analysis in intestines of wild-type and 5-HT(2A) knockout mice. *Am J Physiol Gastrointest Liver Physiol* 2002;282:G877-93.
- Fiorica-Howells E, Maroteaux L, Gershon MD. Serotonin and the 5-HT(2B) receptor in the development of enteric neurons. *J Neurosci* 2000;20(1):294-305.
- Fon EA, Pothos EN, Sun BC, Killeen N, Sulzer D, Edwards RH. Vesicular transport regulates monoamine storage and release but is not essential for amphetamine action. *Neuron* 1997;19:1271-83.
- Fujigaki S, Saito K, Sekikawa K, Tone S, Takikawa O, Fujii H, Wada H, Noma A, Seishima M. Lipopolysaccharide induction of indoleamine 2,3-dioxygenase is mediated dominantly by an IFN-gamma-independent mechanism. *Eur J Immunol* 2001;31:2313-8.
- Fujimiya M, Okumiya K, Kuwahara A. Immunelectron microscopic study of the luminal release of serotonin from rat enterochromaffin cells induced by high intraluminal pressure. *Histochem Cell Biol* 1997;108:105-13.
- Fujiwara M, Shibata M, Nomiya Y, Sugimoto T, Hirata F, Tokuyama T, Senoh S, Hayaishi O. Formation of 5-hydroxykynurenine and 5-hydroxykynurenamine from 5-hydroxytryptophan in rabbit small intestine. *Proc Natl Acad Sci U S A* 1979;76(3):1145-9.
- Fukami MH, Holmsen H, Ugurbil K. Histamine uptake in pig platelets and isolated dense granules. *Biochem Pharmacol* 1984;33(23):3869-74.
- Furness JB, Costa M. Neurons with 5-hydroxytryptamine-like immunoreactivity in the enteric nervous system: their projections in the guinea pig small intestine. *Neurosci* 1982;7:341-50.
- Furness JB, Kunze WA, Bertrand PP, Clerc N, Bornstein JC. Intrinsic primary afferent neurons of the intestine. *Prog Neurobiol* 1998;54:1-18.
- Fuxe K. Evidence for the existence of monoamine neurons in the central nervous system. IV. Distribution of monoamine nerve terminals in the nervous central system. *Aca Physiol Scand* 1965;247:39-85.
- Gaddum JH, Picarelli ZP. Two kinds of tryptamine receptor. *Br J Pharmacol Chemother* 1957;12:323-8.
- Galligan JJ, North RA. Opioid, 5-HT1A and alpha 2 receptors localized to subsets of guinea-pig myenteric neurons. *J Auton Nerv Syst* 1991;32(1):1-11.
- Galli SJ, Hammel I. Unequivocal delayed hypersensitivity in mast cell-deficient and beige mice. *Science*. 1984 Nov 9;226(4675):710-3.
- Gaspar P, Cases O, Maroteaux L. The developmental role of serotonin: news from mouse molecular genetics. *Nature Reviews* 2003;4:1002-12.

- Gershon MD. Review article: roles played by 5-hydroxytryptamine in the physiology of the bowel. *Aliment Pharmacol Ther* 1999;13 Suppl 2:15-30.
- Gershon MD, Drakontides AB, Ross LL. Serotonin: synthesis and release from the myenteric plexus of the mouse intestine. *Science* 1965;149:197-9.
- Gershon MD, Dreyfus CF, Pickel VM, Joh TH, Reis DJ. Serotonergic neurons in the peripheral nervous system: identification in gut by immunohistochemical localization of tryptophan hydroxylase. *Proc Natl Acad Sci U S A* 1977;74:3086-9.
- Gershon MD. Serotonin and its implication for the management of irritable bowel syndrome. *Rev Gastroenterol Disord* 2003;3 Suppl 2:S25-34.
- Goyal RK, Hirano I. The enteric nervous system. *N Engl J Med* 1996;334:1106-15.
- Grabbe S, Schwarz T. Immunoregulatory mechanisms involved in elicitation of allergic contact hypersensitivity. *Immunol Today* 1998;19:37-44.
- Grailhe R, Waeber C, Dulawa SC, Hornung JP, Zhuang X, Brunner D, Geyer MA, Hen R. Increased exploratory activity and altered response to LSD in mice lacking the 5-HT(5A) receptor. *Neuron* 1999;22:581-91.
- Green CB, Beharrese JC. Tryptophan hydroxylase expression is regulated by a circadian clock in *Xenopus laevis* retina. *J Neurochem* 1994;62:2420-8.
- Greenberg CS, Birckbichler PJ, Rice RH. Transglutaminases: multifunctional cross-linking enzymes that stabilize tissues. *FASEB* 1991;5:3071-7.
- Gregory RE, Ettinger DS. 5-HT₃ receptor antagonists for the prevention of chemotherapy-induced nausea and vomiting. A comparison of their pharmacology and clinical efficacy. *Drugs* 1998;55:173-89.
- Grider JR, Kueemmerle JF, Jin JG. 5-HT released by mucosal stimuli initiates peristalsis by activating 5-HT₄/5-HT_{1p} receptors on sensory CGRP neurons. *Am J Physiol* 1996;270 *Am J Physiol*. 1996 May;270(5 Pt 1):G778-82.:G778-82.
- Grider JR, Foxx-Orenstein AE, Jin JG. 5-Hydroxytryptamine₄ receptor agonists initiate the peristaltic reflex in human, rat, and guinea pig intestine. *Gastroenterology* 1998;115(2):370-80.
- Grimsby J, Toth M, Chen K, Kumazawa T, Klaidman L, Adams JD, Karoum F, Gal J, Shih JC. Increased stress response and beta-phenylethylamine in MAOB-deficient mice. *Nat Genet* 1997;17:206-10.
- Gripenberg J. Inhibition of reserpine, guanethidine and imipramine of the uptake of 5-hydroxytryptamine by rat peritoneal mast cells in vitro. *Acta Physiol Scand* 1976;96:407-16.
- Gupta SL, Carlin JM, Pyati P, Dai W, Pfefferkorn ER, Murphy Jr MJ. Antiparasitic and antiproliferative effects of indoleamine 2,3-dioxygenase enzyme expression in human fibroblasts. *Infect Immun*. 1994;62:2277-84.
- Haga K, Asano K, Fukuda T, Kobayakawa T. The function of 5-HT₃ receptors on colonic transit in rats. *Obes Res* 1995;3 Suppl 5:801S-10S.
- Haga N, Mizumoto A, Satoh M, Mochiki E, Mizusawa F, Ohshima K, Itoh Z. Role of endogenous 5-hydroxytryptamine in the regulation of gastric contractions by motilin in dogs. *Am J Physiol* 1996;270:G20-G28.
- Hakanson R, Alumets J, Sundler F. Classification of peptide-hormone-producing cells. *Lancet* 1978;1:997.
- Hamlin KE, Fischer FE. The synthesis of 5-hydroxytryptamine. *J Am Chem Soc* 1951;73:5007-8.

- Hanahan D, Jessee J, Bloom FR. Plasmid transformation of *Escherichia coli* and other bacteria. *Methods Enzymol.* 1991;204:63-113.
- Hansson SR, Mezey E, Hoffman BJ. Serotonin transporter messenger RNA expression in neural crest-derived structures and sensory pathways of the developing rat embryo. *Neuroscience* 1999;89:24-65.
- Hart RP, Yang R, Riley LA, Green TL. Post-transcriptional control of tryptophan hydroxylase gene expression in rat brain stem and pineal gland. *Mol Cell Neurisci* 1991;2:71-7.
- Harvey M, Shink E, Tremblay M, Gagne B, Raymond C, Labbe M, Walther DJ, Bader M, Barden N. Support for the involvement of TPH2 gene in affective disorders. *Mol Psychiatry.* 2004 Nov;9(11):980-1.
- Hasegawa H, Iida Y, Oguro K, Kojima M, Ichiyama A. Tryptophan hydroxylase activity in serotonin producing mast cells. Dependence on intracellular iron concentration manipulated by permeable chelators. *Adv Exp Med Biol* 1996;398:513-7.
- Hasegawa H, Yanagisawa M, Inoue F, Yanaihara N, Ichiyama A. Demonstration of non-neuronal tryptophan 5-mono-oxygenase in mouse intestinal mucosa. *Biochem J* 1987;248:501-9.
- Hayaishi O. Properties and function of indoleamine 2,3-dioxygenase. *J Biochem* 1976;79:13P-21P.
- Haycock JW, Kumer SC, Lewis DA, Vrana KE, Stockmeier CA. A monoclonal antibody to tryptophan hydroxylase: application and identification of the epitope. *J Neurisci Methods* 2002;114:205-12.
- Hedlund PB, Danielson PE, Thomas EA, Slanina K, Carson MJ, Sutcliffe JG. No hypothermic response to serotonin in 5-HT7 receptor knockout mice. *Proc Natl Acad Sci USA* 2003;100:1375-80.
- Hellstrand K, Hermodsson S. Serotonergic 5-HT1A receptors regulate a cell contact-mediated interaction between natural killer cells and monocytes. *Scand J Immunol* 1993 Jan;37:7-18.
- Holland JM. Serotonin deficiency and prolonged bleeding in beige mice. *Proc Soc Exp Biol Med* 1976;151:32-9.
- Holtje M, Winter S, Walther D, Pahner I, Hortnagl H, Ottersen OP, Bader M, Ahnert-Hilger G. The vesicular monoamine content regulates VMAT2 activity through Galphaq in mouse platelets. Evidence for autoregulation of vesicular transmitter uptake. *J Biol Chem* 2003;278(18):15850-8.
- Hommel JD, Sears RM, Georgescu D, Simmons DL, DiLeone RJ. Local gene knockdown in the brain using viral-mediated RNA interference. *Nature Medicine* 2003;9:1539-44.
- Houston DS, Vanhoutte PM. Serotonin and the vascular system. Role in health and disease, and implications for therapy. *Drugs.* 1986 Feb;31(2):149-63.
- Hoyer D, Hannon JP, Martin GR. Molecular, pharmacological and functional diversity of 5-HT receptors. *Pharmacology Biochemistry and Behavior* 2002;71:533-54.
- Hsu SC, Johansson KR, Donahue MJ. The bacterial flora of the intestine of *Ascaris suum* and 5-hydroxytryptamine production. *J Parasitol* 1986;72(4):545-9.
- Huang S, Hendriks W, Althage A, Hemmi S, Bluethmann H, Kamijo R, Vilcek J, Zinkernagel RM, Aguet M. Immune response in mice that lack the interferon-gamma receptor. *Science* 1993;259:1742-5.
- Huizinga JD, Ambrous K, Der-Silaphet T. Co-operation between neural and myogenic mechanisms in the control of distension-induced peristalsis in the mouse small intestine. *J Physiol* 1998;506:843-56.

- Huo Y, Schober A, Forlow SB, Smith DF, Hyman MC, Jung S, Littman DR, Weber C, Ley K. Circulating activated platelets exacerbate atherosclerosis in mice deficient in apolipoprotein E. *Nat Med* 2003;9:61-7.
- Idzko M, Panther E, Stratz C, Muller T, Bayer H, Zissel G, Durk T, Soricther S, Di Virgilio F, Geissler M, Fiebich B, Herouy Y, Elsner P, Norgauer J, Ferrari D. The serotonergic receptors of human dendritic cells: identification and coupling to cytokine release. *J Immunol* 2004;172(10):6011-9.
- Iken K, Chheng S, Fargin A, Goulet AC, Kouassi E. Serotonin upregulates mitogen-stimulated B lymphocyte proliferation through 5-HT_{1A} receptors. *Cell Immunol* 1995;163:1-9.
- Jackson JC, Walker RF, Brooks WH, Roszman TL. Specific uptake of serotonin by murine macrophages. *Life Sci* 1988;42:1641-50.
- Jacobs BL, Azmitia EC. Structure and function of the brain serotonin system. *Physiol Rev* 1992;72:165-229.
- Jaffre F, Callebort J, Sarre A, Etienne N, Nebigil CG, Launay JM, Maroteaux L, Monassier L. Involvement of the serotonin 5-HT_{2B} receptor in cardiac hypertrophy linked to sympathetic stimulation: control of interleukin-6, interleukin-1 β , and tumor necrosis factor- α cytokine production by ventricular fibroblasts. *Circulation*. 2004 Aug 24;110(8):969-74.
- Jin JG, Foxx-Orenstein AE, Grider JR. Propulsion in guinea pig colon induced by 5-hydroxytryptamine (HT) via 5-HT₄ and 5-HT₃ receptors. *J Pharmacol Exp Ther* 1999;288:93-7.
- Jung MJ. Substrates and inhibitors of aromatic amino acid decarboxylase. *Bioorg Chem* 1986;14:429-43.
- Kamimura S, Eguchi K, Yonezawa M, Sekiba K. Localization and developmental change of indoleamine 2,3-dioxygenase activity in the human placenta. *Acta Med Okayama*. 1991;45:135-9.
- Kanai Y, Segawa H, Miyamoto K, Uchino H, Takeda E, Endou H. Expression cloning and characterization of a transporter for large neutral amino acids activated by the heavy chain of 4F2 antigen (CD98). *J Biol Chem* 1998;273(37):23629-32.
- Kato H, Osajima A, Uezono Y, Okazaki M, Tsuda Y, Tanaka H, Oishi Y, Izumi F, Nakashima Y. Involvement of PDGF in pressure-induced mesangial cell proliferation through PKC and tyrosine kinase pathways. *Am J Physiol* 1999;277:F105-12.
- Kellum J, McCabe M, Schneier J, Donowitz M. Neural control of acid-induced serotonin release from rabbit duodenum. *Am J Physiol* 1983;245:G824-31.
- Kelly RB. Storage and release of neurotransmitters. *Cell* 199;72:43-52.
- Khan NA, Poisson JP. 5-HT₃ receptor-channels coupled with Na⁺ influx in human T cells: role in T cell activation. *J Neuroimmunol*. 1999 Sep 1;99(1):53-60.
- Khozai LI, Puchkov VF, Otellin VA. The effect of a serotonin deficiency on mammalian embryonic development. *Ontogenez* 1995;26:350-5.
- Kilpatrick GJ, Butler A, Burridge J, Oxford AW. 1-(m-chlorophenyl)-biguanide, a potent high affinity 5-HT₃ receptor agonist. *Eur J Pharmacol* 1990;182:193-7.
- Kim KS, Wessel TC, Stone DM, Carver CH, Joh TH, Park DH. Molecular cloning and characterization of cDNA encoding tryptophan hydroxylase from rat central serotonergic neurons. *Brain Res Mol Brain Res* 1991;9:277-83.
- Kirchgessner AL, Tamir H, Gershon MD. Identification and stimulation by serotonin of intrinsic sensory neurons of the submucosal plexus of the guinea pig gut: activity-induced expression of Fos immunoreactivity. *J Neurosci* 1992;12:25-49.

- Klinger MH. Platelets and inflammation. *Anat Embryol (Berl)* 1997;196(1):1-11.
- Kuna P, Reddigari SR, Schall TJ, Rucinski D, Viksman MY, Kaplan AP. RANTES, a monocyte and T lymphocyte chemotactic cytokine releases histamine from human basophils. *J Immunol* 1992;149(2):636-42.
- Kurian SS, Ferri GL, De Mey J, Polak JM. Immunocytochemistry of serotonin-containing nerves in the human gut. *Histochemistry* 1983;78:523-9.
- Kurz SM, Diebold SS, Hieronymus T, Gust TC, Bartunek P, Sachs M, Birchmeier W, Zenke M. The impact of c-met/scatter factor receptor on dendritic cell migration. *Eur J Immunol* 2002;32(7):1832-8.
- Kut JL, Young MR, Crayton JW, Wright MA, Young ME. Regulation of murine T-lymphocyte function by spleen cell-derived and exogenous serotonin. *Immunopharmacol Immunotoxicol* 1992;14(4):783-96.
- Laberge S, Cruikshank WW, Beer DJ, Center DM. Secretion of IL-16 (lymphocyte chemoattractant factor) from serotonin-stimulated CD8+ T cells in vitro. *J Immunol* 1996 Jan 1;156(1):310-5.
- Lais LT, Brody MJ. Mechanism of vascular hyperresponsiveness in the spontaneously hypertensive rat. *Circ Res* 1975;36:216-22.
- Lauder JM. Neurotransmitters as growth regulatory signals: role of receptors and second messengers. *Trends Neurosci* 1993;16:233-40.
- Launay G, Costa JL, Da Prada M, Launay JM. Estimation of rate constants for serotonin uptake and compartmentation in normal human platelets. *Am J Physiol* 1994;266:R1061-75.
- Lenschow DJ, Walunas TL, Bluestone JA. CD28/B7 system of T cell costimulation. *Annu Rev Immunol* 1996;14:233-58.
- Lesch KP, Bengel D, Heils A, Sabol SZ, Greenberg BD, Petri S, Benjamin J, Muller CR, Hamer DH, Murphy DL. Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science* 1996;274(5292):1527-31.
- Lewis GP. The inhibition by morphine of the action of smooth muscle stimulants on the guinea-pig intestine. *Br J Pharmacol Chemother* 1960;15:425-31.
- Liang J, Wessel JH 3rd, Iuvone PM, Tosini G, Fukuhara C. Diurnal rhythms of tryptophan hydroxylase 1 and 2 mRNA expression in the rat retina. *Neuroreport* 2004;15(9):1497-500.
- Lomax RB, Gallego S, Novalbos J, Garcia AG, Warhurst G. L-Type calcium channels in enterochromaffin cells from guinea pig and human duodenal crypts: an in situ study. *Gastroenterology* 1999;117(6):1363-9.
- Lucki I. The spectrum of behaviors influenced by serotonin. *Biol Psychiatry* 1998;44:151-62.
- MacKenzie CR, Langen R, Takikawa O, Daubener W. Inhibition of indoleamine 2,3-dioxygenase in human macrophages inhibits interferon-gamma-induced bacteriostasis but does not abrogate toxoplasmatosis. *Eur J Immunol* 1999;29(10):3254-61.
- Mann JJ, Brent DA, Arango V. The neurobiology and genetics of suicide and attempted suicide: a focus on the serotonergic system. *Neuropsychopharmacology* 2001;24:467-77.
- Martin M. The severity and pattern of emesis following different cytotoxic agents. *Oncology* 1996;53 Suppl 1:26-31.
- Masuda M, Betancourt L, Matsusawa T, Kashimoto T, Takao T, Shimonishi Y, Horiguchi Y. Activation of Rho through a cross-link with polyamines catalyzed by Bordetella dermonecrotizing toxin. *EMBO* 2000;19:521-30.

- Mathiau P, Riche D, Behzadi G, Dimitriadou V, Aubineau P. Absence of serotonergic innervation from raphe nuclei in rat cerebral blood vessels – I. Histological evidence. *Neuroscience* 1993;52:645-55.
- Mawe GM, Branchek TA, Gershon MD. Peripheral neural serotonin receptors: identification and characterization with specific antagonists and agonists. *Proc Natl Acad Sci U S A* 1986;83(24):9799-803.
- Mayes PA. Nutrition. In: Harper's Biochemistry, 24th ed. Appleton & Lange 1996.
- Meißner T. Pica-Syndrom: Haare oder Erde füllen den Magen. *Ärzte Zeitung* 2002; Onlineausgabe: www.aerztezeitung.de.
- Melino G, Annicchiarico-Petruzzelli M, Piredda L, Candi E, Gentile V, Davies PJ, Piacentini M. Tissue transglutaminase and apoptosis: sense and antisense transfection studies with human neuroblastoma cells. *Mol Cell Biol* 1994;14:6584-96.
- Mellor AL, Baban B, Chandler P, Marshall B, Jhaver K, Hansen A, Koni PA, Iwashima M, Munn DH. Cutting edge: induced indoleamine 2,3 dioxygenase expression in dendritic cell subsets suppresses T cell clonal expansion. *J Immunol.* 2003 Aug 15;171(4):1652-5.
- Mellor AL, Keskin DB, Johnson T, Chandler P, Munn DH. Cells expressing indoleamine 2,3-dioxygenase inhibit T cell responses. *J Immunol.* 2002;168:3771-6.
- Mellor AL, Sivakumar J, Chandler P, Smith K, Molina H, Mao D, Munn DH. Prevention of T cell-driven complement activation and inflammation by tryptophan catabolism during pregnancy. *Nat Immunol.* 2001;2:64-8.
- Meseguer J, Esteban MA, Rodriguez A. Are thrombocytes and platelets true phagocytes? *Microsc Res Tech* 2002;57(6):491-7.
- Meyniel JP, Khan NA, Ferriere F, Deschaux P. Identification of lymphocyte 5-HT₃ receptor subtype and its implication in fish T-cell proliferation. *Immunol Lett* 1997;55(3):151-60.
- Michel K, Sann H, Schaaf C, Schemann M. Subpopulations of gastric myenteric neurons are differentially activated via distinct serotonin receptors: projection, neurochemical coding, and functional implications. *J Neurosci* 1997;17(20):8009-17.
- Miki T, Sun H, Lee YH, Tandin A, Kovscek AM, Subbotin V, Fung JJ, Valdivia LA. Blockade of tryptophan catabolism prevents spontaneous tolerogenicity of liver allografts. *Transplant Proc.* 2001;33:129-3040
40.
- Missouris CG, Cappuccio FP, Varsamis E, Barron JL, Carr E, Markandu ND, MacGregor GA. Serotonin and heart rate in hypertensive and normotensive subjects. *Am Heart J* 1998;135:838-43.
- Monro RL, Bertrand PP, Bornstein JC. ATP and 5-HT are the principal neurotransmitters in the descending excitatory reflex pathway of the guinea-pig ileum. *Neurogastroenterol Mot* 2002;14:255-64.
- Mossner R, Lesch KP. Role of serotonin in the immune system and in neuroimmune interactions. *Brain Behav Immun.* 1998 Dec;12(4):249-71.
- Munn DH, Shafizadeh E, Attwood JT, Bondarev I, Pashine A, Mellor AL. Inhibition of T cell proliferation by macrophage tryptophan catabolism. *J Exp Med.* 1999;189:1363-72.
- Munn DH, Sharma MD, Lee JR, Jhaver KG, Johnson TS, Keskin DB, Marshall B, Chandler P, Antonia SJ, Burgess R, Slingluff Jr CL, Mellor AL. Potential regulatory function of human dendritic cells expressing indoleamine 2,3-dioxygenase. *Science.* 2002;297:1867-70.

V LITERATURVERZEICHNIS

- Munn DH, Zhou M, Attwood JT, Bondarev I, Conway SJ, Marshall B, Brown C, Mellor AL. Prevention of allogeneic fetal rejection by tryptophan catabolism. *Science*. 1998;281:1191-3.
- Murphy WJ, Taub DD, Anver M, Conlon K, Oppenheim JJ, Kelvin DJ, Longo DL. Human RANTES induces the migration of human T lymphocytes into the peripheral tissues of mice with severe combined immune deficiency. *Eur J Immunol* 1994;24(8):1823-7.
- Nagakura Y, Kontoh A, Tokita K, Tomoi M, Shimomura K, Kadowaki M. Combined blockade of 5-HT₃- and 5-HT₄-serotonin receptors inhibits colonic functions in conscious rats and mice. *J Pharmacol Exp Ther* 1997;281:284-90.
- Nagakura Y, Naitoh Y, Kamato T, Yamano M, Miyata K. Compounds possessing 5-HT₃ receptor antagonistic activity inhibit intestinal propulsion in mice. *Eur J Pharmacol* 1996;311:67-72.
- Nebigil CG, Choi DS, Dierich A, Hickel P, Le Meur M, Messaddeq N, Launay JM, Maroteaux L. Serotonin 2B receptor is required for heart development. *Proc Natl Acad Sci USA* 2000;97:9508-13.
- Nemecek GM, Coughlin SR, Handley DA, Moskowitz MA. Stimulation of aortic smooth muscle cell mitogenesis by serotonin. *Proc Natl Acad Sci U S A* 1986;83(3):674-8.
- Nieuwenhuijs VB, Verheem A, van Duijvenbode-Beumer H *et al*. The role of interdigestive small bowel motility in the regulation of gut microflora, bacterial overgrowth, and bacterial translocation in rats. *Ann Surg* 1998;228:188-93.
- Nilsson O, Cassuto J, Larsson PA, Jodal M, Lidberg P, Ahlman H, Dahlstrom A, Lundgren O. 5-Hydroxytryptamine and cholera secretion: a histochemical and physiological study in cats. *Gut* 1983;24:542-8.
- Ning Y, Zhu JX, Chan HC. Regulation of ion transport by 5-hydroxytryptamine in rat colon. *Clin Exp Pharmacol Physiol* 2004;31:424-8.
- Oda A, Miyakawa Y, Druker BJ, Ozaki K, Yabusaki K, Shirasawa Y, Handa M, Kato T, Miyazaki H, Shimosaka A, Ikeda Y. Thrombopoietin primes human platelet aggregation induced by shear stress and by multiple agonists. *Blood*. 1996 Jun 1;87(11):4664-70.
- Ormsbee III HS, Fondacaro JD. Action of serotonin in the gastrointestinal tract. *Proc Soc Biol Med* 1985;178:333-8.
- Osaka M, Fujita T, Yanatori Y. On the possible role of intestinal hormones as the diarrhoeagenic messenger in cholera. *Virchows Arch B Cell Pathol* 1975;18:287-96.
- Ozaki K, Fujimori H, Nomura S, Nishikawa T, Nishimura M, Pan Hou H, Narama I. Morphologic and hematologic characteristics of storage pool deficiency in beige rats (Chediak-Higashi syndrome of rats). *Lab Anim Sci* 1998;48(5):502-6.
- Pan H, Gershon MD. Activation of intrinsic afferent pathways in submucosal ganglia of the guinea pig small intestine. *J Neurosci* 2000;20(9):3295-309.
- Parks CL, Robinson PS, Sibille E, Shenk T, Toth M. Increased anxiety of mice lacking the serotonin_{1A} receptor. *Proc Natl Acad Sci USA* 1998;95:10734-9.
- Parrat JR, West GB. 5-hydroxytryptamine and tissue mast cells. *J Physiol* 1957;137:169-78.
- Patrono C, Pierucci A. The Use of Antiplatelet Agents in Glomerulonephritis: A Pharmacological Approach. *Nephrol Dial Transplant Suppl*. 1990;1:29-32.
- Peotta VA, Vasquez EC, Meyrelles SS. Cardiovascular neural reflexes in L-NAME-induced hypertension in mice. *Hypertension* 2001;38:555-9.

- Perkins A, Goy MF, Li Z. Uroguanylin is expressed by enterochromaffin cells in the rat gastrointestinal tract. *Gastroenterology* 1997;113(3):1007-14.
- Peroutka S, Snyder SH. Multiple serotonin receptors. Differential binding of [³H]-5-hydroxytryptamine, [³H]-lysergic acid diethylamide and [³H]-spiroperidol. *Mol Pharmacol* 1979;16:687-99.
- Peters LJ. Abdominal pregnancy in a golden hamster (*Mesocricetus auratus*). *Lab Anim Sci* 1982;32:392-3.
- Pfefferkorn ER. Interferon γ blocks the growth of *Toxoplasma gondii* in human fibroblasts by inducing the host cells to degrade tryptophan. *Proc Natl Acad Sci USA*. 1984;81:908-12.
- Piguet PF, Grau GE, Hauser C, Vassalli P. Tumor necrosis factor is a critical mediator in hapten induced irritant and contact hypersensitivity reactions. *J Exp Med* 1991;173(3):673-9.
- Ptak W, Herzog WR, Askenase PW. Delayed-type hypersensitivity initiation by early-acting cells that are antigen mismatched or MHC incompatible with late-acting, delayed-type hypersensitivity effector T cells. *J Immunol* 1991;146:469-75.
- Okuma M, Tokuyama T, Senoh S, Hirata F, Hayaishi O. Antagonism of 5-hydroxykynuren-amine against serotonin action on platelet aggregation. *Proc Natl Acad Sci U S A* 1976;73(2):643-5.
- Qian S, Demetris AJ, Murase N, Rao AS, Fung JJ, Starzl TE. Murine liver allograft transplantation: tolerance and donor cell chimerism. *Hepatology*. 1994;19:916-24.
- Racke K, Reimann A, Schworer H, Kilbinger H. Regulation of 5-HT release from enterochromaffin cells. *Behav Brain Res* 1996;73:83-7.
- Ralevic V, Khalil Z, Helme RD, Dusting GJ. Role of nitric oxide in the actions of substance P and other mediators of inflammation in rat skin microvasculature. *Eur J Pharmacol*. 1995 Sep 25;284(3):231-9.
- Ramage AG. Central cardiovascular regulation and 5-hydroxytryptamine receptors. *Brain Res Bull* 2001;56:425-39.
- Ramboz S, Saudou F, Amara DA, Belzung C, Segu L, Misslin R, Buhot MC, Hen R. 5-HT1B receptor knock out – behavioral consequences. *Behav Brain Res* 1996;7:305-12.
- Rapport MM. Serum vasoconstrictor (serotonin) V. The presence of creatinine in the complex: a proposed study of the vasoconstrictor principle. *J Biol Chem* 1949;180:961-9.
- Rapport MM, Green AA, Page IH. Purification of the substance which is responsible for the vasoconstrictor activity of serum. *Fed Proc* 1947;6:184.
- Rapport MM, Green AA, Page IH. Crystalline Serotonin. *Science* 1948;108:329-30.
- Rapport MM, Green AA, Page IH. Serum vasoconstrictor (serotonin). III. Chemical inactivation. *J Biol Chem* 1948;176:1237.
- Raymond JR, Mukhin YV, Gelasco A, Turner J, Collinsworth G, Gettys TW, Grewal JS, Garnovskaya MN. Multiplicity of mechanisms of serotonin receptor signal transduction. *Pharmacology & Therapeutics* 2001;92:179-212.
- Reading JP, Huffman JL, Wu JC, Palmer FT, Harton GL, Sisson ME, Keyvanfar K, Gresinger TH, Cochrane WJ, Fallon LA. Nucleated erythrocytes in maternal blood: quantity and quality of fetal cells in enriched populations. *Human Reprod*. 1995;10:2510-5.
- Reiter RJ. Pineal melatonin: cell biology of its synthesis and of its physiological interactions. *Endocr Rev* 1991;12:151-80.

- Reiter RJ, Tan DX, Mayo JC, Sainz RM, Leon J, Bandyopadhyay D. Neurally-mediated and neurally-independent beneficial actions of melatonin in the gastrointestinal tract. *J Physiol Pharmacol* 2003;54 Suppl 4:113-25.
- Resnick RH, Gray SJ. Chemical and histologic demonstration of hydrochloric acid-induced release of serotonin from intestinal mucosa. *Gastroenterology* 1962;42:48-55.
- Reuss S. Components and connections of the circadian timing system in mammals. *Cell Tissue Res* 1996;285:353-78.
- Robbiani DF, Finch RA, Jager D, Muller WA, Sartorelli AC, Randolph GJ. The leukotriene C(4) transporter MRP1 regulates CCL19 (MIP-3beta, ELC)-dependent mobilization of dendritic cells to lymph nodes. *Cell* 2000;103(5):757-68.
- Roberts AJ, Krucker T, Levy CL, Slanina KA, Sutcliffe JG, Hedlund PB. Mice lacking 5-HT receptors show specific impairments in contextual learning. *Eur J Neurosci.* 2004 Apr;19(7):1913-22.
- Rodriguez-Iturbe B, Pons H, Herrera-Acosta J, Johnson RJ. Role of immunocompetent cells in nonimmune renal diseases. *Kidney Int.* 2001 May;59(5):1626-40.
- Rot A, Krieger M, Brunner T, Bischoff SC, Schall TJ, Dahinden CA. RANTES and macrophage inflammatory protein 1 alpha induce the migration and activation of normal human eosinophil granulocytes. *J Exp Med* 1992;176(6):1489-95.
- Rubin W, Gershon MD, Ross LL. Electron microscope radioautographic identification of serotonin-synthesizing cells in the mouse gastric mucosa. *J Cell Biol* 1971;50:339-415.
- Sack DA, Sack RB, Nair GB, Siddique AK. Cholera. *Lancet.* 2004;363(9404):223-33.
- Sambrook J, Fritsch EF, Maniatis T. *Molecular cloning: a laboratory manual.* 2nd edition.
- Santucci S, Corazzi G, Francia N, Antonelli A, Aloe L, Alleva E. Neurobehavioural effects of hypergravity conditions in the adult mouse. *Neuroethology* 2000;11:3353-6.
- Saxena PR, Villalon CM. Cardiovascular effects of serotonin agonists and antagonists. *J Cardiovasc Pharmacol* 1990;7:S17-34.
- Schall TJ, Bacon K, Toy KJ, Goeddel DV. Selective attraction of monocytes and T lymphocytes of the memory phenotype by cytokine RANTES. *Nature* 1990;347(6294):669-71.
- Schmidt G, Sehr P, Wilm M, Selzer J, Mann M, Aktories K. Gln 63 of Rho is deamidated by *Escherichia coli* cytotoxic necrotizing factor-1. *Nature* 1997;387:725-9.
- Schmidt G, Goehring UM, Schirmer J, Lerm M, Aktories K. Identification of the C-terminal part of Bordetella dermonecrotic toxin as a transglutaminase for Rho GTPases. *J Biol Chem* 1999;274:31875-81.
- Schworer H, Racke K, Kilbinger H. Cisplatin increases the release of 5-hydroxytryptamine (5-HT) from the isolated vascularly perfused small intestine of the guinea-pig: involvement of 5-HT₃ receptors. *Naunyn Schmiedebergs Arch Pharmacol* 1991;344:143-9.
- Sears CL, Kaper JB. Enteric bacterial toxins: mechanisms of action and linkage to intestinal secretion. *Microbiol Rev* 1996;60:167-215.
- Segawa H, Fukasawa Y, Miyamoto K, Takeda E, Endou H, Kanai Y. Identification and functional characterization of a Na⁺-independent neutral amino acid transporter with broad substrate selectivity. *J Biol Chem* 1999;274(28):19745-51.

- Shimizu T, Abe R, Nishihira J, Shibaki A, Watanabe H, Nakayama T, Taniguchi M, Ishibashi T, Shimizu H. Impaired contact hypersensitivity in macrophage migration inhibitory factor-deficient mice. *Eur J Immunol.* 2003 Jun;33(6):1478-87.
- Shimizu T, Nomiya S, Hirata F, Hayaishi O. Indoleamine 2,3-Dioxygenase. *J Biol Chem* 1978;253:4700-6.
- Shultz PJ, DiCorleto PE, Silver BJ, Abboud HE. Mesangial cells express PDGF mRNAs and proliferate in response to PDGF. *Am J Physiol* 1988;255:F674-84.
- Sibella-Arguelles C. The proliferation of human T lymphoblastic cells induced by 5-HT_{1B} receptors activation is regulated by 5-HT-moduline. *C R Acad Sci III.* 2001 Apr;324(4):365-72.
- Simons FE. H₁-Antihistamines: more relevant than ever in the treatment of allergic disorders. *J Allergy Clin Immunol* 2003;112:S42-52.
- Singh RA, Sodhi A. Antigen presentation by cisplatin-activated macrophages: role of soluble factor(s) and second messengers. *Immunol Cell Biol* 1998;76:513-9.
- Smith CC. Evidence for separate serotonin and catecholamine compartments in human platelets. *Biochim Biophys Acta* 1996;1291(1):1-4.
- Soslau G, Silverberg M, Brodsky I, McCarty PP. ATP modification of serotonin-induced contraction of the rat pulmonary artery. *Proc Soc Exp Biol Med* 1997;214:233-41.
- Spangue A, Forsgren S, El-Salhy M. Effect of diabetic state on co-localization of substance P and serotonin in the gut in animal models. *Histol Histopathol* 2001;16(2):393-8.
- Spangler BD. Structure and function of cholera toxin and the related *Escherichia coli* heat-labile enterotoxin. *Microbiol Rev* 1992;56:622-47.
- Staite ND, Justen JM, Sly LM, Beaudet AL, Bullard DC. Inhibition of delayed-type contact hypersensitivity in mice deficient in both E-selectin and P-selectin. *Blood* 1996;88(8):2973-9.
- Stefulj J, Cicin-Sain L, Schauenstein K, Jernej B. Serotonin and immune response: effect of the amine on in vitro proliferation of rat lymphocytes. *Neuroimmunomodulation.* 2001;9(2):103-8.
- Stefulj J, Jakopcic S, Osmak M, Jernej B. Serotonin and apoptosis: studies on rat lymphocytes. *Neuroimmunomodulation.* 2002-2003;10(3):132-3.
- Stefulj J, Jernej B, Cicin-Sain L, Rinner I, Schauenstein K. mRNA expression of serotonin receptors in cells of the immune tissues of the rat. *Brain Behav Immun* 2000;14:219-24.
- Sternberg EM, Wedner HJ, Leung MK, Parker CW. Effect of serotonin (5-HT) and other monoamines on murine macrophages: modulation of interferon-gamma induced phagocytosis. *J Immunol* 1987;138(12):4360-5.
- Sugden D. Comparison of circadian expression of tryptophan hydroxylase isoform mRNAs in the rat pineal gland using real-time PCR. *J Neurochem.* 2003;86(5):1308-11.
- Szasz R, Dale GL. Thrombospondin and fibrinogen bind serotonin-derivatized proteins on COAT-platelets. *Blood* 2002;100:2827-1.
- Szczepanik M, Anderson LR, Ushio H, Ptak W, Owen MJ, Hayday AC, Askenase PW. $\gamma\delta$ T cells from tolerized $\alpha\beta$ T cell receptor (TCR)-deficient mice inhibit contact sensitivity-effector T cells in vivo, and their interferon- γ production in vitro. *J Exp Med* 1996;184:2129-40.
- Szurszewski J. A migrating electric complex of the canine small intestine. *Am J Physiol* 1996;271:1757-63.

- Tafari A, Alferink J, Moller P, Hammerling GJ, Arnold B. T cell awareness of paternal alloantigens during pregnancy. *Science*. 1995;270:630-3.
- Takahashi N, Miner LL, Sora I, Ujike H, Revay RS, Kostic V, Jackson-Lewis V, Przedborski S, Uhl GR. VMAT2 knockout mice: heterozygotes display reduced amphetamine-conditioned reward, enhanced amphetamine locomotion, and enhanced MPTP toxicity. *Proc Natl Acad Sci USA* 1997;94:9938-43.
- Tanaka T, Mizumoto A, Mochiki E, Haga N, Suzuki H, Itoh Z. Relationship between intraduodenal 5-hydroxytryptamine release and interdigestive contractions in dog. *J Smooth Muscle Res* 2004;40:75-84.
- Taub DD, Conlon K, Lloyd AR, Oppenheim JJ, Kelvin DJ. Preferential migration of activated CD4+ and CD8+ T cells in response to MIP-1 alpha and MIP-1 beta. *Science* 1993;260(5106):355-8.
- Taylor MW, Feng GS. Relationship between interferon-gamma, indoleamine 2,3-dioxygenase, and tryptophan catabolism. *FASEB J* 1991;5:2516-22.
- Tecott LH, Sun LM, Akana SF, Strack AM, Lowenstein DH, Dallman MF, Julius D. Eating disorder and epilepsy in mice lacking 5-HT_{2c} serotonin receptors. *Nature* 1995;374(6522):542-6.
- Terron JA. Role of 5-HT₇ receptors in the long-lasting hypotensive response induced by 5-hydroxytryptamine in the rat. *Br J Pharmacol* 1997;121:563-71.
- Thellin O, Coumans B, Zorzi W, Igout A, Heinen E. Tolerance to the foeto-placental „graft“: ten ways to support a child for nine months. *Curr Opin Immunol*. 2000;12:731-7.
- Toda N, Tokuyama T, Seno S, Hirata F, Hayaishi O. Effects of 5-hydroxykynurenamine, a new serotonin metabolite, on isolated dog basilar arteries. *Proc Natl Acad Sci U S A* 1974;71(1):122-4.
- Trendelenburg P. Physiologische und pharmakologische Versuche über die Dünndarm Peristaltick. *Naunyn-Schmiedebergs Arch Exp Pathol Pharmacol* 1917;81:55-129.
- Turvill JL, Connor P, Farthing MJ. The inhibition of cholera toxin-induced 5-HT release by the 5-HT₃ receptor antagonist, granisetron, in the rat. *Br J Pharmacol* 2000;130(5):1031-6.
- Turvill JL, Mourad FH, Farthing MJ. Crucial role for 5-HT in cholera toxin but not Escherichia coli heat-labile enterotoxin-intestinal secretion in rats. *Gastroenterology* 1998;115(4):883-90.
- Tylicki L, Rutkowski B, Horl WH. Multifactorial determination of hypertensive nephroangiosclerosis. *Kidney Blood Press Res*. 2002;25(6):341-53.
- Tzeng DY, Deuel TF, Huang JS, Baehner RL. Platelet-derived growth factor promotes human peripheral monocyte activation. *Blood* 1985;66(1):179-83.
- Uchida K, Usami M, Bandow H, Harada I. Characteristics of substrates and inhibitors in binding to rat liver L-tryptophan 2,3-dioxygenase: a Fourier transform infrared and kinetic study. *Biochim Biophys Acta* 1992;1121:153-9.
- Uyttenhove C, Pilotte L, Theate I, Stroobant V, Colau D, Parmentier N, Boon T, Van den Eynde BJ. Evidence for a tumoral immune resistance mechanism based on tryptophan degradation by indoleamine 2,3-dioxygenase. *Nat Med* 2003;9:1269-74.
- van Hooff JA, Vijverberg HP. Selection of distinct conformational states of the 5-HT₃ receptor by full and partial agonists. *Br J Pharmacol* 1996;117(5):839-46.
- Vanhoutte P, Amery A, Birkenhager W, Breckenridge A, Buhler F, Distler A, Dormandy J, Doyle A, Frohlich E, Hansson L. Serotonergic mechanisms in hypertension. Focus on the effects of ketanserin. *Hypertension* 1988;11:111-33.

- Vantrappen G, Janssens J, Hellemans J, Ghoois Y. The interdigestive motor complex of normal subjects and patients with bacterial overgrowth of the small intestine. *J Clin Invest* 1977;59:1158-1166.
- van Zwieten PA, Blauw GJ, van Brummelen P. The role of 5-hydroxytryptamine and 5-hydroxytryptaminergic mechanisms in hypertension. *Br J Clin Pharmacol* 1990;30:69S-74S.
- Walsh HA, Daya S. Inhibition of hepatic tryptophan-2,3-dioxygenase: Superior potency of melatonin over serotonin. *J Pineal Res.* 1997;23:20-3.
- Walther DJ. Die Dualität des Serotonergen Systems. Freie Universität Berlin 2000.
- Walther DJ, Bader M. Serotonin synthesis in murine embryonic stem cells. *Mol Brain Res* 1999;68:55-63.
- Walther DJ, Peter JU, Bashammakh S, Hortnagel H, Voits M, Fink H, Bader M. Synthesis of serotonin by a second tryptophan hydroxylase isoform. *Science* 2003a;299:76.
- Walther DJ, Bader M. A unique central tryptophan hydroxylase isoform. *Biochem Pharmacol* 2003b;66:1673-80.
- Walther DJ, Peter JU, Winter S, Höltje M, Paulmann N, Grohmann M, Vowinkel J, Alamo-Bethencourt V, Wilhelm CS, Ahnert-Hilger G, Bader M. Serotonylation of small GTPases is a signal transduction pathway that triggers platelet α -granule release. *Cell* 2003c;115:851-62.
- Wang HY, Fiorica-Howells E, Pan H, Gershon MD, Friedman E. Myenteric ganglionic 5-hydroxytryptamine(1P) signal transmission is mediated via Go protein. *J Pharmacol Exp Ther* 1996;277(1):518-24.
- Wang YM, Gainetdinov RR, Fumagalli F, Xu F, Jones SR, Bock CB, Miller GW, Wightman RM, Caron MG. Knockout of the vesicular monoamine transporter 2 gene results in neonatal death and supersensitivity to cocaine and amphetamine. *Neuron* 1997;19:1285-96.
- Watts SW, Baez M, Webb RC. The 5-hydroxytryptamine_{2B} receptor and 5-HT receptor signal transduction in mesenteric arteries from deoxycorticosterone acetate-salt hypertensive rats. *J Pharmacol Exp Ther* 1996;277:1103-13.
- Weber LJ, Horita A. A study of 5-hydroxytryptamine formation from L-tryptophan in the brain and other tissues. *Biochem Pharmacol* 1965;14:1141-9.
- Weihe E, Schäfer MKH, Erickson JD, Eiden LE. Localization of vesicular monoamine transporter isoforms (VMAT 1 and VMAT2) to endocrine cells and neurons in rat. *J Mol Neurosci* 1994;5:149-64.
- Whalen EJ, Johnson AK, Lewis SJ. Functional evidence for the rapid desensitization of 5-HT₃ receptors on vagal afferents mediating the Bezold-Jarisch reflex. *Brain Res.* 2000 Aug 11;873(2):302-5.
- Wilhelm, CS. Der Einfluß des Serotonergen Systems auf das Immunsystem während der Schwangerschaft. Freie Universität Berlin 2001.
- Winkler T, Sharma HS, Stalberg E, Olsson Y, Dey PK. Impairment of blood-brain barrier function by serotonin induces desynchronization of spontaneous cerebral cortical activity: experimental observations in the anaesthetized rat. *Neuroscience* 1995;68(4):1097-104.
- Yamamoto T, Castell LM, Botella J, Powell H, Hall GM, Young A, Newsholme EA. Changes in the albumin binding of tryptophan during postoperative recovery: a possible link with central fatigue? *Brain Res Bull* 1997;43(1):43-6.
- Yamamoto K, Matsunaga S, Matsui M, Takeda N, Yamatodani A. Pica in mice as a new model for the study of emesis. *Methods Find Exp Clin Pharmacol* 2002a;24:135-8.

- Yamamoto K, Takeda N, Yamatodani A. Establishment of an animal model for radiation-induced vomiting in rats using pica. *J Radiat Res (Tokyo)* 2002b;43:135-41.
- Yavarone MS, Shuey DL, Tamir H, Sadler TW, Lauder JM. Serotonin and cardiac morphogenesis in the mouse embryo. *Teratology* 1993;47:573-84.
- Yilmaz MB, Saricam E, Biyikoglu SF, Guray Y, Guray U, Sasmaz H, Korkmaz S. Mean platelet volume and exercise stress test. *J Thromb Thrombolysis* 2004;17(2):115-20.
- Young MR, Kut JL, Coogan MP, Wright MA, Young ME, Matthews J. Stimulation of splenic T-lymphocyte function by endogenous serotonin and by low-dose exogenous serotonin. *Immunology*. 1993 Nov;80(3):395-400.
- Young MR, Matthews JP. Serotonin regulation of T-cell subpopulations and of macrophage accessory function. *Immunology*. 1995 Jan;84(1):148-52.
- Zeitz KP, Guy N, Malmberg AB, Dirajlal S, Martin WJ, Sun L, Bonhaus DW, Stucky CL, Julius D, Basbaum AI. The 5-HT₃ subtype of serotonin receptor contributes to nociceptive processing via a novel subset of myelinated and unmyelinated nociceptors. *J Neurosci* 2002;22:1010-9.
- Zhou M, Mellor AL.. Expanded cohorts of maternal CD8⁺ T-cells specific for paternal MHC class I accumulate during pregnancy. *J Reprod Immunol*. 1998;40:47-62.
- Zhou QJ, Quaife CJ, Palmiter RD. Targeted disruption of the tyrosin hydroxylase gene reveals that catecholamines are required for mouse fetal development. *Nature* 1995;374:640-3.
- Zill P, Baghai TC, Zwanzger P, Schule C, Eser D, Rupprecht R, Moller HJ, Bondy B, Ackenheil M. SNP and haplotype analysis of a novel tryptophan hydroxylase isoform (TPH2) gene provide evidence for association with major depression. *Mol Psychiatry* 2004;9(11):1030-6.
- Zoja C, Remuzzi G. Role of platelets in progressive glomerular diseases. *Pediatr Nephrol*. 1995;9:495-502.